

Comparative analysis of adipokinetic hormones and their receptors in Blattodea reveals novel patterns of gene evolution

Shixiong Jiang^{1,2}  | Heather G. Marco³  | Nina Scheich^{1,2} | Shulin He⁴ |
Zongqing Wang⁵ | Gerd Gäde³  | Dino P. McMahon^{1,2} 

¹Institute of Biology, Freie Universität Berlin, Berlin, Germany

²Department for Materials and Environment, BAM Federal Institute for Materials Research and Testing, Berlin, Germany

³Department of Biological Sciences, University of Cape Town, Rondebosch, South Africa

⁴College of Life Science, Chongqing Normal University, Chongqing, China

⁵College of Plant Protection, Southwest University, Chongqing, China

Correspondence

Dino P. McMahon, Institute of Biology, Freie Universität Berlin, Schwendenerstr. 1, Berlin 14195, Germany.
Email: dino-peter.mcmahon@bam.de

Funding information

China Scholarship Council, Grant/Award Number: 201906300033; National Research Foundation of South Africa, Grant/Award Numbers: 85768, 10924

Abstract

Adipokinetic hormone (AKH) is a neuropeptide produced in the insect *corpora cardiaca* that plays an essential role in mobilising carbohydrates and lipids from the fat body to the haemolymph. AKH acts by binding to a rhodopsin-like G protein-coupled receptor (GPCR), the adipokinetic hormone receptor (AKHR). In this study, we tackle AKH ligand and receptor gene evolution as well as the evolutionary origins of AKH gene paralogues from the order Blattodea (termites and cockroaches). Phylogenetic analyses of AKH precursor sequences point to an ancient AKH gene duplication event in the common ancestor of Blaberoidea, yielding a new group of putative decapeptides. In total, 16 different AKH peptides from 90 species were obtained. Two octapeptides and seven putatively novel decapeptides are predicted for the first time. AKH receptor sequences from 18 species, spanning solitary cockroaches and subsocial wood roaches as well as lower and higher termites, were subsequently acquired using classical molecular methods and *in silico* approaches employing transcriptomic data. Aligned AKHR open reading frames revealed 7 highly conserved transmembrane regions, a typical arrangement for GPCRs. Phylogenetic analyses based on AKHR sequences support accepted relationships among termite, subsocial (*Cryptocercus* spp.) and solitary cockroach lineages to a large extent, while putative post-translational modification sites do not greatly differ between solitary and subsocial roaches and social termites. Our study provides important information not only for AKH and AKHR functional research but also for further analyses interested in their development as potential candidates for biorational pest control agents against invasive termites and cockroaches.

KEYWORDS

adipokinetic hormone, adipokinetic hormone receptor, 'green' pesticide, neuropeptide, termite

INTRODUCTION

The adipokinetic hormones (AKHs) of insects are well-investigated neuropeptides, synthesised and released from the *corpora cardiaca* (CC). The biologically active peptides belong to the well known AKH/RPCH (red pigment-concentrating hormone) family of peptides,

of which more than 90 different mature AKHs have been identified by primary sequence or predicted from arthropods (Gäde et al., 1997; Gäde, 1997; Gäde, 2004; Gäde & Marco, 2009; Gäde et al., 2013; Gäde and Marco, 2022). In the biosynthesis and processing pathways in the CC, each AKH is derived from an individual preprohormone that is composed of a signal peptide and two potential peptides: the precursor

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Insect Molecular Biology* published by John Wiley & Sons Ltd on behalf of Royal Entomological Society.

of the bioactive AKH and a structurally-unrelated peptide known as the AKH precursor-associated peptide, to which no biological activity has been assigned to date. Following cleavage from the prohormone and post-translational modification, the mature AKH and the unrelated peptide are ready for release from storage/secretory granules (Van der Horst et al., 2001). Peptides of the AKH/RPCH family share typical features: at least 8 but no more than 10 amino acids in length, a blocked pyroGlu (pQ) N-terminus and a C-terminus blocked by amidation. The aromatic residues phenylalanine (Phe) or tyrosine (Tyr) occur at position 4 and tryptophan (Trp) at position 8 (Gäde, 1997; Gäde, 2004; Li et al., 2016). Investigated insect species may synthesise only one AKH peptide, whereas others produce from two to as many as five different AKH peptides in their CC (Gäde, 2009; Gäde et al., 2013; Marciniak et al., 2022; Marco & Gäde, 2019; Veenstra, 2014). However, the evolutionary origins and relationships between AKH gene paralogues in insects are poorly characterised.

It is a well-established fact that in insects, AKHs play a crucial role in the control of energy mobilisation in various physiological processes, such as in locomotory activity, starvation and stress response (Auerswald et al., 2005; Chino et al., 1989; Gäde, 2004; Gäde & Beenackers, 1977; Gäde & Marco, 2013; Goldsworthy et al., 1975; Holwerda et al., 1977; Isabel et al., 2005; Kodrík, 2008; Kodrík et al., 2000; Marco & Gäde, 2017; Robinson & Goldsworthy, 1977; Sajwan et al., 2015; Stone et al., 1976; Tang et al., 2020). These physiological functions are mediated by adipokinetic hormone receptors (AKHRs), specifically, GPCRs related to vertebrate gonadotropin-releasing hormone (GnRH) receptors. During energy-demanding circumstances, AKHs are released from the CC into the haemolymph to reach the AKHRs expressed on fat body cells. AKHR binding and activation initiates a signal transduction cascade that causes the activation of either glycogen phosphorylase or triacylglycerol lipase, which in turn activates specific cellular pathways to release trehalose or diacylglycerol and free fatty acids from the fat body (Gäde & Auerswald, 2003; Gálíková et al., 2015). In 1998, an insect GPCR was cloned from *D. melanogaster*, and 4 years later, it was confirmed to be an AKH receptor (Hauser et al., 1998; Park et al., 2002; Staubli et al., 2002). The AKHR from the silkworm *Bombyx mori* was also identified and described in 2002 (Staubli et al., 2002). Since then, AKHRs have been described and some have been functionally studied from a wide variety of insect species and orders, such as *Anopheles gambiae*, *Sarcophaga crassipalpis*, *Glossina morsitans morsitans* and *Bactrocera dorsalis* in Diptera (Belmont et al., 2006; Bil et al., 2016; Caers et al., 2016; Hou et al., 2017; Kaufmann & Brown, 2006), *Manduca sexta* (Lepidoptera) (Ziegler et al., 2011), *Tribolium castaneum* (Coleoptera) (Li et al., 2008), *Periplaneta americana* and *Blattella germanica* (Blattodea) (Hansen et al., 2006; Huang et al., 2012; Wicher et al., 2006), *Schistocerca gregaria* and *Gryllus bimaculatus* (Orthoptera) (Jackson et al., 2019; Konuma et al., 2012), *Acyrtosiphon pisum* and *Rhodnius prolixus* (Hemiptera) (Alves-Bezerra et al., 2016; Jedlička et al., 2012; Zandawala et al., 2015) and *Carausius morosus* in Phasmatodea (Birgul Iyison et al., 2020).

Termites are a relatively small group of social insects consisting of more than 3000 species (Krishna et al., 2013), which were traditionally classified as the order Isoptera. More recent research demonstrates

that termites appear as the sister clade to the *Cryptocercus* lineage of wood roaches, nesting within cockroaches and, thus, belong to the order Blattodea (Bourguignon et al., 2014; Evangelista et al., 2019; Inward et al., 2007; Klass et al., 2008; Krishna et al., 2013). Blattodea together with Mantodea (mantids) comprise the superorder Dictyoptera, which share a common recent ancestor (Ware et al., 2008).

The most recent study on the structural diversity of AKHs in 13 termite species (Jedličková et al., 2016), used molecular methods (seven species) and *in silico* bioinformatic searches (six species) from five families of Isoptera for phylogenetic analyses between termites and their closest cockroach relatives. These analyses were, however, carried out without sequence data of representatives from all termite families, and in the absence of data from wood roaches (*Cryptocercus*) since the AKH prohormone sequence was not known at the time. In fact, at that time, only the mature AKH sequences were known from three termite species (Liebrich et al., 1995) and over 30 cockroach species (Gäde, 2009; Roth et al., 2009). Despite the short peptide sequences of AKHs, these have been useful in some insect orders, including cockroaches, to investigate phylogenetic trends and evolutionary relationships between insect lineages (Gäde, 1989; Gäde & Marco, 2005; Marco et al., 2020; Roth et al., 2009). However, the occurrence of multiple AKH genes, including within Blattodea, can hamper the interpretation of phylogenetic approaches (Bläser et al., 2020; Bläser & Predel, 2020), particularly when dealing with incomplete or missing data and insufficient taxon sampling. For example, while termites possess just one AKH gene, several cockroach species have 2 AKH peptides (Gäde, 2009; Gäde & Rinehart, 1990; Jedličková et al., 2016).

In contrast to the peptide ligands, only a few AKH receptor sequences are known from Blattodea to date. The AKHR of *P. americana* was the first to be identified from a hemimetabolous insect (Hansen et al., 2006). The AKHR of *Blattella germanica* was cloned in 2012 (Huang et al., 2012), while there are two putative AKHR sequences from termites available on NCBI: *Zootermopsis nevadensis* (XM_022082426.1) and *Cryptotermes secundus* (XM_033755069.1). There is, thus, a paucity of information regarding AKHR sequences of Blattodea. Receptor sequences are fundamentally important for further molecular and pharmacological characterisation of receptor-ligand systems, especially in the identification of potential methods to combat pest insect species.

In addition to representing a number of global pest species, cockroaches and termites play a crucial role in ecosystem services, including in the decomposition of deadwood (Bell et al., 2007; Bignell & Eggleton, 2000). Currently, researchers are considering the use of peptide mimetics (as so-called 'green' insecticides), based on the interaction of neuropeptide ligands with receptors, to negatively influence the physiology and/or behaviour of specific pest insects without harming beneficial insects (Altstein & Nässel, 2010; Gäde et al., 2017; Marco et al., 2018). RNAi represents an alternative potential strategy for pest control, enabling downregulation of specific gene targets via the host's RNAi pathway, whereby dsRNA is converted to small interfering RNA (siRNA), in turn, binding to the RNA-induced silencing complex (RISC), which then efficiently locates and eliminates target mRNA and reducing its translation into protein (Arakane et al., 2008;

TABLE 1 AKH peptides of termites and cockroaches. The taxonomic classification is per Evangelista et al. (2019) and Hellemans et al. (2022). Adipokinetic hormone nomenclature: Bladi-HrTH, *Blaberus discoidalis* hypertrehalosemic hormone; Emppe-AKH, *Empusa pennata* AKH; Manto-CC, Mantophasmatodea CC; Micvi-CC, *Microhodotermes viator* CC; Peram-CAH, *Periplaneta americana* cardioacceleratory hormone; Polae-HrTH, *Polyphaga aegyptiaca* hypertrehalosemic hormone; Pyrap-AKH, *Pyrrhocoris apterus* AKH; Tenmo-HrTH, *Tenebrio molitor* hypertrehalosemic hormone. For peptide source details; see Table S1.

Taxonomy sub-family	Species	AKH name	AKH sequence	References		
Blaberodea	<i>Blattellinae</i>	<i>Lobopterella dimidiatipes</i>	Bladi-HrTH	pQVNFSPGWGTa This study (source: 1KITE)		
		<i>Loboptera decipiens</i>		Roth et al. (2009)		
	<i>Nyctiborinae</i>	<i>Symploce</i> sp., <i>Asiablatta kyotensis</i> , <i>Ischnoptera deropeltiformis</i> , <i>Anallacta methanoides</i>	Bladi-HrTH putative novel 1	pQVNFSPGWGTa pQLNFSPGWGVA	This study (source: 1KITE)	
		<i>Loboptera decipiens</i>			This study (source: 1KITE)	
		<i>Blattella germanica</i>	Bladi-HrTH putative novel 2	pQVNFSPGWGTa pQLNFSPGWGPa	Gäde and Rinehart (1990); Veenstra and Camps (1990); This study (source: Sanger sequencing)	
		<i>Nyctibora</i> sp.	Bladi-HrTH putative novel 3	pQVNFSPGWGTa pQVNFSPGWGVA	This study (source: 1KITE)	
		<i>Blaberidae</i>	<i>Aptera fusca</i> , <i>Archimandrita tessellate</i> , <i>Bantua robusta</i> , <i>Blaptica dubia</i> , <i>Blepharodera discoidalis</i> , <i>Diploptera punctata</i> , <i>Elliptorhina</i> sp., <i>Eulaberus distanti</i> , <i>Gyna caffrorum</i> , <i>Lucihormetica grossei</i> , <i>Panaesthia</i> sp., <i>Panchlora viridis</i> , <i>Princisia vanwaerebeki</i>	Bladi-HrTH	pQVNFSPGWGTa	Roth et al. (2009)
			<i>Blaberus discoidalis</i>			Hayes et al. (1986)
			<i>Gromphadorhina portentosa</i> , <i>Leucophaea maderae</i>			Gäde and Rinehart (1990)
			<i>Nauphoeta cinerea</i>			Gäde and Rinehart (1986)
<i>Panchlora nivea</i>	Bladi-HrTH putative novel 4		pQVNFSPGWGTa pQLNFSPGWGTa	This study (source: 1KITE)		
<i>Blaberus atropos</i>	Bladi-HrTH putative novel 5		pQVNFSPGWGTa pQLNFSPGWGfA	This study (source: 1KITE)		
<i>Princisia vanwaerebeki</i> , <i>Diploptera punctata</i>	Bladi-HrTH		pQVNFSPGWGTa	This study (source: 1KITE)		
<i>Pseudophyllodromiinae</i>	<i>Balta vilis</i> , <i>Ellipsoidion</i> sp., <i>Supella longipalpa</i> , <i>Euthlastoblatta diaphana</i>	Bladi-HrTH putative novel 1	pQVNFSPGWGTa pQLNFSPGWGVA	This study (source: 1KITE)		
	<i>Sundablatta sexpunctata</i>	Bladi-HrTH putative novel 6	pQVNFSPGWGTa pQINFSPGWGVA	This study (source: 1KITE)		
	<i>Cariblatta</i> sp.	putative novel 6 putative novel 7	pQINFSPGWGVA pQVNFSPGWGAA	This study (source: 1KITE)		
	<i>Ectobiinae</i>	<i>Ectobius sylvestris</i>	Bladi-HrTH putative novel 1	pQVNFSPGWGTa pQLNFSPGWGVA	This study (source: 1KITE)	

(Continues)

TABLE 1 (Continued)

Taxonomy sub-family	Species	AKH name	AKH sequence	References		
Solumblattodea	Isoptera	Termitidae		<i>Odontotermes formosanus</i> , <i>Macrotermes natalensis</i> , <i>Nasutitermes takasagoensis</i> , <i>Termes hospes</i>	Jedličková et al. (2016)	
				<i>Trinervitermes trinervoides</i>	Liebrich et al. (1995)	
				<i>Bulbitermes</i> sp., <i>Dicuspiditermes</i> sp., <i>Globitermes</i> sp., <i>Indotermes</i> sp., <i>Macrotermes subhyalinus</i> , <i>Pericapritermes</i> sp., <i>Promicrotermes</i> sp.	This study (source: He et al., 2021)	
	Rhinotermitidae	Peram-CAH-I	pQVNFSPNWa	<i>Coptotermes formosanus</i> , <i>Coptotermes gestroi</i> , <i>Prohinotermes simplex</i> , <i>Reticulitermes flavipes</i> , <i>Reticulitermes speratus</i>	Jedličková et al. (2016)	
				<i>Reticulitermes grassei</i> , <i>Prohinotermes inopinatus</i>	This study (source: He et al., 2021)	
				<i>Neotermes castaneus</i>	Jedlickova et al. (2016)	
	Kalotermitidae			<i>Cryptotermes</i> sp., <i>Cryptotermes domesticus</i> , <i>Incisitermes marginipennis</i>	This study (source: I. marginipennis: 1KITE; C. domesticus: He et al., 2021, <i>Cryptotermes</i> sp. Sanger sequencing)	
				<i>Kalotermes flavicollis</i>	This study (source: Sanger sequencing)	
	Blattodea	Archotermopsidae	Manto-CC	pQVNFSPGWa	<i>Kalotermes flavicollis</i>	This study (source: Sanger sequencing)
					<i>Hodotermodopsis sjostedti</i> , <i>Zootermopsis nevadensis</i>	Jedličková et al. (2016)
Hodotermitidae		Micvi-CC		<i>Microhodotermes viator</i>	Liebrich et al. (1995)	
				<i>Mastotermes darwiniensis</i>	Liebrich et al. (1995); Jedličková et al. (2016)	
Cryptocercidae		Tenmo-HfTH	pQLNFSPNWa	<i>Cryptocercus darwini</i>	Roth et al. (2009)	
				<i>Cryptocercus punctulatus</i>	Gäde et al. (1997)	
				<i>Cryptocercus pudacoensis</i> , <i>Cryptocercus meridianus</i>	This study (source: He et al., 2021)	
Lamproblattidae		Peram-CAH-I	pQVNFSPNWa	<i>Lamproblatta albipalpus</i>	This study (source: 1KITE)	
				<i>Blatta orientalis</i>	Gäde and Rinehart (1990)	
Blattidae				<i>Brinckia hanstroemi</i> , <i>Celatoblatta</i> sp., <i>Deropeltis erythrocephala</i> , <i>Eurycotis floridana</i> , <i>Neostylopyga rhombifolia</i> , <i>Pseudoderopeltis foveolate</i> , <i>Shelfordella lateralis</i>	Predel and Gäde (2005)	
	<i>Periplaneta americana</i>			Peram-CAH-I Peram-CAH-II	pQVNFSPNWa pQLTFTPNWa	Scarborough et al. (1984); Witten et al. (1984); Zeng et al. (2020)
	<i>Shelfordella lateralis</i> , <i>Eurycotis floridana</i> , <i>Methana parva</i>					This study (source: 1KITE)

TABLE 1 (Continued)

Taxonomy sub-family	Species	AKH name	AKH sequence	References
Tryonicidae	<i>Tryonicus parvus</i>	Emppe-AKH	pQINFTPNWa	This study (source: 1KITE)
		Polae-HrTH	pQITFTPNWa	
Corydiidae	<i>Polyphaga aegyptiaca</i>	Tenmo-HrTH	pQLNFSNWA	Gäde and Kellner (1992)
		Polae-HrTH	pQITFTPNWa	
	<i>Ergaula capucina</i>			König et al. (2005)
	<i>Therea petiveriana</i>	Tenmo-HrTH	pQLNFSNWA	Gäde et al. (1997)
	<i>Therea bernhardtii</i>			This study (source: 1KITE)
Nocticolidae	<i>Eucoydia yasumatsui</i>	Pyrap-AKH	pQLNFTPNWa	This study (source: 1KITE)
	<i>Tivia</i> sp.	Peram-CAH-I	pQVNFSPNWA	This study (source: 1KITE)
	<i>Nocticola</i> sp.	Peram-CAH-I	pQVNFSPNWA	This study (source: 1KITE)

Burand & Hunter, 2013; Kapan et al., 2012; Lee et al., 2011; Park et al., 2014; Rewitz et al., 2009; Yu et al., 2023). For such applications, it is imperative to know peptide and receptor sequences and structures in as many pest species as possible, as well as in non-pest species to evaluate the potential collateral damage or success of sequence-based green insecticides.

With the current study, we employ bioinformatic and molecular approaches to characterise AKH precursor sequences and cognate AKHR sequences from a wide diversity of termite and cockroach species, including wood roaches. We combine recently sequenced transcriptomic data with publicly available Sequence Read Archive (SRA) data and classical molecular approaches to conduct a comprehensive sequence analysis of AKH and AKHR in termites and their nearest cockroach relatives. Our aims are to investigate the evolution of AKH peptides and their receptors in Blattodea, as well as resolve the evolutionary origins and relationships of AKH gene duplications in this group. Additionally, we enhance termite AKH and AKHR datasets for comparative analysis, carry out an assessment of patterns of post-translational modification during the evolution of termite sociality and establish a data framework for research aiming to exploit AKH and AKHRs as potential targets for 'green' pesticide development against invasive termites and cockroaches, or for possible RNAi interventions.

RESULTS

AKHs in termites and cockroaches

From the complete list of AKH mature peptides derived from 90 species (Table 1), we were able to obtain sequence information for 85 partial or complete AKH precursors from a total of 62 species spanning diverse families across the order Blattodea. The information was collated from pre-existing (published) data as well as from transcriptome and SRA data and bioinformatic searches of databases, as well as via Sanger sequencing in selected cases to confirm the *in silico* findings. The aligned AKH preprohormone sequences of these species demonstrate that both signal peptides and AKH-associated peptides are conserved at the amino acid level (Figure 1). Precursor sequences in cockroaches, especially for those species in the same family (Table 1), show high levels of sequence conservation, such as only two substitutions between AKH1 of *Shelfordella lateralis* and AKH1 of *P. americana*, or five substitutions between AKH2 of *A. kyotensis* and AKH2 of *S. sexpunctata*.

The mature peptide for each species was classified based on amino acid sequence identity to known AKHs and categorised based on taxonomy. Our new sequence data are compared alongside previously published findings in Table 1. A total of 16 AKH primary sequences from 90 species were identified. Table 1 shows that all predicted bioactive mature peptides consist of 8 or 10 amino acids. In Blaberoidea, only decapeptides were found, with Bladi-HrTH being the most frequently detected peptide (Table 1, Figure 2), which was

found in all species except *Cariblatta* sp. (where two putative novel decapeptides are identified). Further to this, we identified seven putative novel decapeptides in Blaberoidea via *in silico* characterisation.

The presence of the putative novel 2 type in *Blattella germanica* was confirmed via Sanger sequencing. By contrast, species in Solumblatodea (Blattoidea + Corydioidea) contained only octapeptides. The




FIGURE 1 Multiple sequence alignment of 85 termite and cockroach Adipokinetic hormone (AKH) preprohormones from 62 species. The sequence logo beneath depicts the consensus sequence and the solid line under the sequence logo identifies the AKH bioactive peptide. The sequence before the bioactive peptide is referred to as the signal peptide (dashed line), while the sequence following is the AKH-associated peptide. Putative amidation and dibasic cleavage sites are indicated in lowercase based on known homologous precursors in other species. Available sequence accession codes are listed in Data S1.

most frequently detected AKH in Solumblattodea was Peram-CAH-I (Figure 2), which occurs not only in most termite families such as Termitidae, Rhinotermitidae, Mastotermitidae and Kalotermitidae but also in the cockroach family Lamproblattidae, Blattidae, Corydiidae and Nocticolidae. Two AKH octapeptides were recorded in some cockroaches: Peram-CAH-I and Peram-CAH-II in Blattidae, Tenmo-HrTH and Polae-HrTH in Corydiidae (*Polyphaga aegyptiaca*) and Empe-AKH + Polae-HrTH in Tryonicidae (*Tryonicus parvus*). Micvi-CC is exclusively found in the Hodotermitidae (Table 1). The sister group to the termites, the wood roaches (genus *Cryptocercus*) possess a Tenmo-HrTH peptide, which differs only in the occurrence of leucine at position 2 in comparison to valine in Peram-CAH-I. Interestingly, a unique AKH peptide, Manto-CC, in *K. flavicollis* (Kalotermitidae) was identified using Sanger sequencing in this study and further confirmed and validated by mass spectrometry (Marco et al., 2022) for the first time in cockroaches and termites, with a structure that differs from Peram-CAH-I at position 7 by a glycine/asparagine substitution. The other unique octapeptide, Pypap-AKH in *Eucorydia yasumatsui* (Corydiidae), was found for the first time in Blattodea, with a structure that differs from Tenmo-HrTH at position 5 by a serine/threonine substitution (Table 1).

Phylogenetic analysis of AKH in termites and cockroaches

Phylogenetic trees obtained from two different methods, RAxML and MrBayes, rooted between the major Blattodea clades: Blaberoidea and Solumblattodea (Blattoidea + Corydioidea) (Figure 3, Figure S1) support an AKH duplication event in the ancestor of Blaberoidea; leading to a

Bladi-HrTH decapeptide clade (except *Cariblatta* sp., which possesses a novel decapeptide, although this still groups with high support in the Bladi-HrTH clade) and a clade of novel putative decapeptides. Node support for this hypothetical duplication event is high when using nucleotide data and slightly less so when employing amino acid sequences. Termite AKHs nested paraphyletically within Solumblattodea sequences, with limited bootstrap support at deeper nodes in this clade for both nucleotide and amino acid trees. In terms of sequence evolution, termitid plus *Reticulitermes* and *Coptotermes* AKHs appear to diverge from the remaining lower termite and cockroach sequences.

The relationships between Corydiidae + Nocticolidae (Corydioidea), Blattidae and other lineages within Solumblattodea are generally equivocal. The nucleotide phylogeny suggests that a parallel AKH duplication is likely to have taken place in Blattidae, although this receives limited statistical support in both ML and BI approaches and so should be interpreted with a degree of caution.

AKHRs in termites and cockroaches

A total of 18 new AKH receptor sequences were retrieved bioinformatically and/or sequenced using a classical molecular approach. We aligned the sequences for these 18 species and for the other 4 available species to obtain a better understanding of amino acid variation among blattid and termite AKHR sequences (Figure 4). The ORF (open reading frame) of all termite and cockroach AKHRs contains sequences of between 405 and 467 amino acid residues in length. Except for *Cryptotermes* sp. and *C. secundus*, the shortest amino acid sequences belong to the Blattellinae subfamily: *B. germanica*, *Symploce* sp., *I. deropeltiformis*, *A. kytensis* and




FIGURE 2 Simplified phylogenetic scheme of Blattodea modified from Evangelista et al. (2019) and Hellemans et al. (2022). The sequence logo represents the degree of conservation of amino acids in Adipokinetic hormone (AKH) neuropeptides in Blaberoidea and Solumblattodea. N: number of AKH neuropeptide sequences. Mantodea AKH (Empe-AKH; Gäde and Marco (2017)) is used as an outgroup.




FIGURE 3 ML phylogenetic trees constructed from AKH preprohormones from termites and cockroaches. Trees were reconstructed from alignments of the whole ORF sequence, using amino acids (a) or nucleotides (b). Numbers at each node represent bootstrap support values (in percent), with only values above 50 being shown. The putative novel 7 decapeptide of *Cariblatta* sp. AKH2* grouped with Bladi-HrTH. For comparison, equivalent BI trees are shown in Figure S1.

P. coulöniana. The results derived from the aligned sequences demonstrate that termite and cockroach AKH receptors share highly conserved sequence elements, particularly in the seven transmembrane domains (TMs), which corresponds to the typical arrangement found in G-protein-coupled receptors (TM1-TM7 highlighted in Figure 4). Sequence conservation is especially high in TM2, TM6 and TM7, although moderate levels of sequence conservation could be detected across most of the ORF, except close to the N- and particularly the C-termini.

We also compared pairwise identity and similarity at the amino acid level of the AKHRs identified to date of the full 22 Blattodea species (Figure 5). Identities and similarities range from 64.0% to 99.6% and from 68.4% to 99.6%, respectively. Notable levels of identity and similarity include comparisons between solitary cockroaches (Blattidae + Blattellinae) and wood-feeding cockroaches (*Cryptocercus*): 71.6%–81.5%, whereas between solitary

cockroaches and higher termites (Termitidae), identity and similarity values range from 64.0% to 76.0%. Termitidae and *Cryptocercus* are separated by intermediate levels (71.3%–79.0%) of amino acid identity and similarity. AKHR identity and similarity within kalotermitids and rhinotermitids range from 93.6% to 99.3% and 92.2% to 99.6%, respectively, whereas between these families, values range from 81.1% to 88.2%. All other pairwise comparisons are shown in Figure 5.

Phylogenetic analysis of AKHRs in termites and cockroaches

Phylogenetic reconstructions of 22 AKHR sequences are depicted in Figure 6. The evolution of AKHR largely reflects the accepted view of termites and cockroach lineage diversification (Figure 6).



FIGURE 4 Comparative amino acid sequence alignment of termite and cockroach AKHR sequences. Intracellular (ICL) and extracellular (ECL) loops, N-terminus and C-terminus are labelled above the alignment. Transmembrane domains 1–7 are highlighted in grey. Blue and red lines under the alignment show putatively conserved myristoylation and glycosylation sites, respectively.

The topologies of the ML and BI phylogenetic trees are identical. Five AKHRs from Blaberidae cluster together as a monophyletic group. *B. orientalis* and *P. americana* in the Blattidae family are closely clustered with high support and *Cryptocercus* occurs as the nearest lineage to the termites. Within the termites, *Mastotermes*, *Zootermopsis* and *Kalotermitidae* occur in expected positions in the phylogenies, as do the AKHR sequences belonging to the Neoisoptera (Rhinotermitidae + Termitidae). The rhinotermitids *Reticulitermes* and *Coptotermes* are paraphyletic with respect to Termitidae (instead of appearing as a monophyletic clade). But overall the evolution of AKHR in termites and cockroaches mostly reflects the accepted view of termite lineage diversification and the majority of ancestral nodes receive unequivocal support.

Predicted AKHR post-translational modifications

Post-translational modifications (PTMs) and their potential interactions contribute significantly to the biological functions of proteins, and GPCRs are an important class of proteins that are regulated by PTMs. To investigate the putative PTMs of AKHRs in termites, three modifications, glycosylation, myristoylation and phosphorylation were explored. The results reveal that all 22 species contain putative glycosylation sites in at least two domains (in the ECL1 and C-terminus, Figure 7). Among them, cockroaches in the Blattidae family (*B. orientalis* and *P. americana*) are the most widely distributed glycosylation sites, being found in the following domains: N-terminus, TM1, ECL1, ECL2, ICL3 and the C-terminus.

Two subsocial cockroaches in the genus *Cryptocercus* (*C. pudacoensis* and *C. meridianus*) and in Archotermopsidae (*Z. nevadensis*) have five glycosylation domains in the TM1, ECL1, ECL2, ICL3 and C-terminus. By contrast, *C. formosanus* contains two (ECL1 and C-terminus) and *Indotermes* sp. only three (ECL1, ECL2 and C-terminus). All 22 species contain one glycosylation site in the ECL1. In addition, all cockroach species, except *Symptloce* sp. and *Z. nevadensis* contain one glycosylation site in ICL3 (Figure 7a).

In terms of myristoylation sites, all AKHR sequences have at least two putative domains in TM3 and the C-terminus. In addition, *Cryptotermes* sp. and *C. secundus* (Kalotermitidae) contain an other site localised in the N-terminus, while *C. pudacoensis*, *C. meridianus* and *Z. nevadensis* (*Cryptocercus* + Archotermopsidae) contain a further myristoylation site in the TM5 (Figure 7b).

Three putative phosphorylation sites (serine, threonine and Tyr residues) were quantified in the intracellular domains. In ICL1, two sites in *B. orientalis* and *P. americana* (Blattidae), *N. castaneus*,




FIGURE 5 Pairwise comparison (%) of amino acid identity (lower triangular) and similarity (upper triangular) from 22 AKHRs of termites and cockroaches.




FIGURE 6 Phylogenetic reconstruction of adipokinetic hormone receptors (AKHRs) in termites and cockroaches using amino acids (a) or nucleotides (b). Numbers at each node represent bootstrap support values/posterior probabilities (ML/BI respectively). Different colours indicate traditional classifications of termites and cockroaches (Evangelista et al., 2019). Bootstrap support/posterior probability values above 50/0.8 are displayed.




FIGURE 7 Distribution of putative glycosylation (a) and myristoylation sites (b) from AKHRs in 22 blattodean species (13 termite and 9 cockroach species). C_ter, C-terminus; ECL, Extracellular loop; ICL, Intracellular loop; N_ter, N-terminus.




FIGURE 8 Phosphorylation features of cockroach and termite AKHR intracellular domains. (a) distribution of phosphorylation sites (Ser, Thr and Tyr); (b) varieties of putative protein kinase phosphorylation sites and (c) their distribution. C_ter, C-terminus; ICL, Intracellular loop; InsR, Insulin receptor; PKA, Protein kinase A; PKB, Protein kinase B; PKC, Protein kinase C; UNSP, Unspecified kinase.

C. secundus, *Cryptotermes* sp. and *K. flavicollis* (Kalotermitidae) were detected. One site was found in *R. grassei* and *R. flavipes* (Rhinotermitidae) and *Globitermes* sp. (Termitidae). In ICL2, two sites were found for *A. kyotensis* and *P. coulouana*, and one site was detected in *B. orientalis*, *P. americana*, *B. germanica*, *I. deropeltiformis*, *C. pudacoensis*, *C. meridianus*, *M. darwiniensis* and *Z. nevadensis*. In ICL3, four or five sites in all roach species were recovered, as well as in *Z. nevadensis* and all kalotermitid species, while other termites possess six sites. Regarding the C-terminus, *P. simplex* contains the most sites, followed by *P. inopinatus*. With respect to putative kinase phosphorylation sites, *M. darwiniensis* has the highest number in the ICL and in the C-terminus (Figure 8a, b).

Further analysis of specific kinases indicates that putative PKA and UNSP sites are distributed among three ICL domains and the C-terminus. Most protein kinase phosphorylation sites were found in the C-terminus, while PKB, PKC and InsR were restricted to two or three ICL domains. Termite species possess up to 4 additional protein kinase phosphorylation sites (except *Z. nevadensis*) in the ICL3 compared with wood and solitary cockroaches (Figure 8c, Table S3), while they have lost 1 such site in the ICL 2 (except *M. darwiniensis* and *Z. nevadensis*) compared to wood and solitary cockroaches. Otherwise, PKA, PKC and UNSP sites were regularly predicted to occur in both the ICL3 and the C-terminus of all 22 species, indicating their potentially conserved role in intracellular signalling among blattodean AKHRs.

DISCUSSION

We present 16 structurally different mature AKH peptides from 90 species distributed across a wide range of cockroach and termite families. Evolutionary trees based on phylogenetic analyses using AKH precursor sequences reveal a new family of putative decapeptides in Blaberoidea. Phylogenetic trees indicate that an ancient duplication of AKH genes occurred in the ancestor of this group, yielding the previously described Bladi-HrTH on the one hand, and a group of novel decapeptides on the other. The decapeptide AKH, Bladi-HrTH, was until now the only decapeptide to have been reported from Blattodea (Gäde & Rinehart Jr, 1986; Hayes et al., 1986). Our study identifies seven putative novel decapeptides, significantly increasing the known diversity of decapeptides in Blattodea. The remaining AKHRs of Solumblattodea are found to be solely octapeptides in nature, with Peram-CAH-I (pQVNFSPNWamide) accounting for the largest proportion in this clade. Further species with two AKH peptides were also identified in Blattidae, Tryonicidae and Corydiidae. Phylogenies suggest that at least one parallel duplication event also took place at some point during the evolution of the Solumblattodea, although statistical support is low and inconsistently retrieved between nucleotide and amino acid-based analyses.

The presence of more than one mature AKH in a single species may indicate that each AKH plays a role in different regulatory processes (Bártů et al., 2010; Goldsworthy et al., 1997; Kaufmann & Brown, 2008). Indeed, the retention of both AKH genes in many extant

species found across Blaberoidea suggests that both peptides serve an important adaptive purpose, potentially serving different physiological functions in these insects. While the functional significance of AKH peptide duplicates in Blaberoidea, and potentially also Solumblattodea, remain under investigation, we hypothesise that these may be related to the capacity of cockroaches to live in diverse and challenging niches, where adaptive responses to diverse pathogens, toxins or fluctuating environments may have necessitated important alterations to AKH-mediated energy, stress or starvation responses.

Multiple AKHRs have been reported in other insects, such as one decapeptide and two octapeptides in the migratory locust, *Locusta migratoria* (Oudejans et al., 1991; Siegert et al., 1985; Stone et al., 1976). One octapeptide and two decapeptides are also described from the African frog hopper, *Ptyelus flavescens* (Gäde et al., 2017), while certain grasshoppers have three octapeptides (Gäde, 2006) and even five mature peptides (octa-, nona- and decapeptides) have been isolated and sequenced from two species of the lepidopteran genus *Hippotion* (Gäde et al., 2013). We also confirm the presence of Manto-CC in termites, being previously only recorded from the order Mantophasmatodea (Gäde et al., 2005) and also recently verified by mass spectrometry in the termite *K. flavicollis* (Marco et al., 2022). Another surprising discovery from the current work is that Pyrap-AKH is also present in cockroaches, where previously it had been found in Hemiptera (Kodrík et al., 2000; Gäde and Marco, 2022), Orthoptera (Caelifera) (Gäde, 2006) and in some beetle species (Gäde et al., 2019).

A study by Evangelista et al. (2019) provides a recent phylogenomic analysis of Blattodea, while a further study analysing 17 neuro-peptide precursor sequences (AKH precursor sequences were not included) revealed consistent topologies (Bläser et al., 2020). Our evolutionary trees based on AKHR genes were broadly in agreement with accepted topologies. An interesting question to ponder here is how to interpret divergent versus conserved patterns of ligand and receptor gene evolution, respectively. Multiple ligand binding may be permitted by the availability of large AKHR binding pockets which could facilitate promiscuous receptor-ligand interactions (Marchal et al., 2018; Marco & Gäde, 2019; Rios et al., 2001; Schwartz, 1994; Stank et al., 2016; Venkatakrishnan et al., 2013; Zhu et al., 2009). A recent study by Jackson et al. (2019) determined the structure of three AKHRs in the desert locust, *Schistocerca gregaria* using NMR (Nuclear magnetic resonance) techniques, finding that they interact with the same receptor residues despite having varying chain lengths and sequences. Such flexibility could depend on the receptor's ligand affinity properties, the intrinsic stability of receptor states and long-range allosteric coupling dynamics between the binding pocket and receptor regions residing on the cytoplasmic side (Chen et al., 2020; Cong et al., 2022; Möller et al., 2001; Xia et al., 2022).

We next chose to examine predicted post-translational modification signatures in AKHR. We explored this by characterising putative glycosylation, myristoylation and phosphorylation sites in corresponding TM regions, as well as intra- and extracellular domains of the AKHR amino acid sequence. A principal aim of this analysis was to explore whether patterns of post-translational modification differed between

social termites and their solitary cockroach relatives. There are some patterns of note, such as 1 fewer and up to 4 additional protein kinase phosphorylation sites in the ICL2 and ICL3 of termites, compared with cockroaches. Neoisopteran sequences also appear to possess an additional predicted PKA site and an additional UNSP site in ICL3 compared with other termites and cockroaches. Kalotermitids (dry wood termites) appear to have fewer putative sites of modifications in general, which may reflect a truncated C-terminus in these species. A study by Yang et al. (2018) implied potential differences in post-translational modification of AKHRs between solitary and social bees. Aside from the minor changes outlined above, our study did not detect obvious phosphorylation site pattern differences between social termites and subsocial or solitary cockroaches. This is perhaps surprising, given the important role of post-translational modification in diverse functions including protein interaction and stability, signalling, β -arrestin recruitment and receptor trafficking (Chou, 2020; Withers & Dong, 2017).

Bioactive neuropeptides and their GPCRs in insects are under investigation for their potential use as a more environmentally friendly alternative to conventional pesticides. This concept of a 'green insecticide' is based on the potentially disruptive consequences of interference at the level of neuropeptide-cognate receptor binding, as discussed elsewhere (Gäde et al., 2017; Gäde & Goldsworthy, 2010; Whetstone & Hammock, 2007). One of the first steps towards identifying such a lead for the chemical development of an AKH peptide mimetic with which to disrupt normal endocrine signalling in crucial metabolic pathways in so-called pest insects is to ascertain the complement of AKHs in insects and to establish whether cross-activity of a ligand is possible also in other insects.

In the case of Blattodea, cockroaches and termites are ecologically important detritivores, especially in the subtropical and tropical regions where they recycle wood and plant matter and represent a major driver of carbon cycles (Bignell, 2019; Ulyshen, 2016). Nevertheless, approximately 10% of termite species are considered severe pest insects where they infect and feed on dry timber in man-made structures and are very costly to combat worldwide (Khan & Ahmad, 2018; Su & Scheffrahn, 2000). Furthermore, less than 1% of cockroach species are recognised as omnivorous pest insects, when they come into close contact with human habitation (Cochran, 1999), presenting potentially serious health threats (Rosenstreich et al., 1997). Table 1 reveals that the AKH octapeptide, Peram-CAH-I, is not pest-insect specific, being shared by 36 cockroach and termite species. In fact, Peram-CAH-I is prevalent also in other insect orders, such as Hemiptera (Gäde and Marco, 2022), Coleoptera (Gäde et al., 2019) and Archaeognatha (Marco et al., 2014) and is, thus, not a suitable lead compound as cross-activity is likely to occur across a wide spectrum of insects. The remaining 6 octapeptide AKHs so far detected in the Blattodea are also not unique (Gäde, 2009), except for Polae-HrTH, which is synthesised in the CC of only 2 cockroach species (Table 1). However, our current study has uncovered a potential wealth in decapeptide AKHs, characterised by both *in silico* approaches as well as Sanger sequencing in the case of *B. germanica*: Table 1 lists a total of 8 decapeptides, 7 of which are novel structures, and all are present only in the Blaberoidea. These AKHs may be sufficiently different to warrant

such a decapeptide as a lead peptide. The strategy is clear and partially shown for the locust pair of AKHs and the cognate AKH receptor where a non-peptide (mimetic) has already been found to act as a competitive substance on the receptor (Jackson et al., 2022), and a case study in which a mimetic could bind the locust AKH receptor but not that of the honeybee (Abdulganiyyu et al., 2020).

Co-application of AKH with pathogens could yet further enhance these effects by interfering with the immune response in some insect species (Adamo et al., 2008; Gautam et al., 2020; Ibrahim et al., 2017; Ibrahim et al., 2018), potentially making them more susceptible to pathogen-mediated control, which in termites has proven difficult to implement, in part due to their social immune defence traits, which prevent the spread of infectious disease (He et al., 2018; Davis et al., 2018; Liu et al., 2019; Hassan et al., 2021).

In conclusion, our study provides valuable comparative data, not only for further research exploring the interactions between AKH and AKHR in termites and cockroaches but also in providing necessary sequence information for functional research in an economically and scientifically important group of non-model insects, with potential applications in the development of mimetic or RNAi-based approaches to pest control. The data also constitute an important molecular framework for future studies seeking to exploit neuropeptides as a sustainable means of controlling globally important termite and cockroach pest organisms.

EXPERIMENTAL PROCEDURES

Insect samples

The termite species *Coptotermes formosanus*, *Kaloterms flavicollis*, *Mastotermes darwiniensis*, *Cryptotermes* sp., *Prorethotermes inopinatus* and *Reticulitermes flavipes*, as well as cockroach species, *Blattella germanica* were bred from laboratory colonies of the Federal Institute of Materials Research and Testing (BAM), Berlin, Germany. One species of subsocial wood roaches, *Cryptocercus meridianus* was collected in Yunshanping (27°14'N, 100°23'E), Yulongxueshan, Lijiang, Yunnan, China, and one higher social termite, *Indotermes* sp., was collected in Mengla County, Xishuangbanna Dai Autonomous Prefecture, Yunnan, China (21°62'N, 101°58'E).

Sequence retrieval

AKH preprohormone sequences were retrieved from the published literature (Jedličková et al., 2016) and accessible SRA datasets for *Cryptotermes domesticus* (SRR2039534) and *Reticulitermes grassei* (SRR13251 [02–10]). The following AKHR nucleotide sequences were retrieved from SRA datasets and the nucleotide collection from GenBank: *Periplaneta americana* (DQ217786.1), *Blattella germanica* (GU591493.1), *Zootermopsis nevadensis* (XM_022082426.1) and *Cryptotermes secundus* (XM_033755069.1).

In addition, AKHs and AKHRs were searched from assemblies of 18 cockroach and termite transcriptomes (He et al., 2021) and

transcriptome sequence assemblies (TSA) from ‘The 1KITE project: evolution of insects’ (BioProject: PRJNA183205). We adopted a hidden Markov model (HMM) approach, as described by He et al. (2020) with slight modifications (Data S2). Briefly, candidate AKH and AKHR sequences were searched in the transcriptome assemblies by querying against a reference dataset of diverse AKH precursor and AKHR sequences compiled from NCBI (Table S1). The online tool SignalP 5.0 Server (<https://services.healthtech.dtu.dk/service.php?SignalP-5.0>) was then used to predict the signal peptide cleavage site from each candidate AKH precursor sequence (Almagro Armenteros et al., 2019).

Sequencing the AKH precursor gene and AKH receptor gene

For some species, the complete open reading frame (ORF) of AKHR could be obtained from assembled transcriptome data alone, including *C. wrighti*, *Globitermes* sp., *B. orientalis*, *P. simplex*, *N. castaneus*, *R. grassei*, *Symploce* sp., *Asiablatta kyotensis*, *Paratemnopteryx coulouiana* and *Ischnoptera deropeltiformis*. However, none or only partial AKHR fragment(s) were retrieved from *Indotermes* sp., *C. formosanus*, *C. meridianus*, *K. flavicollis*, *M. darwiniensis*, *P. inopinatus*, *Indotermes* sp. and *R. flavipes*. To obtain these, and to verify the presence of the novel AKH peptide type in *Blattella germanica*, we employed classical molecular methods to attain the full-length AKH/AKHR sequences of these remaining species. ORFfinder (<https://www.ncbi.nlm.nih.gov/orffinder/>) was used to determine ORFs in the sequences.

Apart from *C. meridianus*, whose abdomens were cut with sterile scissors, the whole body of other termite species was used for total RNA isolation. Pre-cooled Trizol reagent (Thermo Fisher Scientific) was used to preserve insect tissues and then homogenised with a FastPrep®-24 homogeniser (MP Biomedicals). Recovery of RNA was followed according to the manufacturer’s recommendation for Trizol (Thermo Fisher Scientific), with chloroform extraction and isopropanol precipitation, followed by re-dissolving RNA in Nuclease-free water, and subsequent incubation with TurboDNase for 30 min at 37°C to remove remaining DNA (TURBO DNA-free Kit, Ambion). cDNA synthesis was performed with oligo-dT primers using the M-MLV Reverse Transcriptase (Promega). PCR was performed using Taq DNA Polymerase (Red Load Taq Master (5x), NEB) with specific or degenerate forward and reverse primers (Table S2).

Primers were manually selected based on conserved homologous nucleotide sequences or designed online: <https://www.ncbi.nlm.nih.gov/tools/primer-blast/> (Ye et al., 2012) using the assembled transcripts and available full-length gene sequences from the species *B. germanica*, *P. americana*, *C. wrighti* and *C. secundus* to guide primer sequence design. PCR products were either cleaned directly with a Monarch® PCR & DNA Cleanup Kit (NEB) (for a specific PCR product) or separated on a 1.5% agarose gel (Agarose NEEO ultra-quality Roth®, Karlsruhe, Germany) and the expected size band excised and extracted using the QIAquick Gel Extraction Kit (Qiagen). Cleaned-up products were then sent for Sanger sequencing (Eurofins Genomic).

Sequence alignment and phylogenetic analysis

The multiple sequence alignment programme MAFFT (Katoh & Standley, 2013) with the E-INS-I algorithm was used to perform multiple sequence alignments for AKH precursor sequences and AKHRs. Alignments were visualised using Jalview (Waterhouse et al., 2009), and the transmembrane domains of the AKHR receptors were predicted with the TMHMM server (<https://services.healthtech.dtu.dk/service.php?TMHMM-2.0>). Pairwise sequence comparisons for AKHR amino acid identity and similarity were computed using the online tool: SIAS (<http://imed.med.ucm.es/Tools/sias.html>) using the ‘length of smallest sequence’ option.

To explore the evolutionary relationships of AKH precursor sequences and AKHR sequences, we carried out phylogenetic reconstruction on amino acid or nucleotide-based alignments. Aligned sequences were trimmed using trimAl v1.2 (Capella-Gutiérrez et al., 2009) with the function -gappout prior to use in phylogenetic tree reconstruction. We employed a maximum likelihood (ML) approach in RAxML v8.2.12 (Stamatakis, 2014) and a Bayesian inference (BI) approach in MrBayes v3.2.7a (Huelsenbeck & Ronquist, 2001) to reconstruct phylogenies.

For ML, we used 1000 rapid bootstrap (RB) replicates with the PROTGAMMAAUTO model (amino acid alignment) or the GTRGAMMAI model (nucleotide alignment). For BI, we employed a model-jumping approach with the command ‘prset aamodelpr = mixed’ (amino acid alignment), or ‘lset nst = mixed’ (nucleotide alignment). Four chains of the Markov Chain Monte Carlo (MCMC) with two independent runs of 15 million generations were conducted, sampling every 500 generations. The average standard deviation of split frequencies was inspected to ensure these were less than 0.01. Posterior probabilities (PPs) were calculated from the posterior distribution of trees after discarding 25% as burn-in.

Prediction of AKHR post-translational modifications (PTMs)

Putative glycosylation and myristoylation sites of AKHRs were detected using MotifScan (https://myhits.sib.swiss/cgi-bin/motif_scan) using default settings (Sigrist et al., 2010) and phosphorylation sites of the intracellular domains of AKHRs were predicted using the NetPhos server (<https://services.healthtech.dtu.dk/service.php?NetPhos-3.1>), displaying only scores higher than a threshold of 0.5 (Blom et al., 1999; Blom et al., 2004).

AUTHOR CONTRIBUTIONS

Shixiong Jiang: Investigation; writing – original draft; methodology; visualization; writing – review and editing; software; formal analysis; data curation; resources. **Heather G. Marco:** Funding acquisition; writing – original draft; validation; writing – review and editing. **Nina Scheich:** Methodology; investigation. **Shulin He:** Methodology; software; writing – review and editing. **Zongqing Wang:** Resources. **Gerd Gäde:** Writing – original draft; validation; writing – review and editing; funding acquisition.

Dino P. McMahon: Conceptualization; investigation; writing – original draft; methodology; validation; writing – review and editing; formal analysis; supervision; resources; project administration; funding acquisition.

ACKNOWLEDGEMENTS

The authors thank the high-performance computing (HPC) Service offered by ZEDAT at Freie Universität Berlin for computing resources (Bennett et al., 2020). SJ was funded by the China Scholarship Council (CSC) (grant no 201906300033). Part of this research was funded by the National Research Foundation of South Africa (grant no 85768 [IFR13020116790] to GG and grant no 10924 to HGM), and via The Research Council of the University of Cape Town as staff awards to GG and HGM. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article

ORCID

Shixiong Jiang  <https://orcid.org/0000-0002-7140-9909>

Heather G. Marco  <https://orcid.org/0000-0003-3000-3010>

Gerd Gäde  <https://orcid.org/0000-0002-7048-9660>

Dino P. McMahon  <https://orcid.org/0000-0003-1119-5299>

REFERENCES

- Abdulganiyyu, I.A., Kaczmarek, K., Zabrocki, J., Nachman, R.J., Marchal, E., Schellens, S. et al. (2020) Conformational analysis of a cyclic AKH neuropeptide analog that elicits selective activity on locust versus honeybee receptor. *Insect Biochemistry and Molecular Biology*, 125, 103362.
- Adamo, S., Roberts, J., Easy, R. & Ross, N. (2008) Competition between immune function and lipid transport for the protein apolipoprotein III leads to stress-induced immunosuppression in crickets. *Journal of Experimental Biology*, 211, 531–538.
- Almagro Armenteros, J.J., Tsirigos, K.D., Sønderby, C.K., Petersen, T.N., Winther, O., Brunak, S. et al. (2019) SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature Biotechnology*, 37, 420–423.
- Altstein, M. & Nässel, D.R. (2010) Neuropeptide signaling in insects. *Advances in Experimental Medicine and Biology*, 692, 155–165.
- Alves-Bezerra, M., De Paula, I.F., Medina, J.M., Silva-Oliveira, G., Medeiros, J.S., Gäde, G. et al. (2016) Adipokinetic hormone receptor gene identification and its role in triacylglycerol metabolism in the blood-sucking insect *Rhodnius prolixus*. *Insect Biochemistry and Molecular Biology*, 69, 51–60.
- Arakane, Y., Li, B., Muthukrishnan, S., Beeman, R.W., Kramer, K.J. & Park, Y. (2008) Functional analysis of four neuropeptides, EH, ETH, CCAP and bursicon, and their receptors in adult ecdysis behavior of the red flour beetle, *Tribolium castaneum*. *Mechanisms of Development*, 125, 984–995.
- Auerswald, L., Siegert, K.J. & Gäde, G. (2005) Activation of triacylglycerol lipase in the fat body of a beetle by adipokinetic hormone. *Insect Biochemistry and Molecular Biology*, 35, 461–470.
- Bártů, I., Tomčala, A., Socha, R., Šimek, P. & Kodrů, D. (2010) Analysis of the lipids mobilized by adipokinetic hormones in the firebug *Pyrhocoris apterus* (Heteroptera: Pyrrhocoridae). *European Journal of Entomology*, 107, 509–520.
- Bell, W.J., Roth, L.M. & Nalepa, C.A. (2007) *Cockroaches: ecology, behavior, and natural history*. Baltimore: Johns Hopkins University Press.
- Belmont, M., Cazzamali, G., Williamson, M., Hauser, F. & Grimmelikhuijzen, C.J. (2006) Identification of four evolutionarily related G protein-coupled receptors from the malaria mosquito *Anopheles gambiae*. *Biochemical and Biophysical Research Communications*, 344, 160–165.
- Bennett, L., Melchers, B. & Proppe, B. (2020) *Curta: a general-purpose high-performance computer at ZEDAT*. Berlin: Freie Universität Berlin.
- Bignell, D.E. (2019) Termite ecology in the first two decades of the 21st century: a review of reviews. *Insects*, 10, 60.
- Bignell, D.E. & Eggleton, P. (2000) Termites in ecosystems. In: *Termites: Evolution, sociality, symbioses, ecology*. Dordrecht: Springer Netherlands.
- Bil, M., Timmermans, I., Verlinden, H. & Huybrechts, R. (2016) Characterization of the adipokinetic hormone receptor of the anautogenous flesh fly, *Sarcophaga crassipalpis*. *Journal of Insect Physiology*, 89, 52–59.
- Birgul Iyison, N., Sinmaz, M.G., Sahbaz, B.D., Shahraki, A., Aksoydan, B., & Durdagi, S. (2020) *In silico* characterization of adipokinetic hormone receptor and screening for pesticide candidates against stick insect, *Carausius morosus*. *Journal of Molecular Graphics and Modelling*, 101, 107720.
- Bläser, M., Misof, B. & Predel, R. (2020) The power of neuropeptide precursor sequences to reveal phylogenetic relationships in insects: a case study on Blattodea. *Molecular Phylogenetics and Evolution*, 143, 106686.
- Bläser, M. & Predel, R. (2020) Evolution of neuropeptide precursors in Polyneoptera (Insecta). *Frontiers in Endocrinology*, 11, 197.
- Blom, N., Gammeltoft, S. & Brunak, S.O.R. (1999) Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *Journal of Molecular Biology*, 294, 1351–1362.
- Blom, N., Sicheritz-Pontén, T., Gupta, R., Gammeltoft, S. & Brunak, S. (2004) Prediction of post-translational glycosylation and phosphorylation of proteins from the amino acid sequence. *Proteomics*, 4, 1633–1649.
- Bourguignon, T., Lo, N., Cameron, S.L., Šobotnik, J., Hayashi, Y., Shigenobu, S. et al. (2014) The evolutionary history of termites as inferred from 66 mitochondrial genomes. *Molecular Biology and Evolution*, 32, 406–421.
- Burand, J.P. & Hunter, W.B. (2013) RNAi: Future in insect management. *Journal of Invertebrate Pathology*, 112, S68–S74.
- Caers, J., Janssen, T., Van Rompay, L., Broeckx, V., Van Den Abbeele, J., Gäde, G. et al. (2016) Characterization and pharmacological analysis of two adipokinetic hormone receptor variants of the tsetse fly, *Glossina morsitans morsitans*. *Insect Biochemistry and Molecular Biology*, 70, 73–84.
- Capella-Gutiérrez, S., Silla-Martínez, J.M. & Gabaldón, T. (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25, 1972–1973.
- Chen, K.Y.M., Keri, D. & Barth, P. (2020) Computational design of G protein-coupled receptor allosteric signal transductions. *Nature Chemical Biology*, 16, 77–86.
- Chino, H., Kiyomoto, Y. & Takahashi, K. (1989) In vitro study of the action of adipokinetic hormone in locusts. *Journal of Lipid Research*, 30, 571–578.
- Chou, K.C. (2020) Progresses in predicting post-translational modification. *International Journal of Peptide Research and Therapeutics*, 26, 873–888.
- Cochran, D.G. (1999) *Cockroaches: their biology, distribution and control*. Document WHO/CDS/WHOPES/99.3. Geneva: World Health Organization.
- Cong, X., Ren, W., Pacalon, J., Xu, R., Xu, L., Li, X. et al. (2022) Large-scale G protein-coupled olfactory receptor-ligand pairing. *ACS Central Science*, 8, 379–387.

- Davis, H.E., Meconcelli, S., Radek, R. & McMahon, D.P. (2018) Termites shape their collective behavioural response based on stage of infection. *Scientific Reports*, 8, 14433.
- Evangelista, D.A., Wipfler, B., Béthoux, O., Donath, A., Fujita, M., Kohli, M.K. et al. (2019) An integrative phylogenomic approach illuminates the evolutionary history of cockroaches and termites (Blattodea). *Proceedings of the Royal Society B: Biological Sciences*, 286, 20182076.
- Gäde, G. (1989) The hypertrehalosaemic peptides of cockroaches: a phylogenetic study. *General and Comparative Endocrinology*, 75, 287–300.
- Gäde, G., Grandcolas, P. & Kellner, R. (1997) Structural data on hypertrehalosaemic neuropeptides from *Cryptocercus punctulatus* and *Theresa petiveriana*: how do they fit into the phylogeny of cockroaches? *Proceedings of the Royal Society B: Biological Sciences*, 264, 763–768.
- Gäde, G. (1997) The explosion of structural information on insect neuropeptides. *Progress in the Chemistry of Organic Natural Products*, 71, 1–128.
- Gäde, G. (2004) Regulation of intermediary metabolism and water balance of insects by neuropeptides. *Annual Review of Entomology*, 49, 93–113.
- Gäde, G. (2006) Unusual complement of three AKH octapeptides in two species of grasshoppers (Caelifera). *European Journal of Entomology*, 103, 297–304.
- Gäde, G. (2009) Peptides of the adipokinetic hormone/red pigment-concentrating hormone family. *Annals of the New York Academy of Sciences*, 1163, 125–136.
- Gäde, G. & Beenackers, A.M.T. (1977) Adipokinetic hormone-induced lipid mobilization and cyclic AMP accumulation in the fat body of *Locusta migratoria* during development. *General and Comparative Endocrinology*, 32, 481–487.
- Gäde, G. & Auerswald, L. (2003) Mode of action of neuropeptides from the adipokinetic hormone family. *General and Comparative Endocrinology*, 132, 10–20.
- Gäde, G. & Goldsworthy, G.J. (2010) Insect peptide hormones: a selective review of their physiology and potential application for pest control. *Pest Management Science*, 59, 1063–1075.
- Gäde, G. & Kellner, R. (1992) Primary structures of the hypertrehalosemic peptides from corpora cardiaca of the primitive cockroach *Polyphaga aegyptiaca*. *General and Comparative Endocrinology*, 86, 119–127.
- Gäde, G. & Marco, H.G. (2005) The adipokinetic hormones of Odonata: a phylogenetic approach. *Journal of Insect Physiology*, 51, 333–341.
- Gäde, G. & Marco, H.G. (2009) Flight-related metabolism and its regulatory peptides in the spittle bug *Locris arithmetica* (Cicadomorpha: Cercopidae) and the stink bugs *Nezara viridula* (Heteroptera: Pentatomidae) and *Encosternum delegorguei* (Heteroptera: Tesseratomidae). *Journal of Insect Physiology*, 55, 1134–1144.
- Gäde, G. & Marco, H.G. (2013) The adipokinetic hormones of African water bugs of the Heteropteran families Nepidae and Belostomatidae. *Physiological Entomology*, 38, 279–291.
- Gäde, G. & Marco, H.G. (2017) The adipokinetic hormone of Mantodea in comparison to other Dictyoptera. *Archives of Insect Biochemistry and Physiology*, 94, e21376.
- Gäde, G. & Marco, H.G. (2022) The adipokinetic peptides of Hemiptera: structure, function, and evolutionary trends. *Frontiers in Insect Science*, 2, 891615.
- Gäde, G., Marco, H.G., Šimek, P. & Marais, E. (2005) The newly discovered insect order Mantophasmatodea contains a novel member of the adipokinetic hormone family of peptides. *Biochemical and Biophysical Research Communications*, 330, 598–603.
- Gäde, G. & Rinehart, K.L., Jr. (1986) Amino acid sequence of a hypertrehalosaemic neuropeptide from the corpus cardiacum of the cockroach, *Nauphoeta cinerea*. *Biochemical and Biophysical Research Communications*, 141, 774–781.
- Gäde, G. & Rinehart, K.L. (1990) Primary structures of hypertrehalosaemic neuropeptides isolated from the corpora cardiaca of the cockroaches *Leucophaea maderae*, *Gromphadorhina portentosa*, *Blattella germanica* and *Blatta orientalis* and of the stick insect *Extatosoma tiaratum* assigned by tandem fast atom bombardment mass spectrometry. *Biological Chemistry Hoppe-Seyler*, 371, 345–354.
- Gäde, G., Šimek, P., Clark, K.D. & Marco, H.G. (2013) Five functional adipokinetic peptides expressed in the corpus cardiacum of the moth genus *Hippotion* (Lepidoptera, Sphingidae). *Regulatory Peptides*, 184, 85–95.
- Gäde, G., Šimek, P. & Marco, H.G. (2017) The African froghopper *Ptyelus flavescens* (suborder: Cicadomorpha) contains two novel and one known peptides of the adipokinetic hormone (AKH) family: structure, function and comparison with aphid AKH (suborder: Sternorrhyncha). *Amino Acids*, 49, 1679–1690.
- Gäde, G., Šimek, P. & Marco, H.G. (2019) Structural diversity of adipokinetic hormones in the hyperdiverse coleopteran Cucujiformia. *Archives of Insect Biochemistry and Physiology*, 102, e21611.
- Gáliková, M., Diesner, M., Klepsatel, P., Hehlert, P., Xu, Y., Bickmeyer, I. et al. (2015) Energy homeostasis control in *Drosophila* adipokinetic hormone mutants. *Genetics*, 201, 665–683.
- Gautam, U.K., Bohatá, A., Shaik, H.A., Zemek, R. & Kodrík, D. (2020) Adipokinetic hormone promotes infection with entomopathogenic fungus *Isaria fumosorosea* in the cockroach *Periplaneta americana*. *Comparative Biochemistry and Physiology Part C Toxicology and Pharmacology*, 229, 108677.
- Goldsworthy, G.J., Jutsum, A.R. & Robinson, N.L. (1975) Proceedings: Adipokinetic hormone and flight metabolism in locusts. *Journal of Endocrinology*, 64, 66–67.
- Goldsworthy, G.J., Lee, M.J., Luswata, R., Drake, A.F. & Hyde, D. (1997) Structures, assays and receptors for locust adipokinetic hormones. *Comparative Biochemistry and Physiology. B: Biochemistry and Molecular Biology*, 117, 483–496.
- Hansen, K.K., Hauser, F., Cazzamali, G., Williamson, M. & Gimmelikhuijzen, C.J. (2006) Cloning and characterization of the adipokinetic hormone receptor from the cockroach *Periplaneta americana*. *Biochemical and Biophysical Research Communications*, 343, 638–643.
- Hassan, A., Huang, Q., Mehmood, N., Xu, H., Zhou, W. & Gao, Y. (2021) Alteration of termite locomotion and allogrooming in response to infection by pathogenic fungi. *Journal of Economic Entomology*, 114, 1256–1263.
- Hauser, F., Søndergaard, L. & Gimmelikhuijzen, C.J. (1998) Molecular cloning, genomic organization and developmental regulation of a novel receptor from *Drosophila melanogaster* structurally related to gonadotropin-releasing hormone receptors from vertebrates. *Biochemical and Biophysical Research Communications*, 249, 822–828.
- Hayes, T.K., Keeley, L.L. & Knight, D.W. (1986) Insect hypertrehalosemic hormone: isolation and primary structure from *Blaberus discoidalis* cockroaches. *Biochemical and Biophysical Research Communications*, 140, 674–678.
- He, S., Johnston, P.R., Kuroпка, B., Lokatis, S., Weise, C., Plarre, R. et al. (2018) Termite soldiers contribute to social immunity by synthesizing potent oral secretions. *Insect Molecular Biology*, 27, 564–576.
- He, S., Johnston, P.R. & McMahon, D.P. (2020) Analyzing immunity in non-model insects using de novo transcriptomics. In: *Immunity in insects*. New York, USA: Springer Protocols Handbooks; Humana, pp. 35–51.
- He, S., Sieksmeyer, T., Che, Y., Mora, M.A.E., Stiblik, P., Banasiak, R. et al. (2021) Evidence for reduced immune gene diversity and activity during the evolution of termites. *Proceedings of the Royal Society B: Biological Sciences*, 288, 20203168.
- Hellemans, S., Wang, M., Hasegawa, N., Šobotník, J., Scheffrahn, R.H. & Bourguignon, T. (2022) Using ultraconserved elements to reconstruct the termite tree of life. *Molecular Phylogenetics and Evolution*, 173, 107520.
- Holwerda, D.A., Van Doorn, J. & Beenackers, A.M. (1977) Characterization of the adipokinetic and hyperglycaemic substances from the locust corpus cardiacum. *Insect Biochemistry*, 7, 151–157.

- Hou, Q.L., Chen, E.H., Jiang, H.B., Wei, D.D., Gui, S.H., Wang, J.J. et al. (2017) Adipokinetic hormone receptor gene identification and its role in triacylglycerol mobilization and sexual behavior in the oriental fruit fly (*Bactrocera dorsalis*). *Insect Biochemistry and Molecular Biology*, 90, 1–13.
- Huang, J.H., Bellés, X. & Lee, H.J. (2012) Functional characterization of hypertrehalosemic hormone receptor in relation to hemolymph trehalose and to oxidative stress in the cockroach *Blattella germanica*. *Frontiers in Endocrinology*, 2, 114.
- Huelsenbeck, J.P. & Ronquist, F. (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17, 754–755.
- Ibrahim, E., Dobeš, P., Kunc, M., Hyršl, P. & Kodrík, D. (2018) Adipokinetic hormone and adenosine interfere with nematobacterial infection and locomotion in *Drosophila melanogaster*. *Journal of Insect Physiology*, 107, 167–174.
- Ibrahim, E., Hejníková, M., Shaik, H.A., Doležel, D. & Kodrík, D. (2017) Adipokinetic hormone activities in insect body infected by entomopathogenic nematode. *Journal of Insect Physiology*, 98, 347–355.
- Inward, D., Beccaloni, G. & Eggleton, P. (2007) Death of an order: a comprehensive molecular phylogenetic study confirms that termites are eusocial cockroaches. *Biology Letters*, 3, 331–335.
- Isabel, G., Martin, J.R.E., Chidami, S., Veenstra, J.A. & Rosay, P. (2005) AKH-producing neuroendocrine cell ablation decreases trehalose and induces behavioral changes in *Drosophila*. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 288, R531–R538.
- Jackson, G.E., Gäde, G. & Marco, H.G. (2022) *In silico* screening for pesticide candidates against the desert locust *Schistocerca gregaria*. *Life (Basel, Switzerland)*, 12, 387.
- Jackson, G.E., Pavadai, E., Gäde, G. & Andersen, N.H. (2019) The adipokinetic hormones and their cognate receptor from the desert locust, *Schistocerca gregaria*: solution structure of endogenous peptides and models of their binding to the receptor. *PeerJ*, 7, e7514.
- Jedlička, P., Steinbauerová, V., Šimek, P. & Zahradníčková, H. (2012) Functional characterization of the adipokinetic hormone in the pea aphid, *Acyrtosiphon pisum*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 162, 51–58.
- Jedlička, V., Jedlička, P. & Špuláková, B. (2016) Diversity of adipokinetic hormones in termites (Isoptera). *European Journal of Entomology*, 113, 364.
- Kapan, N., Lushchak, O.V., Luo, J. & Nässel, D.R. (2012) Identified peptidergic neurons in the *Drosophila* brain regulate insulin-producing cells, stress responses and metabolism by coexpressed short neuropeptide F and corazonin. *Cellular and Molecular Life Sciences*, 69, 4051–4066.
- Katoh, K. & Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30, 772–780.
- Kaufmann, C. & Brown, M.R. (2006) Adipokinetic hormones in the African malaria mosquito, *Anopheles gambiae*: identification and expression of genes for two peptides and a putative receptor. *Insect Biochemistry and Molecular Biology*, 36, 466–481.
- Kaufmann, C. & Brown, M.R. (2008) Regulation of carbohydrate metabolism and flight performance by a hypertrehalosaemic hormone in the mosquito *Anopheles gambiae*. *Journal of Insect Physiology*, 54, 367–377.
- Khan, M.A. & Ahmad, W. (2018) Termites: An Overview. Termites and Sustainable Management. In: *Volume 1-Biology, social behaviour and economic importance*, pp. 1–25. Switzerland: Springer.
- Klass, K.D., Nalepa, C. & Lo, N. (2008) Wood-feeding cockroaches as models for termite evolution (Insecta: Dictyoptera): *Cryptocercus* vs. *Parasphaeria boleiriana*. *Molecular Phylogenetics and Evolution*, 46, 809–817.
- Kodrík, D. (2008) Adipokinetic hormone functions that are not associated with insect flight. *Physiological Entomology*, 33, 171–180.
- Kodrík, D., Socha, R., Šimek, P., Zemek, R. & Goldsworthy, G.J. (2000) A new member of the AKH/RPCH family that stimulates locomotory activity in the firebug, *Pyrrhocoris apterus* (Heteroptera). *Insect Biochemistry and Molecular Biology*, 30, 489–498.
- König, S., Albers, C. & Gäde, G. (2005) Mass spectral signature for insect adipokinetic hormones. *Rapid Communications in Mass Spectrometry*, 19, 3021–3024.
- Konuma, T., Morooka, N., Nagasawa, H. & Nagata, S. (2012) Knockdown of the adipokinetic hormone receptor increases feeding frequency in the two-spotted cricket *Gryllus bimaculatus*. *Endocrinology*, 153, 3111–3122.
- Krishna, K., Grimaldi, D.A., Krishna, V. & Engel, M.S. (2013) Treatise on the Isoptera of the world: Basal families. *Bulletin of the American Museum of Natural History*, 2013, 200–623.
- Lee, D.W., Shrestha, S., Kim, A.Y., Park, S.J., Yang, C.Y., Kim, Y. et al. (2011) RNA interference of pheromone biosynthesis-activating neuropeptide receptor suppresses mating behavior by inhibiting sex pheromone production in *Plutella xylostella* (L). *Insect Biochemistry and Molecular Biology*, 41, 236–243.
- Li, B., Predel, R., Neupert, S., Hauser, F., Tanaka, Y., Cazzamali, G. et al. (2008) Genomics, transcriptomics, and peptidomics of neuropeptides and protein hormones in the red flour beetle *Tribolium castaneum*. *Genome Research*, 18, 113–122.
- Li, S., Hauser, F., Skadborg, S.K., Nielsen, S.V., Kirketerp-Møller, N. & Grimmelikhuijzen, C.J. (2016) Adipokinetic hormones and their G protein-coupled receptors emerged in Lophotrochozoa. *Scientific Reports*, 6, 32789.
- Liebrich, W., Kellner, R. & Gäde, G. (1995) Isolation and primary structures of neuropeptides of the AKH/RPCH family from various termite species. *Peptides*, 16, 559–564.
- Liu, L., Wang, W., Liu, Y., Sun, P., Lei, C. & Huang, Q. (2019) The influence of allogrooming behavior on individual innate immunity in the subterranean termite *Reticulitermes chinensis* (Isoptera: Rhinotermitidae). *Journal of Insect Science*, 19, 6.
- Marchal, E., Schellens, S., Monjon, E., Bruyninckx, E., Marco, H.G., Gäde, G. et al. (2018) Analysis of peptide ligand specificity of different insect adipokinetic hormone receptors. *International Journal of Molecular Sciences*, 19, 542.
- Marciniak, P.L., Pacholska-Bogalska, J. & Ragionieri, L. (2022) Neuropeptidomes of *Tenebrio molitor* L. and *Zophobas atratus* Fab. (Coleoptera, Polyphaga: Tenebrionidae). *Journal of Proteome Research*, 21, 2247–2260.
- Marco, H.G. & Gäde, G. (2017) Structure and function of adipokinetic hormones of the large white butterfly *Pieris brassicae*. *Physiological Entomology*, 42, 103–112.
- Marco, H.G. & Gäde, G. (2019) Five neuropeptide ligands meet one receptor: How does this tally? A structure-activity relationship study using adipokinetic bioassays with the sphingid moth, *Hippotion eson*. *Frontiers in Endocrinology*, 10, 231.
- Marco, H.G., Katali, O. & Gäde, G. (2018) Influence of aminergic and peptidergic substances on heart beat frequency in the stick insect *Carausius morosus* (Insecta, Phasmatodea). *Archives of Insect Biochemistry and Physiology*, 98, e21469.
- Marco, H.G., König, S. & Gäde, G. (2022) Mass spectrometric proof of predicted peptides: novel adipokinetic hormones in insects. *Molecules*, 27, 6469.
- Marco, H.G., Šimek, P. & Gäde, G. (2014) Adipokinetic hormones of the two extant apterygotan insect orders, Archaeognatha and Zygentoma. *Journal of Insect Physiology*, 60, 17–24.
- Marco, H.G., Šimek, P. & Gäde, G. (2020) Unique members of the adipokinetic hormone family in butterflies and moths (Insecta, Lepidoptera). *Frontiers in Physiology*, 11, 614552.

- Möller, S., Vilo, J. & Croning, M.D. (2001) Prediction of the coupling specificity of G protein coupled receptors to their G proteins. *Bioinformatics*, 17, S174–S181.
- Oudejans, R.C., Kooiman, F.P., Heerma, W., Versluis, C., Slotboom, A.J. & Beenakkers, A.M.T. (1991) Isolation and structure elucidation of a novel adipokinetic hormone (Lom-AKH-III) from the glandular lobes of the corpus cardiacum of the migratory locust, *Locusta migratoria*. *European Journal of Biochemistry*, 195, 351–359.
- Park, S., Sonn, J.Y., Oh, Y., Lim, C. & Choe, J. (2014) SIFamide and SIFamide receptor define a novel neuropeptide signaling to promote sleep in *Drosophila*. *Molecules and Cells*, 37, 295.
- Park, Y., Kim, Y.J. & Adams, M.E. (2002) Identification of G protein-coupled receptors for *Drosophila* PRXamide peptides, CCAP, corazonin, and AKH supports a theory of ligand-receptor coevolution. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 11423–11428.
- Predel, R. & Gäde, G. (2005) Peptidomics of neurohemal organs from species of the cockroach family Blattidae: How do neuropeptides of closely related species differ. *Peptides*, 26, 3–9.
- Rewitz, K.F., Yamanaka, N., Gilbert, L.I. & O'Connor, M.B. (2009) The insect neuropeptide PTTH activates receptor tyrosine kinase torso to initiate metamorphosis. *Science*, 326, 1403–1405.
- Rios, C., Jordan, B., Gomes, I. & Devi, L. (2001) G-protein-coupled receptor dimerization: Modulation of receptor function. *Pharmacology and Therapeutics*, 92, 71–87.
- Robinson, N. & Goldsworthy, G. (1977) A possible site of action for adipokinetic hormone on the flight muscle of locusts. *Journal of Insect Physiology*, 23, 153–158.
- Rosenstreich, D.L., Eggleston, P., Kattan, M., Baker, D., Slavin, R.G., Gergen, P. et al. (1997) The role of cockroach allergy and exposure to cockroach allergen in causing morbidity among inner-city children with asthma. *New England Journal of Medicine*, 336, 1356–1363.
- Roth, S., Fromm, B., Gäde, G. & Predel, R. (2009) A proteomic approach for studying insect phylogeny: CAPA peptides of ancient insect taxa (Dictyoptera, Blattoptera) as a test case. *BMC Ecology and Evolution*, 9, 1–12.
- Sajwan, S., Sidorov, R., Stašková, T., Žaloudíková, A., Takasu, Y., Kodrík, D. et al. (2015) Targeted mutagenesis and functional analysis of adipokinetic hormone-encoding gene in *Drosophila*. *Insect Biochemistry and Molecular Biology*, 61, 79–86.
- Scarborough, R.M., Jamieson, G.C., Kalish, F., Kramer, S.J., Mckenroe, G.A., Miller, C.A. et al. (1984) Isolation and primary structure of two peptides with cardioacceleratory and hyperglycemic activity from the corpora cardiaca of *Periplaneta americana*. *Proceedings of the National Academy of Sciences*, 81, 5575–5579.
- Schwartz, T.W. (1994) Locating ligand-binding sites in 7TM receptors by protein engineering. *Current Opinion in Biotechnology*, 5, 434–444.
- Siegert, K., Morgan, P. & Mordue, W. (1985) Primary structures of locust adipokinetic hormones II. *Biological Chemistry Hoppe-Seyler*, 366, 723–727.
- Sigrist, C.J., Cerutti, L., De Castro, E., Langendijk-Genevaux, P.S., Bulliard, V., Bairoch, A. et al. (2010) PROSITE, a protein domain database for functional characterization and annotation. *Nucleic Acids Research*, 38, D161–D166.
- Stamatakis, A. (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30, 1312–1313.
- Stank, A., Kokh, D.B., Fuller, J.C. & Wade, R.C. (2016) Protein binding pocket dynamics. *Accounts of Chemical Research*, 49, 809–815.
- Staubli, F., Jørgensen, T.J., Cazzamali, G., Williamson, M., Lenz, C., Søndergaard, L. et al. (2002) Molecular identification of the insect adipokinetic hormone receptors. *Proceedings of the National Academy of Sciences*, 99, 3446–3451.
- Stone, J.V., Mordue, W., Batley, K.E. & Morris, H.R. (1976) Structure of locust adipokinetic hormone, a neurohormone that regulates lipid utilisation during flight. *Nature*, 263, 207–211.
- Su, N.Y. & Scheffrahn, R.H. (2000) Termites as pests of buildings. In: *Termites: Evolution, Sociality, Symbioses, Ecology*. Dordrecht: Springer Netherlands.
- Tang, B., Cheng, Y., Li, Y., Li, W., Ma, Y., Zhou, Q. et al. (2020) Adipokinetic hormone regulates cytochrome P450-mediated imidacloprid resistance in the brown planthopper, *Nilaparvata lugens*. *Chemosphere*, 259, 127490.
- Ulyshen, M.D. (2016) Wood decomposition as influenced by invertebrates. *Biological Reviews*, 91, 70–85.
- Van Der Horst, D.J., Van Marrewijk, W.J. & Diederens, J.H. (2001) Adipokinetic hormones of insect: release, signal transduction, and responses. *International Review of Cytology*, 211, 179–240.
- Veenstra, J.A. & Camps, F. (1990) Structure of the hypertrehalosaemic neuropeptide of the German cockroach, *Blattella germanica*. *Neuropeptides*, 15, 107–109.
- Veenstra, J.A. (2014) The contribution of the genomes of a termite and a locust to our understanding of insect neuropeptides and neurohormones. *Frontiers in Physiology*, 5, 454.
- Venkatakrishnan, A., Deupi, X., Lebon, G., Tate, C.G., Schertler, G.F. & Babu, M.M. (2013) Molecular signatures of G-protein-coupled receptors. *Nature*, 494, 185–194.
- Ware, J.L., Litman, J., Klass, K.D. & Spearman, L.A. (2008) Relationships among the major lineages of Dictyoptera: the effect of outgroup selection on dictyopteran tree topology. *Systematic Entomology*, 33, 429–450.
- Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M. & Barton, G.J. (2009) Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics*, 25, 1189–1191.
- Whetstone, P.A. & Hammock, B.D. (2007) Delivery methods for peptide and protein toxins in insect control. *Toxicol*, 49, 576–596.
- Wicher, D., Agricola, H.J., Söhler, S., Gundel, M., Heinemann, S.H., Wollweber, L. et al. (2006) Differential receptor activation by cockroach adipokinetic hormones produces differential effects on ion currents, neuronal activity, and locomotion. *Journal of Neurophysiology*, 95, 2314–2325.
- Withers, J. & Dong, X. (2017) Post-translational regulation of plant immunity. *Current Opinion in Plant Biology*, 38, 124–132.
- Witten, J.L., Schaffer, M.H., O'Shea, M., Cook, J.C., Hemling, M.E. & Rinehart, K.L., Jr. (1984) Structures of two cockroach neuropeptides assigned by fast atom bombardment mass spectrometry. *Biochemical and Biophysical Research Communications*, 124, 350–358.
- Xia, Y., Xia, C., Pan, X. & Shen, H.B. (2022) BindWeb: a web server for ligand binding residue and pocket prediction from protein structures. *Protein Science*, 31, e4462.
- Yang, H., Huang, J., Liu, Y., Li, J., Luo, S. & Wu, J. (2018) Prediction of the post-translational modifications of adipokinetic hormone receptors from solitary to eusocial bees. *Sociobiology*, 65, 271–279.
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S. & Madden, T.L. (2012) Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, 13, 1–11.
- Yu, Z., Shi, J., Jiang, X., Song, Y., Du, J. & Zhao, Z. (2023) Neuropeptide F regulates feeding via the juvenile hormone pathway in *Ostrinia furnacalis* larvae. *Pest Management Science*, 79, 1193–1203.
- Zandawala, M., Hamoudi, Z., Lange, A.B. & Orchard, I. (2015) Adipokinetic hormone signalling system in the Chagas disease vector, *Rhodnius prolixus*. *Insect Molecular Biology*, 24, 264–276.
- Zeng, H., Qin, Y., Du, E., Wei, Q., Li, Y., Huang, D. et al. (2020) Genomics- and peptidomics-based discovery of conserved and novel neuropeptides in the American cockroach. *Journal of Proteome Research*, 20, 1217–1228.
- Zhu, C., Huang, H., Hua, R., Li, G., Yang, D., Luo, J. et al. (2009) Molecular and functional characterization of adipokinetic hormone receptor and its peptide ligands in *Bombyx mori*. *FEBS Letters*, 583, 1463–1468.

Ziegler, R., Isoe, J., Moore, W., Riehle, M. & Wells, M. (2011) The putative AKH receptor of the tobacco hornworm, *Manduca sexta*, and its expression. *Journal of Insect Science*, 11, 12–18.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Data S1. Supporting Information.

How to cite this article: Jiang, S., Marco, H.G., Scheich, N., He, S., Wang, Z., Gäde, G. et al. (2023) Comparative analysis of adipokinetic hormones and their receptors in Blattodea reveals novel patterns of gene evolution. *Insect Molecular Biology*, 1–19. Available from: <https://doi.org/10.1111/imb.12861>