Identification of a Glucose Permease from *Mycobacterium smegmatis* mc² 155

Elisângela F. Pimentel-Schmitt^a Knut Jahreis^b Mike P. Eddy^c Johannes Amon^a
Andreas Burkovski^a Fritz Titgemeyer^a

^aDepartment of Microbiology, Friedrich Alexander University Erlangen-Nürnberg, Erlangen;

^bDepartment of Biology and Chemistry, University of Osnabrück, Osnabrück, Germany;

^cBioTechHome.com, Del Mar, Calif., USA

Key Words

Sugar transport · Membrane topology · *Mycobacterium* · Pathogenicity · Bacterial lifestyle · Carbon regulation

Abstract

We report here the molecular identification of a glucose permease from *Mycobacterium smegmatis*, a model organism for our understanding of the life patterns of the major pathogens *Mycobacterium tuberculosis* and *Mycobacterium leprae*. A computer-based search of the available genome of *M. smegmatis* mc² 155 with the sequences of well-characterized glucose transporters revealed the gene *msmeg4187* as a possible candidate. The deduced protein belongs to the major facilitator superfamily of proton symporters and facilitators and exhibits up to 53% of amino acid identity to other members of this family. Heterologous expression of *msmeg4187* in an *Escherichia coli* glucose-negative mutant led to the restoration of growth on glucose. The determination of the biochemical features characterize MSMEG4187 (GlcP) as a high affinity (K_m of 19 μ M), glucose-specific permease. The results represent the first molecular characterization of a sugar permease in mycobacteria, and thus supply fundamental data for further in-depth analysis on the nutritional lifestyle of these bacteria.

Copyright © 2008 S. Karger AG, Basel

Introduction

Mycobacterium tuberculosis remains a serious threat to human health [Kaufmann, 2006]. About one-third of the world's population is thought to be infected with this pathogenic microorganism. In spite of significant investments in research, the mechanisms of its pathogenicity are still not fully understood. One aspect that is inextricably linked to pathogenicity is nutrition. This has been recently demonstrated by the identification of central metabolic genes, including carbohydrate transporter genes, which are essential for entering into and proliferating inside the macrophage [Joshi et al., 2006]. The current literature reveals limited information about the intermediary metabolism, utilization of organic compounds in the process of growth, and carbon catabolite control of mycobacteria. Efforts have been undertaken to understand virulence in mycobacteria by using the fast-growing *Mycobacterium smegmatis*. This model organism is infrequently reported as pathogenic and causes skin or soft tissue infections [Brinton et al., 1991; Lustgarten, 1884].

We reported previously the identification of a glucose kinase gene from *M. smegmatis* mc² 155, which could be a potential signalling factor for carbon regulation [Pimentel-Schmitt et al., 2006]. Since the transport of a carbon source represents, for many pathways, the decisive

step for the flux through the pathway [Ruyter et al., 1991], we aimed to unravel the glucose uptake system(s). A putative glucose transporter that is encoded by the gene *msmeg4187* was recently detected by us in the course of a genome analysis of the carbohydrate permeases of *M. smegmatis* and *M. tuberculosis* [Titgemeyer et al., 2007].

In this communication, we demonstrate the function of the possible glucose transporter by heterologous complementation in an appropriate mutant of *Escherichia coli*. Furthermore, we provide biochemical details on the catalytic function of GlcP (MSMEG4187) and present a detailed topology model that includes the coordinates of conserved amino acid residues among related bacterial glucose permeases. GlcP^{Msm} is to the best of our knowledge the first molecular described sugar permease in the genus mycobacteria.

Results

msmeg4187 Encodes a Putative Glucose Permease

The protein database of the *M. smegmatis* genome was scanned with several known glucose permease protein sequences using BLASTP as implemented at The Institute for Genomic Research (TIGR) [Parche et al., 2006; van Wezel et al., 2005]. We found only one gene (*msmeg4187*) that encodes a predicted glucose permease of the major facilitator super family (MFS) [Pao et al., 1998]. The deduced protein shares 53% amino acid identity to the well-characterized glucose permease GlcP of *Streptomyces coelicolor* [Titgemeyer et al., 2007; van Wezel et al., 2005]. Analysis of the gene locus of *msmeg4187* (hereafter named *glcP*) indicated that the gene is transcribed as a monocistronic mRNA (fig. 1). *glcP* is flanked downstream by a putative dicistronic *hisEG* operon of genes involved in histidine biosynthesis, and upstream by an unknown open reading frame. While the *hisEG* operon is conserved in mycobacteria, it appears that the slow-growing mycobacteria, e.g. *M. tuberculosis*, *M. bovis*, and *M. avium*, do not have an obvious homologue of *glcP*.

The multiple alignment shown in figure 1b includes four homologous glucose permeases which have been experimentally described [Parche et al., 2006; van Wezel et al., 2005; Weisser et al., 1995; Zhang et al., 1989]. The range of protein identity is from 34 to 53%, showing 67 conserved residues in these sequences. The two typical signature sequences of the MFS are found in GlcP^{Msm} at positions 126–151 (VgGIGvGVasviapayiaETspggiR) and

331 to 348 (AIALIDKIGRKpllligS). Hydrophobicity profiles showed that all five sequences have 12 predicted transmembrane helices that are at equivalent positions (fig. 1b). To establish the phylogenetic relationships of GlcP^{Msm}, we selected 19 protein sequences of the MFS which have been characterized by experimental investigation and those from diverse mycobacteria (fig. 1c). GlcP^{Msm} clusters with the *S. coelicolor* homolog and the one from the cyanobacterium *Synechocystis* [van Wezel et al., 2005; Zhang et al., 1989]. Interestingly, the xylose permease Xyle of *E. coli* is more closely related to GlcP^{Msm} than expected, while the bifidobacterial GlcP is more distant but closely associated with another xylose permease, XylT of *Lactobacillus brevis* [Chaillou et al., 1998]. This suggests that glucose and xylose transporter genes may have evolved following two different branches from the common origins.

Heterologous Expression of *GlcP* in *E. coli*

The sequence of *msmeg4187* was amplified from chromosomal DNA and cloned into plasmid pSU2719 in orientation of the *lacZ* gene, yielding plasmid pFT283 (*glcP*⁺). The plasmid was introduced into *E. coli* LJ140 (Δ *ptsHICrr::kan*), which does not transport glucose to appreciable

Fig. 1. Genetic organization of the *glcP* gene in *M. smegmatis*, protein alignment, and phylogenetic tree. **a** Genetic map. The depicted region of *glcP* (black) in *M. smegmatis* includes the genes *hisG*, *hisE* (both histidine biosynthetic genes), and a gene encoding a protein of unknown function (*urf*, MSMEG4188). The genome of *M. smegmatis* (mc² 155) was sequenced by TIGR (<http://www.tigr.org/>). The arrows indicate length and transcriptional orientation of annotated genes which are depicted by the respective *orf* number. Numbers in square brackets show the lengths of intergenic regions in bp. **b** The amino acid sequences of *glcP* in *M. smegmatis* were aligned with representative homologues. Residues identical in all proteins are shaded black and those that are conserved are shown in grey. Above the alignment, the residues present in all proteins are marked by asterisks. Abbreviations and protein designations are as follows. Msm, *M. smegmatis* (MSMEG4187); Sco, *S. coelicolor* (SCO5578); Syn, *Synechocystis* sp. PCC6803 (sl10771); Blo, *B. longum* (BL1631); Zmo, *Z. mobilis* (ZMO0366). **c** Phylogenetic tree of the *glcP* family. An unrooted phylogenetic tree was computed with the CLUSTALW software making use of the implemented neighbor joining method with the function for evolutionary distance correction. Evolutionary distances are proportional to the branch length. 19 protein sequences were selected as indicated in the figure. The references for experimentally verified transporters are as follows: Chaillou et al., 1998; Krispin and Allmansberger, 1998; Martin et al., 1994; Parche et al., 2006; van Wezel et al., 2005; Weisser et al., 1995; Zhang et al., 1989.

Fig. 2. Heterologous complementation and transport kinetics. **a** Complementation of the glucose-negative phenotype in *E. coli* LJ140 (pFT283 *glcP*⁺) is shown on a MacConkey agar base plate supplemented with 50 mM glucose. Pigmented red (here grey) colonies indicate the fermentation of glucose, whereas white colonies reflect a deficiency in glucose fermentation. DH5 α (pSU2719) was used as a positive control (dark grey) and LJ140 (pSU2719) as the congenic negative control. **b** Time-dependent uptake of 100 μ M [¹⁴C]glucose using cells of DH5 α (pSU2719) (circles, positive control), LJ140 (pFT283) (triangles) and LJ140 (pSU2719) negative control (squares). Standard deviations of triplicate data points are shown by error bars but are barely visible due to minimal data variation. **c** [¹⁴C]glucose uptake at different concentrations were fit to a non-linear regression and **(d)** graphed into a Lineweaver-Burk plot revealing a K_m of 19 μ M and V_{max} of 0.63 nmol min⁻¹ OD₆₀₀⁻¹.

amounts due to a chromosomal deletion of the *pts* operon. LJ140(pFT283), and several control strains were streaked onto MacConkey agar plates supplemented with 50 mM glucose (fig. 2a). The presence of *M. smegmatis glcP* led to the restoration of glucose fermentation as indicated by the formation of gray colonies. In comparison, the congenic strain LJ140(pSU2719) that carries the same cloning vector without the *glcP* gene grew as white colonies.

GlcP Is a High-Affinity Glucose Permease with Narrow Substrate Specificity

Glucose uptake experiments were performed to further characterize the biochemical features of the GlcP^{Msm} permease (fig. 2b). While LJ140(pSU2719) showed negligible uptake kinetics, transport of glucose was readily detectable when *msmeg4187* was expressed in LJ140(pFT283), reaching about 50% of the intrinsic *E. coli* wild-type glucose transport activity that was monitored in *E. coli* DH5 α (pSU2719). A series of uptake experiments with various glucose concentrations was then performed to determine the kinetic parameters of GlcP (fig. 2c, d). We determined an affinity constant (K_m) of 19 μ M and a velocity (V_{max}) of 0.63 nmol min⁻¹ OD₆₀₀⁻¹. The substrate specificity was investigated by a competitive inhibition of glucose transport. Therefore, we added a hundredfold excess (10 mM) of nonradioactive sugar. While the presence

of excess glucose and the derivative 2-deoxyglucose led to a reduction of more than 80% of the initial uptake rate, no or very small effects (<10%) were observed, when methyl α -D-glucopyranoside, xylose, arabinose, fructose, and trehalose were tested. In conclusion, the biochemical analysis of GlcP^{Msm} revealed a high-affinity permease with a narrow substrate specificity.

Discussion

The MFS family is one of the largest transporter families with members in bacteria and in eukaryotes like the GLUT facilitator transporters of mammals [Pao et al., 1998]. The transmembrane proteins of the GLUT family exhibit different substrate specificities, kinetic properties and tissue expression profiles. GLUT1 for example is expressed as a high-affinity permease in erythrocytes, the perineurium of peripheral nerves, and capillary endothelial cells of the blood-brain barrier [Gould et al., 1991; Mueckler et al., 1985; Salas-Burgos et al., 2004]. GLUT2 is a low-affinity glucose transporter, expressed in the liver, pancreas, intestine, and kidneys [Fukumoto et al., 1988; Gould et al., 1991]. GLUT3 and GLUT4 are high-affinity symporters present in brain, heart, muscle, and brown adipose tissue [Fukumoto et al., 1989; Kayano et

Fig. 3. Topology model of *M. smegmatis* GlcP. The drawing displays the predicted two-dimensional topology structure of GlcP, which is composed of 12 putative transmembrane segments. Conserved residues present in bacterial GlcPs (fig. 1b) are highlighted by black circles. Charged amino acids are marked by (+) and (−), showing a charge distribution of more residues on the inside of a transmembrane helix compared to the outside as defined by the positive-inside rule described by Heijne [1986].

al., 1988; Wood and Trayhurn, 2003]. Thus, this type of glucose permease has obviously been very successful in evolution.

In this study, we report by molecular means the first mycobacterial sugar permease and established that heterologous gene expression of a mycobacterial integral membrane protein in *E. coli* is feasible. The permease is the glucose-specific transport protein GlcP from *M. smegmatis* mc² 155 encoded by open reading frame *msmeg4187*. GlcP^{Msm} exhibited an apparent K_m of 19 μ M when expressed in *E. coli*. This correlates with the corresponding values of 41 μ M of *S. coelicolor*'s GlcP [van Wezel et al., 2005] and 70 μ M of *B. longum*'s GlcP [Parche et al., 2006]. GlcP^{Msm} shows the closest phylogenetic relationship to glucose permeases from streptomycetes and cyanobacteria [van Wezel et al., 2005; Zhang et al., 1989], which differ in their substrate specificity for fructose. GlcP^{Msm} was found specific for glucose and 2-deoxyglucose and not for fructose. Thus, it resembles in its substrate specificity profile the homolog GlcP^{Sco} from *S. coelicolor* [van Wezel et al., 2005]. In fact, our recent genome-based compilations of all carbohydrate transport systems of *S. coelicolor* and *M. smegmatis* shows that these two soil-dwelling actinomycetes have many homologous sugar uptake systems in common [Bertram et al., 2004; Titgemeyer et al., 2007].

Our bioinformatic analysis on the membrane topology of GlcP^{Msm} led to the model presented in figure 3. GlcP^{Msm} has a predicted composition of 12 transmembrane helices typical for MFS porters [Pao et al., 1998]. The model was obtained by aligning the predictions of possible transmembrane segments of the five GlcP proteins depicted in figure 1b assuming that they all share the same overall topology. This showed that 12 transmembrane segments are probably present in all of them, while sequence variations in terms of deletions and inser-

tions occurred only in regions which were predicted as flexible loops. Figure 3 displays that helices I, IV, and VII exhibit (with eight, six, and seven conserved residues) the highest conservation suggesting that these may be most relevant for substrate recognition.

Interestingly, no GlcP homolog is present in slow-growing mycobacteria, notably *M. tuberculosis* and *M. leprae* [Titgemeyer et al., 2007]. In these bacteria, glucose may enter either by simple diffusion, which could be efficient enough to meet the nutritional demands of these extremely slow-growing bacteria (generation time of weeks), or through another, yet unknown permease [Titgemeyer et al., 2007]. *M. smegmatis* and also the slow-growing mycobacterial species possess a glucose kinase that converts the incoming glucose to glucose-6-phosphate [Pimentel-Schmitt et al., 2006; Titgemeyer et al., 2007]. Apart from its catalytic function, this enzyme confers global carbon regulation in closely related streptomycetes [Angell et al., 1994]. We could recently show in *S. coelicolor* that glucose kinase binds to glucose permease GlcP [van Wezel et al., 2007]. This protein-protein interaction is thus thought to influence the regulatory function of glucose kinase. For this reason, it would be worthwhile to study, whether GlcP and GlkA of *M. smegmatis* also interact with each other and whether this has similar consequences regarding nutritional control. This in turn would help to understand how nutrition and virulence may be linked in mycobacteria.

Experimental Procedures

Bacterial Strains and Growth Conditions

Cells of *E. coli* were grown at 37°C under vigorous shaking in Luria-Bertani broth (LB) and supplemented with the appropriate antibiotics when required. *E. coli* DH5 α served as the host strain for gene cloning purposes [Sambrook et al., 1989]. LJ140 (Δ *ptsHI*::

kan) is a derivative of *E. coli* LJ110 [Zeppenfeld et al., 2000], in which a deletion of the *pts* operon was introduced by common P1(kc) transduction through selection of a gene for kanamycin resistance which substitutes the *pts* locus.

Heterologous Complementation

Chromosomal DNA was prepared from 100-ml cultures grown to saturation [Belisle and Sonnenberg, 1998]. 400 ng was used as template for a polymerase chain reaction with oligonucleotides FS12 ggctgtctagatcattgggccgcaggaaga and FPS21 tcg-cgcgatgggctctagatccgtcagt (engineered *Hind*III and *Xba*I restriction sites are underlined). Primer FS12 contained stop codons in all three reading frames and an artificial ribosome binding site for *glcP* to ensure efficient translation. The amplified DNA fragment was digested and ligated in pSU2719 in the same orientation as the *lacZ* gene [Chandler, 1991]. The ligation mixture was transformed into DH5 α . Recombinant plasmids were isolated and confirmed by DNA sequencing. The resulting *glcP^{Msm}* expression vector was designated pFT283. LJ140 was transformed with pFT283 or pSU2719 (control), and transformant colonies were streaked out on MacConkey agar plates supplemented with 50 mM glucose in order to examine heterologous complementation of the glucose transport-negative phenotype.

Transport Assays

Cells of *E. coli* LJ140 bearing either plasmid pSU2719 or pFT283 *glcP⁺* and *E. coli* DH5 α (pSU2719) were grown at 37°C in 20 ml LB supplemented with chloramphenicol to an OD₆₀₀ of 1.0. Cells were pre-equilibrated to 37°C before uptake was initiated by the addition of [¹⁴C]glucose at a final concentration of 100 μ M (100,000 cpm/ml) [Nothaft et al., 2003]. Samples of 0.5 ml were taken at different time points, rapidly filtered through nitrocellulose filters, and washed with 5 ml chilled 0.1 M LiCl. Radioactivity was determined by scintillation oscillography with 3 ml scintillation solution per filter. Uptake rates are expressed in nanomoles min⁻¹ OD₆₀₀⁻¹. For determination of the kinetic parameters,

glucose uptake experiments were conducted with various glucose concentrations (12.5–400 μ M). To determine V_{max} and K_m, the data were fit to the Michaelis-Menten equation by a nonlinear regression program and graphed on a Lineweaver-Burk plot. The substrate specificity was studied by glucose uptake competition assay using a 100-fold excess of nonradioactive sugar (10 mM) as described previously [van Wezel et al., 2005]. Transport activity without additional sugar was set to 100% (control).

Computational Analyses

The genome data of *M. smegmatis* mc² 155 were collected from the primary annotation database (<http://cmr.tigr.org/tigr-scripts/CMR/shared/Genomes.cgi>) available at The Institute for Genomic Research (TIGR). Protein databank searches were carried out at the BLAST server of the National Center for Biotechnology Information at the National Institutes of Health, Bethesda, Md., USA (<http://www.ncbi.nlm.nih.gov>), the BLAST server of the Transport Classification Databank, TCDB (www.tcdb.org), and at TIGR. Multiple alignment analysis and graphical optimization were achieved with CLUSTALW (www.ebi.ac.uk/clustalw) and BOXSHADE (www.ch.embnet.org/software/BOX_form.html) software. Hydrophobicity profiles and topology predictions were calculated with the TopPred II software [Claros and von Heijne, 1994]. The phylogenetic tree was calculated with the neighbor joining method as implemented in CLUSTALW. The tree was graphically visualized by importing phylip (*.ph) files into the TREEVIEW (<http://taxonomy.zoology.gla.ac.uk/rod/treeview.html>) software that was kindly provided by Rod Page.

Acknowledgements

This study was supported through grants of SFB473 to F.T., GK805 to E.F.P.-S., and SFB431 to K.J. of the Deutsche Forschungsgemeinschaft.

References

- Angell S, Lewis CG, Buttner MJ, Bibb MJ: Glucose repression in *Streptomyces coelicolor* A3(2): a likely regulatory role for glucose kinase. *Mol Gen Genet* 1994;244:135–143.
- Belisle JT, Sonnenberg MG: Isolation of genomic DNA from mycobacteria. *Methods Mol Biol* 1998;101:31–44.
- Bertram R, Schlicht M, Mahr K, Nothaft H, Sailer MH Jr, Titgemeyer F: In silico and transcriptional analysis of carbohydrate uptake systems of *Streptomyces coelicolor* A3(2). *J Bacteriol* 2004;186:1362–1373.
- Brinton LA, Li JY, Rong SD, Huang S, Xiao BS, Shi BG, Zhu ZJ, Schiffman MH, Dawsey S: Risk factors for penile cancer: results from a case-control study in China. *Int J Cancer* 1991;47:504–509.
- Chaillou S, Bor YC, Batt CA, Postma PW, Pouwels PH: Molecular cloning and functional expression in *Lactobacillus plantarum* 80 of *xylT*, encoding the D-xylose-H⁺ symporter of *Lactobacillus brevis*. *Appl Environ Microbiol* 1998;64:4720–4728.
- Chandler MS: New shuttle vectors for *Haemophilus influenzae* and *Escherichia coli*: P15A-derived plasmids replicate in *H. influenzae* Rd. *Plasmid* 1991;25:221–224.
- Claros MG, von Heijne G: TopPred II: an improved software for membrane protein structure predictions. *Comput Appl Biosci* 1994;10:685–686.
- Fukumoto H, Seino S, Imura H, Seino Y, Eddy RL, Fukushima Y, Byers MG, Shows TB, Bell GI: Sequence, tissue distribution, and chromosomal localization of mRNA encoding a human glucose transporter-like protein. *Proc Natl Acad Sci USA* 1988;85:5434–5438.
- Fukumoto H, Kayano T, Buse JB, Edwards Y, Pilch PF, Bell GI, Seino S: Cloning and characterization of the major insulin-responsive glucose transporter expressed in human skeletal muscle and other insulin-responsive tissues. *J Biol Chem* 1989;264:7776–7779.
- Gould GW, Thomas HM, Jess TJ, Bell GI: Expression of human glucose transporters in *Xenopus* oocytes: kinetic characterization and substrate specificities of the erythrocyte, liver, and brain isoforms. *Biochemistry* 1991;30:5139–5145.
- Heijne G: The distribution of positively charged residues in bacterial inner membrane proteins correlates with the trans-membrane topology. *EMBO J* 1986;5:3021–3027.

- Joshi SM, Pandey AK, Capite N, Fortune SM, Rubin EJ, Sasseti CM: Characterization of mycobacterial virulence genes through genetic interaction mapping. *Proc Natl Acad Sci USA* 2006;103:11760–11765.
- Kaufmann SH: Envisioning future strategies for vaccination against tuberculosis. *Nat Rev Immunol* 2006.
- Kayano T, Fukumoto H, Eddy RL, Fan YS, Byers MG, Shows TB, Bell GI: Evidence for a family of human glucose transporter-like proteins: sequence and gene localization of a protein expressed in fetal skeletal muscle and other tissues. *J Biol Chem* 1988;263:15245–15248.
- Krispin O, Allmansberger R: The *Bacillus subtilis* AraE protein displays a broad substrate specificity for several different sugars. *J Bacteriol* 1998;180:3250–3252.
- Lustgarten S: Über spezifische Bacillen in syphilitischen Krankheitsprodukten. *Wienerische Medizinische Wochenschrift* 1884;1.
- Martin GE, Seamon KB, Brown FM, Shanahan MF, Roberts PE, Henderson PJ: Forskolin specifically inhibits the bacterial galactose- H^+ transport protein, GalP. *J Biol Chem* 1994;269:24870–24877.
- Mueckler M, Caruso C, Baldwin SA, Panico M, Blench I, Morris HR, Allard WJ, Lienhard GE, Lodish HF: Sequence and structure of a human glucose transporter. *Science* 1985;229:941–945.
- Nothaft H, Parche S, Kamionka A, Titgemeyer F: In vivo analysis of HPr reveals a fructose-specific phosphotransferase system that confers high-affinity uptake in *Streptomyces coelicolor*. *J Bacteriol* 2003;185:929–937.
- Pao SS, Paulsen IT, Saier MH Jr: Major facilitator superfamily. *Microbiol Mol Biol Rev* 1998;62:1–34.
- Parche S, Beleut M, Rezzonico E, Jacobs D, Arigoni F, Titgemeyer F, Jankovic I: Lactose-over-glucose preference in *Bifidobacterium longum* NCC2705:glcP, encoding a glucose transporter, is subject to lactose repression. *J Bacteriol* 2006;188:1260–1265.
- Pimentel-Schmitt EF, Thomae AW, Amon J, Klieber AM, Roth H-M, Muller YA, Jahreis K, Burkovski A, Titgemeyer F: A glucose kinase from *Mycobacterium smegmatis*. *J Mol Microbiol Biotechnol* 2007;12:75–81.
- Ruyter GJ, Postma PW, van Dam K: Control of glucose metabolism by enzyme II^{Glc} of the phosphoenolpyruvate-dependent phosphotransferase system in *Escherichia coli*. *J Bacteriol* 1991;173:6184–6191.
- Salas-Burgos A, Iserovich P, Zuniga F, Vera JC, Fischbarg J: Predicting the three-dimensional structure of the human facilitative glucose transporter glut1 by a novel evolutionary homology strategy: insights on the molecular mechanism of substrate migration, and binding sites for glucose and inhibitory molecules. *Biophys J* 2004;87:2990–2999.
- Sambrook J, Fritsch EF, Maniatis T: Molecular Cloning: A Laboratory Manual, ed 2. Cold Spring Harbor, Cold Spring Harbor Laboratory Press, 1989.
- Titgemeyer F, Amon J, Parche S, Mahfoud M, Bail J, Schlicht M, Rehm N, Hillmann D, Stephan J, Walter B, Burkovski A, Niederweis M: A genomic view of sugar transport in *Mycobacterium smegmatis* and *Mycobacterium tuberculosis*. *J Bacteriol* 2007;189:5903–5915.
- van Wezel GP, Mahr K, König M, Traag BA, Pimentel-Schmitt EF, Willimek A, Titgemeyer F: GlcP constitutes the major glucose uptake system of *Streptomyces coelicolor* A3(2). *Mol Microbiol* 2005;55:624–636.
- van Wezel GP, König M, Mahr K, Nothaft H, Thomae AW, Bibb MJ, Titgemeyer F: A new piece of an old jigsaw: glucose kinase is activated posttranslationally in a glucose transport-dependent manner in *Streptomyces coelicolor* A3 (2). *J Mol Microbiol Biotechnol* 2007;12:65–72.
- Weisser P, Krämer R, Sahm H, Sprenger GA: Functional expression of the glucose transporter of *Zymomonas mobilis* leads to restoration of glucose and fructose uptake in *Escherichia coli* mutants and provides evidence for its facilitator action. *J Bacteriol* 1995;177:3351–3354.
- Wood IS, Trayhurn P: Glucose transporters (GLUT and SGLT): expanded families of sugar transport proteins. *Br J Nutr* 2003;89:3–9.
- Zeppenfeld T, Larisch C, Lengeler JW, Jahreis K: Glucose transporter mutants of *Escherichia coli* K-12 with changes in substrate recognition of IICB^{Glc} and induction behavior of the *ptsG* gene. *J Bacteriol* 2000;182:4443–4452.
- Zhang CC, Durand MC, Jeanjean R, Joset F: Molecular and genetical analysis of the fructose-glucose transport system in the cyanobacterium *Synechocystis* PCC6803. *Mol Microbiol* 1989;3:1221–1229.