Cite this: *Chem. Commun.*, 2015, 51, 14877

Received 9th July 2015,
Accepted 7th August 2015

DOI: 10.1039/c5cc05679a

www.rsc.org/chemcomm

Thermal dimerization of $[n]$ cumulenes ($n = 5, 7, 9$)†

Dominik Wendinger, Johanna A. Januszewski, Frank Hampel and Rik R. Tykwinski*

The thermal dimerization of three $[n]$ cumulenes ($n = 5, 7$, and 9) has been investigated, and a common reactivity pattern is observed that gives radialenes and expanded radialenes through regioselective cycloaddition reactions; all three products are characterized using X-ray crystallography.

Cycloaddition reactions of polyynes have been used to create conjugated molecules that are otherwise difficult to obtain, and the regioselectivity of these reactions can often be controlled by the electronic nature of endcapping groups.^{1–3} On the other hand, this same level of control has not yet been achieved for reactions for the sp-carbon chain of long $[n]$ cumulenes ($n \geq 5$).⁴ There have been a handful of reports in which $[5]$ cumulenes react with electron-poor olefins or alkynes, but the regiochemistry of cycloaddition reactions is difficult to predict.^{5–8} Dimerization reactions under thermal, photochemical, or metal-catalysed conditions that give cyclic oligoynes or radialenes are also known.⁹ The regiochemistry observed for these cycloaddition reactions has been typically attributed to either steric requirements or to the electronic structure of endcapping groups.^{10–14} For example, the thermal dimerization of a $[5]$ cumulene with bulky alkyl substituents, such as $[5]tBu$ (Scheme 1), results in dimerization at the central γ -bond to form the symmetric $[4]$ radialene **1a** (Scheme 1).⁷ $[5]$ Cumulenes functionalized with smaller groups (e.g., $[5]H$ or $[5]Me$) show dimerization at the α -bonds, to give cyclynes **2**.^{13,15,16} In the case of aryl substituents, the thermal reaction of $[5]Ph$ forms **3** (Scheme 1), presumably *via* $[4]$ radialene **1b** and a subsequent $[4+2]$ cycloaddition with $[5]Ph$ followed by a $[2+2]$ electrocyclicization.¹⁷ Finally, the dimerization of $[5]$ cumulenes under Ni-catalysis has been reported by Iyoda and coworkers,¹² and the regiochemical outcome of these reactions

centers on the reaction at either the β - or γ -bond, and is dependent on the size and electronic makeup of the end groups. The reactivity of longer “odd” $[n]$ cumulenes (*i.e.*, $n = 7$ or 9) in cycloaddition reactions has not, to our knowledge, been reported to date.

We describe herein the dimerization of $[n]$ cumulenes, including the first reported reactions of $[7]$ - and $[9]$ cumulenes. These transformations follow a similar reactivity pattern, and the products provide a unique series of expanded radialenes that have been characterized by X-ray crystallography.

Intrigued by the formation of **3** from $[5]Ph$ as reported by Kawamura and coworkers, the thermal reaction of $[5]tBuPh$, recently synthesized in our laboratory,¹⁸ was examined. After heating in toluene at reflux for 4 d (Scheme 2), the crude reaction mixture showed many products (by TLC analysis). The main product could be isolated by flash chromatography on silica gel, although NMR spectroscopy indicated the presence of several compounds. On the other hand, mass spectrometry showed a signal consistent with the mass expected for a dimer of $[5]tBuPh$ (m/z 1657.3). Single crystals could be obtained by

Scheme 1 Examples of thermal di- and trimerization reactions of $[5]$ cumulenes.

Department of Chemistry and Pharmacy & Interdisciplinary Center for Molecular Materials (ICMM), Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Henkestrasse 42, 91054 Erlangen, Germany. E-mail: rik.tykwinski@fau.de; Web: <http://www.chemie.uni-erlangen.de/tykwinski>

† Electronic supplementary information (ESI) available. CCDC 1402907–1402909. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5cc05679a

Scheme 2 Thermal dimerization reactions of [5]tBuPh and [5]Tol.

layering a CH_2Cl_2 solution with hexanes, and X-ray crystallography of the colorless crystals (*vide infra*) revealed the formation of the cross-conjugated,¹⁹ four-fold symmetric [4]radialene **4a**. Further optimization of this reaction was explored, but, quite surprisingly, all attempts to repeat the formation of **4a** were unsuccessful. It seemed clear that the increased steric bulk of the di-*tert*-butyl phenyl groups of [5]tBuPh was a problem in comparison to the phenyl groups in [5]Ph. Thus, the use of a less sterically shielded [5]cumulene was explored, [5]Tol, a member of a new series of [n]cumulenes encapped by *ortho*-tolyl substituents (Scheme 2).²⁰

Scheme 3 Dimerization of [7]- and [9]Tol to expanded radialenes **5** and **6**, respectively.

All attempts to date toward converting [5]Tol to radialene **4b** have been frustratingly unsuccessful (Scheme 2), and we have not been able to expand on the preliminary formation of **4a**.

Although the investigations using [5]cumulenes were not generally successful, subsequent work with [7]Tol gave an interesting result.²⁰ When handling [7]Tol in solution at rt, it was obvious upon TLC analysis that a second compound was being formed. ^1H and ^{13}C NMR spectroscopy offered only inconclusive evidence that the “by-product” contained a plethora of sp - and sp^2 -hybridized carbon atoms, in addition to the presence of methyl groups. Attempted purification of [7]Tol *via* crystallization at rt gave red crystals, and X-ray crystallographic analysis confirmed the formation of the dimeric, expanded radialene **5** (Scheme 3). Unlike the reactions with [5]tBuPh, however, the dimerization of [7]Tol was quite reproducible. In a solution of CH_2Cl_2 , the yield of **5** increased with longer reaction time, but the overall conversion was limited by the stability of [7]Tol, which decomposes in solution over a period of hours to give a light brown, insoluble precipitate. In the best case, the yield of **5** was 24%. Drawing from the seminal work by Iyoda and coworkers,¹² $[\text{Ni}(\text{CO})_2\text{Cl}_2]$ was explored as a catalyst toward improving the yield of **5**, but these attempts were not successful. Finally, it is worth noting that the dimerization reaction could be prevented if solutions of [7]Tol were maintained at a low temperature (-20°C).

The synthesis of [9]Tol provided an equally interesting outcome.²⁰ It was quickly noted that the stability of [9]Tol was reduced compared to that of [5]Tol and [7]Tol, and the purple solutions of [9]Tol in CH_2Cl_2 discolored within minutes at rt. Thus, a solution of [9]Tol in CH_2Cl_2 was layered with MeOH and stored at -20°C . After several days grey needles were formed, and X-ray structural analysis indicated that the dimerization of [9]Tol had occurred to give expanded radialene **6** (Scheme 3). Similar to the dimerization of [7]Tol to **5**, the formation of expanded radialene **6** was also reproducible, giving yields of 5–10%.

With the X-ray crystallographic data of **4a**, **5**, and **6** in hand, general structural characteristics were examined in comparison to other analogous macrocycles (Fig. 1–3). The exocyclic allene units of **5** and **6** are, in general, slightly bent from linearity, showing angles of *ca.* 173 – 177° . This is, more or less, consistent with the values for **7**¹² and **8**,²¹ which fall between 174° and 178° . In the case of **4a**, the bending of the allene results in

Fig. 1 ORTEP drawing of radialene **4a** (ellipsoids at 30% probability level; hydrogen atoms not shown), in comparison to **7**–**9**.

Fig. 2 ORTEP drawing of expanded radialene **5** (ellipsoids at 30% probability level; hydrogen atoms and cocrystallized solvent molecules not shown), in comparison to **10–13**.

Fig. 3 ORTEP drawing of expanded radialene **6** (ellipsoids at 30% probability level; hydrogen atoms not shown), in comparison to **2b**, **14**, and **15**.

dispersive interactions between the *tert*-butyl group and the neighboring aromatic ring, with the shortest C–H··· π distance of 2.95 Å (Fig. 1).²²

For symmetrical [4]radialenes **4a**, **7**, and **9**,²³ the vinylidene/allenylidene angles (133–137°) deviate little from the expected values of 135°. In unsymmetric radialene **8**, on the other hand, the steric bulk of the *tert*-butyl groups seems to play a role, and the two allene units are slightly expanded away from the vinyl groups, giving reduced bond angles on the allene “side” (129.0°). Finally, the presence of the vinylidene phenyl rings in the plane of the octaphenyl[4]radialene **9** produces a puckered structure (34.7°) that is similar to cyclobutane (*ca.* 32–35°, see ref. 24).

The structure of expanded radialene **5** (Fig. 2) is most similar to that of **10**, reported by Diederich and coworkers.²⁵ Compound **5** also shares structural similarities with cyclooctadiene **11**,²⁶ as well as annulenes **12**^{27,28} and **13**.²⁹ The acetylene units of **5** are significantly bent at angles of 156.8° and 156.9°, which are in the same range as the values determined for molecules **10–12** (155.3–159.3°). It is interesting to consider the transannular distances between the acetylene carbons in **5**, from the standpoint of homoconjugation. The closest distance in **5**, C4–C5', is 2.64 Å, which is in good agreement with other known cyclooctadienes **10** (2.59 Å), **11** (2.60 Å), **12** (2.62 Å), and **13** (2.57 and 2.60 Å). In all cases, this distance is less than the van der Waals radii, and, according to Gleiter and coworkers, a transannular distance of 3 Å or less is necessary for through-space interactions to occur.³⁰ It is, however, hard to determine if through-space conjugation plays a significant role.³¹

A comparison of the structure of **6** to that of the antiaromatic annulene **14**,³² dodecatetrayne **2b**,^{10,33} and benzoannulenes such as **15**³⁴ shows that all four molecules share a very similar

geometry (Fig. 3). Namely, the C–C–C bonds fall within a narrow range of 164.5–167.9°.

A couple of comments regarding the ¹³C NMR shifts of the sp-hybridized allene carbon of compounds **5** and **6** are noteworthy, since the resonances fall in a unique range³⁵ (the ¹³C NMR spectrum of **4b** was, unfortunately, not available). In compounds **5** and **6**, the allene carbons resonate at 207.9 and 209.7 ppm, respectively, consistent with other 1,1-diarylallenes (210–200 ppm).³⁶ Tetraalkylallenes, on the other hand, generally show allene resonances in the range of 200–165 ppm.³⁷ The more the ring strain present in the terminating alkyl rings, the farther upfield the signal shifts, as evident by a comparison of resonances in **7** (cyclopentylidene endgroup) and **8** (cyclopropylidene endgroup) at 189.8 and 178.7 ppm, respectively. Thus, ring strain might be reflected in the chemical shift of the allene carbon of **5** and **6**, although the effect is small.

Finally, it is perhaps expected that the molecules such as **5** and **6** feature low energy absorptions due to their conjugated, carbon-rich framework. Like other radialenes and expanded radialenes without donor and/or acceptor groups, however, expanded radialenes **5** and **6** absorb in the high-energy region of the UV/Vis spectrum, with lowest energy absorption maxima (λ_{max}) at 275 and 359 nm, respectively.¹⁹ This is at much higher energy compared to, for example, dehydroannulenes such as **14** (λ_{max} = 660 nm),³² where linear-conjugation extends throughout the macrocycle.

In summary, we have reported the formation of three new radialenes (**4a**, **5**, and **6**) from the dimerization of [*n*]cumulenes (*n* = 5, 7, and 9). The formation of **5** and **6** from [7]**Tol** and [9]**Tol**, respectively, offers the first examples of expanded radialenes with allenylidene groups at the vertices. All three products are

characterized using X-ray crystallography, and the most interesting structural feature is the significant strain in the acetylene unit (C3–C4–C5) in **5**, which is bent 23.2° from linearity. The dimerization reactions of all three derivatives, **[5]tBuPh**, **[7]Tol**, and **[9]Tol**, follow a similar pattern, and we are currently exploring if these reactions follow a concerted (*i.e.*, pericyclic) or radical pathway.

Funding from the University of Erlangen-Nuremberg and the Deutsche Forschungsgemeinschaft (DFG – SFB 953, “Synthetic Carbon Allotropes”) is gratefully acknowledged.

Notes and references

- 1 P. Reutenauer, M. Kivala, P. D. Jarowski, C. Boudon, J.-P. Gisselbrecht, M. Gross and F. Diederich, *Chem. Commun.*, 2007, 4898; M. Kivala, C. Boudon, J.-P. Gisselbrecht, P. Seiler, M. Gross and F. Diederich, *Angew. Chem., Int. Ed.*, 2007, **46**, 6357.
- 2 M. Kivala and F. Diederich, *Acc. Chem. Res.*, 2009, **42**, 235.
- 3 R. M. Bettison, P. B. Hitchcock and D. R. M. Walton, *J. Organomet. Chem.*, 1988, **341**, 247.
- 4 The discussion here centers on longer “odd” $[n]$ cumulenes, *i.e.*, those derivatives with $n = 5, 7, 9$.
- 5 J. A. Januszewski, F. Hampel, C. Neiss, A. Goerling and R. R. Tykwinski, *Angew. Chem., Int. Ed.*, 2014, **53**, 3743.
- 6 B. Bildstein, *Coord. Chem. Rev.*, 2000, **206–207**, 369.
- 7 H. D. Hartzler, *J. Am. Chem. Soc.*, 1971, **93**, 4527.
- 8 H. D. Hartzler, *J. Am. Chem. Soc.*, 1966, **88**, 3155.
- 9 J. A. Januszewski and R. R. Tykwinski, *Chem. Soc. Rev.*, 2014, **43**, 3184.
- 10 L. T. Scott and G. J. DeCicco, *Tetrahedron Lett.*, 1976, **17**, 2663.
- 11 H. Hopf and G. Maas, *Angew. Chem., Int. Ed.*, 1992, **31**, 931.
- 12 Y. Kuwatani, G. Yamamoto, M. Oda and M. Iyoda, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 2188.
- 13 M. Kaftory, I. Agmon, M. Ladika and P. J. Stang, *J. Am. Chem. Soc.*, 1987, **109**, 782.
- 14 S. Basak, S. Srivastava and W. J. LeNoble, *J. Org. Chem.*, 1987, **52**, 5095.
- 15 L. T. Scott and G. J. DeCicco, *J. Org. Chem.*, 1980, **45**, 4055.
- 16 C. Werner, H. Hopf, I. Dix, P. Bubenitschek and P. G. Jones, *Chem. – Eur. J.*, 2007, **13**, 9462.
- 17 N. Islam, T. Ooi, T. Iwasawa, M. Nishiuchi and Y. Kawamura, *Chem. Commun.*, 2009, 574.
- 18 J. A. Januszewski, D. Wendinger, C. D. Methfessel, F. Hampel and R. R. Tykwinski, *Angew. Chem., Int. Ed.*, 2013, **52**, 1817.
- 19 M. Gholami and R. R. Tykwinski, *Chem. Rev.*, 2006, **106**, 4997.
- 20 See the ESI† for the synthesis of **[5]**-, **[7]**-, and **[9]Tol**.
- 21 A. E. Learned, A. M. Arif and P. J. Stang, *J. Org. Chem.*, 1988, **53**, 3122.
- 22 Similar CH– π interactions were also observed in the solid-state structure of **[3]tBuPh**, see ref. 18.
- 23 M. Iyoda, H. Otani, M. Oda, Y. Kai, Y. Baba and N. Kasai, *J. Am. Chem. Soc.*, 1986, **108**, 5371.
- 24 A. Stein, C. W. Lehmann and P. Luger, *J. Am. Chem. Soc.*, 1992, **114**, 7684.
- 25 S. Kammermeier, R. R. Tykwinski, P. Siemsen, P. Seiler and F. Diederich, *Chem. Commun.*, 1998, 1285.
- 26 E. Kloster-Jensen and J. Wirz, *Helv. Chim. Acta*, 1975, **58**, 162.
- 27 For the synthesis of **12** see: H. N. C. Wong, P. J. Garratt and F. Sondheimer, *J. Am. Chem. Soc.*, 1974, **96**, 5604.
- 28 R. Destro, T. Pilati and M. Simonetta, *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.*, 1977, **33**, 447.
- 29 C.-Y. Leung, T. C. W. Mak and H. N. C. Wong, *J. Chem. Crystallogr.*, 1996, **26**, 227.
- 30 R. Gleiter, D. Kratz, W. Schaefer and V. Schehlmann, *J. Am. Chem. Soc.*, 1991, **113**, 9258; see also ref. 31.
- 31 S. Eisler, R. McDonald, G. R. Loppnow and R. R. Tykwinski, *J. Am. Chem. Soc.*, 2000, **122**, 6917.
- 32 J. Anthony, C. B. Knobler and F. Diederich, *Angew. Chem., Int. Ed.*, 1993, **32**, 406.
- 33 K. N. Houk, L. T. Scott, N. G. Rondan, D. C. Spellmeyer, G. Reinhardt, J. L. Hyun, G. J. DeCicco, R. Weiss, M. H. M. Chen, L. S. Bass, J. Clardy, F. S. Jorgensen, T. A. Eaton, V. Sarkozi, C. M. Petit, L. Ng and K. D. Jordan, *J. Am. Chem. Soc.*, 1985, **107**, 6556.
- 34 U. H. F. Bunz and V. Enkelmann, *Chem. – Eur. J.*, 1999, **5**, 263.
- 35 K. B. Wiberg, J. D. Hammer, K. W. Zilm and J. R. Cheeseman, *J. Org. Chem.*, 1999, **64**, 6394.
- 36 See, for examples: S. Ma and Q. He, *Tetrahedron*, 2006, **62**, 2769.
- 37 See, for examples: W. Ando, H. Hayakawa and N. Tokitoh, *Tetrahedron Lett.*, 1986, **27**, 6357; S. I. Kozhushkov, T. Haumann, R. Boese and A. de Meijere, *Angew. Chem., Int. Ed.*, 1993, **32**, 401; M. Eckert-Maksic, S. Zoellner, W. Goethling, R. Boese, L. Maksimovic, R. Machinek and A. de Meijere, *Chem. Ber.*, 1991, **124**, 1591.