

## Supplement

Table S1. Mean  $\pm$  SD (Min–Max) values for measurements of morphological characters used for field identification and calculated wing morphometry of captured individuals of *P. auritus* ( $n = 51$ ) and *P. austriacus* ( $n = 26$ ); \* significant ( $p \leq 0.005$ ) and \*\* highly significant ( $p \leq 0.001$ ) variation between species.

	<i>Plecotus auritus</i>		<i>Plecotus austriacus</i>	
	<i>mean <math>\pm</math> SD</i>	<i>min–max</i>	<i>mean <math>\pm</math> SD</i>	<i>min–max</i>
Head-Body-Length, <i>HBL</i> [mm]	52.43 $\pm$ 4.95	42.0–59.0	54.34 $\pm$ 3.84	42.5–60.5
Thumb length, <i>TL</i> [mm]*	6.79 $\pm$ 0.45	5.5–7.1	6.16 $\pm$ 0.58	5.4–7.1
Claw length, <i>CL</i> [mm]*	2.55 $\pm$ 0.28	1.9–3.0	1.84 $\pm$ 0.33	1.5–2.1
Hind foot length, <i>HF</i> [mm]	8.3 $\pm$ 1.19	6.9–10.6	8.24 $\pm$ 0.59	7.0–9.0
Ear length, <i>EL</i> [mm]	36.77 $\pm$ 2.88	26.2–38.5	36.92 $\pm$ 2.91	28.6–41.0
Tragus length, <i>TrL</i> [mm]*	15.57 $\pm$ 0.42	14.9–16.2	15.86 $\pm$ 2.97	14.5–17.0
Tragus width, <i>TrW</i> [mm]*	4.91 $\pm$ 0.38	4.2–5.5	5.93 $\pm$ 0.18	5.5–6.3
Body mass, <i>m</i> [g]	8.97 $\pm$ 1.03	6.9–10.8	9.49 $\pm$ 0.04	7.9–11.0
Forearm length, <i>FA</i> [mm]	40.66 $\pm$ 2.1	35.1–43.2	42.1 $\pm$ 1.64	37.0–43.2
Length of 3 <sup>rd</sup> finger, <i>D3</i> [mm]**	65.48 $\pm$ 0.98	64.0–67.0	68.48 $\pm$ 2.23	64.5–71
Length of 5 <sup>th</sup> finger, <i>D5</i> [mm]**	53.6 $\pm$ 1.3	49.0–55.0	52.68 $\pm$ 1.26	49.0–54.5
Wing span, <i>WS</i> [mm]*	259.61 $\pm$ 7.1	240.0–271.0	268.34 $\pm$ 5.95	255.0–281.0
Wing area, <i>S</i> [ $\text{m}^2$ ]*	0.0093 $\pm$ 0.0003	0.0084–0.0099	0.0093 $\pm$ 0.0004	0.0083–0.01
Aspect ratio, <i>AR</i> **	7.29 $\pm$ 0.37	5.97–7.85	7.73 $\pm$ 0.29	6.9–8.14
Wing loading, <i>WL</i> [ $\text{N}/\text{m}^2$ ]**	8.61 $\pm$ 0.78	6.84–10.9	9.98 $\pm$ 0.84	8.37–11.61