

---

# Enlargement of filtrations and continuous Girsanov-type embeddings

Stefan Ankirchner<sup>1</sup>, Steffen Dereich<sup>2</sup>, and Peter Imkeller<sup>1</sup>

<sup>1</sup> Institut für Mathematik  
Humboldt-Universität zu Berlin  
Unter den Linden 6  
10099 Berlin  
Germany  
[ankirch@math.tu-berlin.de](mailto:ankirch@math.tu-berlin.de)   [imkeller@mathematik.hu-berlin.de](mailto:imkeller@mathematik.hu-berlin.de)

<sup>2</sup> Fachbereich Mathematik  
Technische Universität Berlin  
Strasse des 17. Juni 136  
10623 Berlin  
Germany  
[dereich@math.tu-berlin.de](mailto:dereich@math.tu-berlin.de)

**Summary.** Let  $(\mathcal{G}_t)$  be an enlargement of the filtration  $(\mathcal{F}_t)$ . Jeulin and Jacod discussed a sufficient criterion for the inheritance of the semimartingale property when passing to the larger filtration. We provide alternative proofs of their results in a more general setting by using decoupling measures and Girsanov's changes of measure. We derive necessary and sufficient conditions for the embedding of vector spaces of  $(\mathcal{F}_t)$ -semimartingales into spaces of  $(\mathcal{G}_t)$ -semimartingales to be continuous in terms of generalized entropies of the information increment.

**2000 AMS subject classifications:** primary 60G44; secondary 60G48, 94A17, 62B10.

**Key words and phrases:** enlargement of filtration; semimartingale; decoupling measure; Girsanov's theorem; Shannon information; entropy.

## Introduction

On a probability space, let  $(\mathcal{G}_t)$  be a filtration containing a smaller filtration  $(\mathcal{F}_t)$ . The basic question of the well known theory of enlargement of filtrations (see [JY85]) with some relevance in simple models of financial markets with asymmetric information (see for instance [Imk03]) is this: under which conditions every  $(\mathcal{F}_t)$ -semimartingale remains a semimartingale relative to  $(\mathcal{G}_t)$ ? In the pioneering papers of [JY85] this inheritance property has been called 'Hypothèse (H)'. Jacod [Jac85] gives a sufficient criterion for it to hold and studies semimartingales' Doob-Meyer decompositions relative to  $(\mathcal{G}_t)$ . With

respect to vector space topologies on the set of  $(\mathcal{F}_t)$ - and  $(\mathcal{G}_t)$ -semimartingales Yor [Yor85] investigates continuity properties of the associated mapping of  $(\mathcal{F}_t)$ -semimartingales into the space of  $(\mathcal{G}_t)$ -semimartingales.

In this paper we reconsider the problem of the inheritance of the semimartingale property from a different and more general perspective. In fact, in section 1 we derive inheritance results generalizing Jacod's [Jac85], which were proved in the setting of initial enlargements by the information stored in random elements with values in Lusin spaces. Our proofs are based on the concept of the *decoupling measure*, which allows an independent view on the additional information contained in the enlarged filtration, specified in  $\sigma$ -fields  $\mathcal{H}_t$  enlarging  $\mathcal{F}_t$  to obtain  $\mathcal{G}_t = \mathcal{F}_t \vee \mathcal{H}_t$ . The key observation is that under the decoupling measure every  $(\mathcal{F}_t)$ -martingale is a  $(\mathcal{G}_t)$ -martingale. Hence, enlarging the filtration can be seen as stepping from a view of processes through the decoupling measure to a view by the original measure. In particular, the associated Girsanov transform can be used to obtain explicit representations of the Doob-Meyer decomposition w.r.t. the larger filtration. This idea goes back to [FI93], where this method was used to analyze initial enlargements of the Wiener filtration by some random variable  $G$ . Later [AIS98] and [GP98] extended these techniques to more general stochastic bases and semimartingales. In more recent approaches it was rediscovered in terms of a Bayesian interpretation of simple models of insider trading by Gasbarra and Valkeila [GV03]. Of course, the cost of this approach consists in the very assumption of the existence of the decoupling measure. It restricts generality to a non-trivial extent, as is seen if compared for example to the setting of [ADI04]. For instance, if the *information drift* to be deducted from a martingale in the larger filtration does not generate an equivalent martingale measure capturing the change of views from the small to the large filtration, then there will be no decoupling measure. In order to tackle the problem, as Yoeurp [Yoe85] for the analysis of progressive enlargements, we choose a formulation a product space: the first marginal contains the original information, while the second describes the additional information. Under the product measure both marginals are independent. Therefore it will be the appropriate candidate for our decoupling measure.

Here is an outline of the structure of the material presented. Our main occupation in section 1 consists in showing how objects are transferred from the original space into the artificial product space and vice versa. Once this is handled, an application of the Girsanov transform leads to explicit Doob-Meyer decompositions. In section 2 we provide estimates of the strength of the information drift by appropriate generalized entropies. These are used in section 3 in order to prove continuity properties of the embedding of the  $(\mathcal{F}_t)$ -semimartingales into the set of  $(\mathcal{G}_t)$ -semimartingales with respect to well known vector space topologies. These results generalize continuity results obtained by Yor [Yor85].

## 1 Enlargement of filtrations and Girsanov's theorem

Let  $(\Omega, \mathcal{F}, P)$  be a probability space with right-continuous filtrations  $(\mathcal{F}_t)_{t \geq 0}$  and  $(\mathcal{H}_t)_{t \geq 0}$ . Moreover, let  $\mathcal{F}_\infty = \bigvee_{t \geq 0} \mathcal{F}_t$  and  $\mathcal{H}_\infty = \bigvee_{t \geq 0} \mathcal{H}_t$ .

Our objective is to study the enlarged filtration

$$\mathcal{G}_t = \bigcap_{s > t} (\mathcal{F}_s \vee \mathcal{H}_s), \quad t \geq 0.$$

We relate this enlargement to a measure change on the product space

$$\bar{\Omega} = \Omega \times \Omega$$

equipped with the  $\sigma$ -field

$$\bar{\mathcal{F}} = \mathcal{F}_\infty \otimes \mathcal{H}_\infty.$$

We endow  $\bar{\Omega}$  with the filtration

$$\bar{\mathcal{F}}_t = \bigcap_{s > t} (\mathcal{F}_s \otimes \mathcal{H}_s), \quad t \geq 0.$$

$\Omega$  will be embedded into  $\bar{\Omega}$  by the map

$$\psi : (\Omega, \mathcal{F}) \rightarrow (\bar{\Omega}, \bar{\mathcal{F}}), \quad \omega \mapsto (\omega, \omega).$$

We denote by  $\bar{P}$  the image of the measure  $P$  under  $\psi$ , i.e.

$$\bar{P} = P_\psi.$$

Hence for all  $\bar{\mathcal{F}}$ -measurable functions  $f : \bar{\Omega} \rightarrow \mathbb{R}$  we have

$$\int f(\omega, \omega') d\bar{P}(\omega, \omega') = \int f(\omega, \omega) dP(\omega). \quad (1)$$

In the following measure the two components in  $\bar{\Omega}$  are decoupled, and weighted according to  $P$ :

$$\bar{Q} = P|_{\mathcal{F}_\infty} \otimes P|_{\mathcal{H}_\infty}.$$

We use notations and concepts of stochastic analysis as explained in the book by Protter [Pro04]. Most of our results only hold for completed filtrations. Since we consider different measures relative to which completions are taken, we use the following notation. Let  $(\mathcal{K}_t)$  be a filtration and  $R$  a probability measure. We denote by  $(\mathcal{K}_t^R)$  the filtration  $(\mathcal{K}_t)$  completed by the  $R$ -negligible sets.

We start with a simple observation.

**Lemma 1.** *If  $\bar{f} : \bar{\Omega} \rightarrow \mathbb{R}$  is  $\bar{\mathcal{F}}_t^{\bar{P}}$ -measurable, then the map  $\bar{f} \circ \psi$  is  $\mathcal{G}_t^P$ -measurable.*

*Proof.* First observe that

$$\begin{aligned}
\mathcal{G}_t &= \bigcap_{s>t} \sigma(A \cap B : A \in \mathcal{F}_s, B \in \mathcal{H}_s) \\
&= \bigcap_{s>t} \sigma(\psi^{-1}(A \times B) : A \in \mathcal{F}_s, B \in \mathcal{H}_s) \\
&= \psi^{-1} \left( \bigcap_{s>t} (\mathcal{F}_s \otimes \mathcal{H}_s) \right) = \psi^{-1}(\bar{\mathcal{F}}_t).
\end{aligned}$$

Now let  $\bar{f} = 1_A$  with  $A \in \bar{\mathcal{F}}_t^{\bar{P}}$ . There is a set  $B \in \bar{\mathcal{F}}_t$  such that  $\bar{P}(A \Delta B) = 0$ . From the first part we deduce that the map  $1_B \circ \psi$  is  $\mathcal{G}_t$ -measurable. Since we have  $P$ -almost surely

$$1_A \circ \psi = 1_B \circ \psi,$$

the map  $1_A \circ \psi$  is  $\mathcal{G}_t^P$ -measurable. By standard arguments one can show the statement for arbitrary  $\bar{\mathcal{F}}_t^{\bar{P}}$ -measurable functions.

**Lemma 2.** *If  $\bar{X}$  is  $(\bar{\mathcal{F}}_t^{\bar{P}})$ -predictable, then  $\bar{X} \circ \psi$  is  $(\mathcal{G}_t^P)$ -predictable.*

*Proof.* Let  $0 < s \leq t$ ,  $A \in \bar{\mathcal{F}}_s^{\bar{P}}$  and

$$\bar{\theta} = 1_A 1_{]s,t]}$$

Then, by Lemma 1,  $\bar{\theta} \circ \psi = (1_A \circ \psi) 1_{]s,t]}$  is  $(\mathcal{G}_t^P)$ -predictable. The proof may be completed by a monotone class argument.

**Lemma 3.** *Let  $\bar{Y}$  be  $(\bar{\mathcal{F}}_t^{\bar{P}})$ -adapted. Then the process*

$$Y = \bar{Y} \circ \psi$$

*is  $(\mathcal{G}_t^P)$ -adapted. Moreover, if  $\bar{Y}$  is a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -local martingale, then  $Y$  is a  $(\mathcal{G}_t^P, P)$ -local martingale.*

*Proof.* The first statement follows immediately from Lemma 1. Now suppose that  $\bar{Y}$  is a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -martingale. Let  $0 \leq s < t$  and  $A \in \mathcal{G}_s$ . Then there is a set  $B \in \bar{\mathcal{F}}_s$  such that  $\psi^{-1}(B) = A$  and hence

$$E^P[1_A(Y_t - Y_s)] = E^{\bar{P}}[1_B(\bar{Y}_t - \bar{Y}_s)] = 0.$$

Thus  $Y$  is a  $(\mathcal{G}_t^P)$ -martingale.

Finally, let  $\bar{Y}$  be a  $(\bar{\mathcal{F}}_t^{\bar{P}})$ -local martingale and  $\bar{T}$  a localizing stopping time. The random time  $T = \bar{T} \circ \psi$  is a  $(\mathcal{G}_t^P)$ -stopping time, since

$$\{T \leq t\} = \psi^{-1}\{\bar{T} \leq t\} \in \psi^{-1}(\bar{\mathcal{F}}_t^{\bar{P}}) \subset \mathcal{G}_t^P.$$

Now it is straightforward to show that  $Y$  is a  $(\mathcal{G}_t^P)$ -local martingale.

**Theorem 1.** *Let  $\bar{Y}$  be a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -semimartingale. Then the process  $Y = \bar{Y} \circ \psi$  is a  $(\mathcal{G}_t^P, P)$ -semimartingale.*

*Proof.* Let  $\bar{Y}$  be a  $(\bar{\mathcal{F}}_t^P)$ -semimartingale and  $Y = \bar{Y} \circ \psi$ . Obviously  $Y$  has cadlag paths  $P$ -a.s. and Lemma 3 implies that  $Y$  is  $(\mathcal{G}_t^P)$ -adapted. By the theorem of Bichteler-Dellacherie-Mokobodzki it is sufficient to show that if  $(\theta^n)$  is a sequence of simple  $(\mathcal{G}_t)$ -adapted integrands converging uniformly to 0, then the simple integrals  $(\theta^n \cdot Y)$  converge to 0 in probability relative to  $P$ . Recall that any  $(\mathcal{G}_t)$ -simple integrand is of the form  $\sum_{1 \leq i \leq n} 1_{]t_i, t_{i+1}]}\theta_i$ , where  $\theta_i$  is  $\mathcal{G}_{t_i}$ -measurable. Since  $\mathcal{G}_t = \psi^{-1}(\bar{\mathcal{F}}_t)$  one can find simple  $(\bar{\mathcal{F}}_t)$ -adapted processes  $(\bar{\theta}^n)$  converging uniformly to 0 such that  $\bar{\theta}^n \circ \psi = \theta^n$ .  $\bar{Y}$  being a semimartingale implies that the sequence  $(\bar{\theta}^n \cdot \bar{Y})$  converges to 0 in probability relative to  $\bar{P}$ , and hence  $(\theta^n \cdot Y)$  converges to 0 in probability relative to  $P$ .

So far we have seen how objects can be translated from  $\bar{\Omega}$  to  $\Omega$ . Now we look at the reverse transfer.

**Lemma 4.** *Let  $M$  be a right-continuous  $(\mathcal{F}_t^P, P)$ -local martingale. Then the process  $\bar{M}(\omega, \omega') = M(\omega)$  is a  $(\bar{\mathcal{F}}_t^Q, \bar{Q})$ -local martingale.*

*Proof.* It is immediate that  $\bar{M}$  is  $(\bar{\mathcal{F}}_t^Q)$ -adapted. For  $0 \leq s < t$  and sets  $A \in \mathcal{F}_s, B \in \mathcal{H}_s$  we have

$$E^{\bar{Q}}[1_A(\omega)1_B(\omega')(\bar{M}_t - \bar{M}_s)] = P(B)E^P[1_A(M_t - M_s)] = 0.$$

By the monotone class theorem, for all bounded  $(\mathcal{F}_s \otimes \mathcal{H}_s)$ -measurable functions  $\theta$  we have

$$E^{\bar{Q}}[\theta(\bar{M}_t - \bar{M}_s)] = 0.$$

Since  $\bar{M}$  is right-continuous, this remains true for all bounded  $\bigcap_{u>s}(\mathcal{F}_u \otimes \mathcal{H}_u)$ -measurable  $\theta$ , and hence  $\bar{M}$  is a martingale with respect to  $(\bar{\mathcal{F}}_t^Q)$ .

Via  $\bar{T}(\omega, \omega') = T(\omega)$  stopping times can be trivially extended to the product space. This finally shows that the local martingale property translates to  $\bar{\Omega}$  with respect to  $\bar{Q}$ .

In the sequel we will always assume that  $\bar{P}$  is absolutely continuous with respect to  $\bar{Q}$ , i.e.

**Assumption 1.**

$$\bar{P} \ll \bar{Q} \text{ on } \bar{\mathcal{F}}.$$

Note that this assumption is always satisfied if  $(\mathcal{G}_t)$  is obtained by an initial enlargement by some discrete random variable  $G$ , i.e.  $\mathcal{H}_t = \sigma(G)$  for all  $t \geq 0$ .

Now let  $M$  be a  $(\mathcal{F}_t^P, P)$ -local martingale and  $\bar{M}$  its extension to  $\bar{\Omega}$  as in Lemma 4. Since  $\bar{P} \ll \bar{Q}$ ,  $\bar{M}$  is a  $(\bar{\mathcal{F}}_t^P, \bar{P})$ -semimartingale and hence, by Theorem 1,  $M$  is a  $(\mathcal{G}_t^P, P)$ -semimartingale. Thus, clearly hypothesis (H') is satisfied. But what is its Doob-Meyer decomposition relative to  $(\mathcal{G}_t^P, P)$ ?

Essentially the change of filtrations corresponds to changing the measure from  $\bar{Q}$  to  $\bar{P}$  on the product space  $\bar{\Omega}$ . Girsanov's theorem applies on  $\bar{\Omega}$ , since the measure  $\bar{P}$  is absolutely continuous with respect to  $\bar{Q}$ . As a consequence we obtain a Girsanov-type result for the corresponding change of filtrations.

For its explicit description we introduce the density process. Let  $(\bar{Z}_t)$  denote a cadlag  $(\bar{\mathcal{F}}_t^{\bar{Q}})$ -adapted process with

$$\bar{Z}_t = \frac{d\bar{P}}{d\bar{Q}} \Big|_{\bar{\mathcal{F}}_t^{\bar{Q}}}.$$

Note that we need to consider the completed filtration in order to assure the existence of a cadlag density process. Theorem 1 implies that the process  $Z$  defined by

$$Z = \bar{Z} \circ \psi$$

is a  $(\mathcal{G}_t^P, P)$ -semimartingale. Before giving the Girsanov-type results, we show how the quadratic variation processes behave under the projection  $\psi$ .

**Lemma 5.** *Let  $\bar{X}$  and  $\bar{Y}$  be  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -semimartingales. If  $X = \bar{X} \circ \psi$  and  $Y = \bar{Y} \circ \psi$ , then*

$$[\bar{X}, \bar{Y}] \circ \psi = [X, Y]$$

up to indistinguishability relative to  $P$ .

*Proof.* Put  $X = \bar{X} \circ \psi$  and  $Y = \bar{Y} \circ \psi$ . Let  $t > 0$  and  $t_i^n = t \frac{i}{2^n}$  for all  $i = 0, 1, \dots, 2^n$ . It is known that the sums

$$\bar{X}_0 \bar{Y}_0 + \sum_{0 \leq i < 2^n} (\bar{X}_{t_{i+1}^n} - \bar{X}_{t_i^n})(\bar{Y}_{t_{i+1}^n} - \bar{Y}_{t_i^n})$$

converge to  $[\bar{X}, \bar{Y}]_t$  in probability relative to  $\bar{P}$  (see Theorem 20, Chapter VIII in [DM82]). Hence  $[\bar{X}, \bar{Y}]_t \circ \psi$  is the limit (in probability) of the sums

$$X_0 Y_0 + \sum_{0 \leq i < 2^n} (X_{t_{i+1}^n} - X_{t_i^n})(Y_{t_{i+1}^n} - Y_{t_i^n})$$

relative to  $P$ . Obviously the limit is also equal to  $[X, Y]_t$ , and hence we have

$$[\bar{X}, \bar{Y}]_t \circ \psi = [X, Y]_t.$$

Since both processes are cadlag, they coincide up to indistinguishability relative to  $P$ .

Let  $\bar{M}$  be a  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -semimartingale and  $M = \bar{M} \circ \psi$ . Since  $\bar{P}$  is absolutely continuous with respect to  $\bar{Q}$ ,  $\bar{M}$  is also a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -semimartingale. Moreover, the bracket  $[\bar{M}, \bar{Z}]$  relative to  $\bar{Q}$  is  $\bar{P}$ -indistinguishable from the bracket relative to  $\bar{P}$ . Similarly, Lemma 5 implies that the bracket  $[M, Z]$  of the  $(\mathcal{G}_t^P, P)$ -semimartingales  $M$  and  $Z$  coincides with  $[\bar{M}, \bar{Z}] \circ \psi$ .

We are now in a position to state the first Girsanov-type result. We begin with some definitions. Let

$$\bar{T} = \inf\{t > 0 : \bar{Z}_t = 0, \bar{Z}_{t-} > 0\}$$

and  $\bar{U}_t = \Delta \bar{M}_{\bar{T}} 1_{\{t \geq \bar{T}\}}$ . We further denote by  $\tilde{U}$  the compensator of  $\bar{U}$ , i.e. the  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -predictable projection of  $\bar{U}$ . Moreover, we will use the abbreviation  $\hat{U} = \tilde{U} \circ \psi$ .

**Theorem 2.** *If  $M$  is a  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ , then*

$$M - \frac{1}{Z} \cdot [M, Z] + \hat{U} \quad (2)$$

*is a  $(\mathcal{G}_t^P, P)$ -local martingale.*

*Proof.* Let  $M$  be an  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ . We may assume that  $M$  has cadlag paths. Lemma 4 implies that the process defined by

$$\bar{M}(\omega, \omega') = M(\omega)$$

is a  $(\bar{\mathcal{F}}_t^{\bar{Q}})$ -local martingale and the Lenglart-Girsanov Theorem yields that

$$\bar{M} - \frac{1}{\bar{Z}} \cdot [\bar{M}, \bar{Z}] + \tilde{U}_t$$

is a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -local martingale (see Théorème 3 in [Len77] or Chapter III in [Pro04]). Since the bracket process  $[\bar{M}, \bar{Z}] \circ \psi$  is  $P$ -indistinguishable from  $[M, Z]$  (see Lemma 5), we have

$$\left(\frac{1}{\bar{Z}} \cdot [\bar{M}, \bar{Z}]\right) \circ \psi = \frac{1}{Z} \cdot [M, Z]$$

up to indistinguishability. With Lemma 3 we conclude that

$$M - \frac{1}{Z} \cdot [M, Z] + \hat{U}_t$$

is a  $(\mathcal{G}_t^P, P)$ -local martingale.

In case  $M$  is continuous, the preceding decomposition in the larger filtration simplifies.

**Theorem 3.** *If  $M$  is a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ , then*

$$M - \frac{1}{Z} \cdot [M, Z]$$

*is a  $(\mathcal{G}_t^P, P)$ -local martingale.*

*Proof.* Let  $M$  be a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$  and put  $\bar{M}(\omega, \omega') = M(\omega)$ . The related process  $\bar{U}$  vanishes, and hence  $\hat{U}$  vanishes as well. The result follows now from Theorem 2.

The preceding may also be formulated in terms of the stochastic logarithm of the density process  $\bar{Z}$ . To this end set  $\bar{S} = \inf\{t > 0 : \bar{Z}_t = 0, \Delta\bar{Z}_t = 0\}$  and define

$$\bar{L} = \int_{0+}^{\cdot} \frac{1}{\bar{Z}_-} d\bar{Z} \quad \text{on } [0, \bar{S}[. \quad (3)$$

So far, the process  $\bar{L}$  is determined  $\bar{P}$ -, but *not*  $\bar{Q}$ -almost everywhere. (In order to define it everywhere we may put  $\bar{L} = 0$  on  $[\bar{S}, \infty[$ .) Then  $\bar{L}$  is an  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -semimartingale but not necessarily an  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -semimartingale. However, restricted to the time interval  $[0, \bar{S}[$  it is an  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale. As usual

we write  $L = \bar{L} \circ \psi$ . Alternatively, one can define  $L$  through the stochastic integral

$$L = \int_{0+}^{\cdot} \frac{1}{Z_-} dZ.$$

Since the process  $\bar{L}$  is a  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale on the interval  $[0, \bar{S}]$ , it can be decomposed into a unique local-martingale part  $\bar{L}^c$  and a sum of compensated jumps  $\bar{L}^d$ . As before, we consider the processes  $L^c = \bar{L}^c \circ \psi$  and  $L^d = \bar{L}^d \circ \psi$ .

Theorem 3 can now be reformulated as follows.

**Theorem 4.** *If  $M$  is a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ , then*

$$M - [M, L]$$

*is a  $(\mathcal{G}_t^P, P)$ -local martingale.*

*Proof.* Let  $M$  be a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ . Since  $M$  is continuous, the bracket process  $[M, Z]$  is continuous and Theorem 3 implies that

$$M - \frac{1}{Z} \cdot [M, Z] = M - \frac{1}{Z_-} \cdot [M, Z]$$

is a  $(\mathcal{G}_t^P, P)$ -local martingale. Moreover, the definition of  $L$  implies that  $\frac{1}{Z_-} \cdot [M, Z] = [M, L]$ ,  $P$ -a.s., so that  $M - [M, L]$  is a  $(\mathcal{G}_t^P, P)$ -local martingale.

Finally, we will need the following formula, in which the subtracted drift is represented in terms of the quadrature variation of the given local martingale.

**Theorem 5.** *If  $M$  is a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ , then there is a  $(\mathcal{G}_t^P)$ -predictable process  $\alpha$  such that  $P$ -a.s.*

$$\int_0^\infty \alpha_t^2 d[M, M]_t \leq [L, L]_\infty^c < \infty,$$

*and*

$$M - \alpha \cdot [M, M]$$

*is a  $(\mathcal{G}_t^P)$ -local martingale.*

*Proof.* Let  $M$  be a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ . By the Kunita-Watanabe Inequality one has for  $0 \leq s < t$ ,

$$[M, L]_t - [M, L]_s \leq [L, L]_t^{1/2} ([M, M]_t - [M, M]_s)^{1/2}.$$

Since  $[L, L]_t$  is finite for all  $t \geq 0$ , the measure  $d[M, L]$  is absolutely continuous with respect to  $d[M, M]$  and there exists a  $(\mathcal{G}_t^P)$ -predictable process  $\alpha$  with

$$\alpha \cdot [M, M] = [M, L] = [M, L^c]$$

(see Lemme 1.36 in [Jac79]). Moreover, the processes  $M$  and  $O = L^c - \alpha \cdot M$  are orthogonal w.r.t.  $[\cdot, \cdot]$ . Consequently,



$$\alpha^2 \cdot [M, M] = [\alpha \cdot M, \alpha \cdot M] \leq [L^c, L^c] = [L, L]^c.$$

Recall that

$$[L, L] = \left( \frac{1}{\bar{Z}_-^2} \cdot [\bar{Z}, \bar{Z}] \right) \circ \psi$$

and that  $\bar{Z}$  is a uniformly integrable nonnegative  $(\bar{\mathcal{F}}_t^Q, \bar{Q})$ -martingale. Since  $\bar{P}$ -a.s.  $\bar{Z}_\infty > 0$ , one has also  $\inf_{t \geq 0} \bar{Z}_t > 0$ ,  $\bar{P}$ -a.s. Moreover,  $[\bar{Z}, \bar{Z}]_\infty < \infty$ ,  $\bar{Q}$ -a.s. Therefore,  $[\bar{L}, \bar{L}]$  is  $\bar{P}$ -a.s. bounded and consequently  $[L, L]^c_t$  converges as  $t \rightarrow \infty$   $P$ -a.s. to some real value which we denote by  $[L, L]^c_\infty$ .

*Remark 1.* In the literature the process  $\alpha$  is sometimes called *information drift* (see for example [Imk03]).

*Remark 2.* Due to the previous theorem the information drift obtained via the Girsanov approach under Assumption 1 is always locally square integrable. It was shown in [ADI04] that in case  $\Omega$  is standard Borel and each  $(\mathcal{F}_t)$ -martingale has a continuous modification, square integrability on the product space  $\Omega \times [0, T]$  with respect to the measure  $d[M, M] \otimes P$  implies the absolute continuity of the kernels  $k_t(\cdot, d\omega')$  with respect to the conditional laws  $P_t(\cdot, d\omega')$  of  $\mathcal{G}_{t-}$  with respect to  $\mathcal{F}_t$ , where

$$P_t(\cdot, A) = P(A) + \int_0^t k_s(\cdot, A) dM_s + L_t^A, \quad A \in \mathcal{G}_{t-}, t \in [0, T],$$

$L^A$  being orthogonal to  $M$ . In this case the R-N density process  $\gamma_t(\omega, \omega') = \frac{k_t(\cdot, d\omega')}{P_t(\cdot, d\omega')}$  is identical to  $\alpha$  if restricted to the diagonal  $\omega = \omega'$ . Hence this absolute continuity condition (*ACL*) is implied by Assumption 1. Enlargements with locally integrable but not square integrable information drifts are beyond the scope of this article. But they provide examples for which (*ACL*) does not imply Assumption 1. One example is obtained for instance by enlarging the Wiener filtration by the maximum of the Wiener process over some finite time interval. In this case Malliavin's calculus can be applied and an explicit representation of the information drift is obtained via the Clark-Ocone formula (see [IPW01] and [Imk03]). In case  $\Omega$  is not standard Borel we do not know at the moment whether 1 is more restrictive than (*ACL*). The methods of [ADI04] allow in a more general setting the description of information drifts which are not necessarily locally square integrable.

### Comparison with Jacod's condition

In Jacod's paper (see [Jac85]) the filtration  $(\mathcal{F}_t)$  is supposed to be enlarged by some random variable  $G$  taking values in a Lusin space  $(E, \mathcal{E})$ . As a consequence, for  $t \in [0, T]$  regular conditional distributions  $Q_t$  of  $G$  relative to  $\mathcal{F}_t$  exist. The following condition is assumed to be satisfied:

(A') For every  $t \geq 0$  and  $P$ -a.a.  $\omega$  the measure  $Q_t(\omega, \cdot)$  is absolutely continuous with respect to the law  $\eta$  of  $G$ .

We will show that condition (A') implies our assumption 1. More precisely, with  $\mathcal{H}_t = \sigma(G)$ , we have the following.

**Lemma 6.** *If (A') is satisfied, then  $\bar{P} \ll \bar{Q}$  on  $\bar{\mathcal{F}}_s$  for all  $s \geq 0$ .*

*Proof.* Let  $t \geq 0$ ,  $A \in \mathcal{F}_t$  and  $B \in \mathcal{E}$ . We put

$$\tilde{P}(A \times G^{-1}[B]) = \int_A Q_t(\omega, B) dP(\omega),$$

and extend  $\tilde{P}$  to a probability measure on  $\mathcal{F}_t \otimes \sigma(G)$ . Note that for  $A \in \mathcal{F}_t$  and  $B \in \mathcal{E}$

$$\tilde{P}(A \times G^{-1}[B]) = P(A \cap G^{-1}[B]) = \bar{P}(A \times G^{-1}B),$$

and hence  $\tilde{P} = \bar{P}$  on  $\mathcal{F}_t \otimes \sigma(G)$ . Now let  $s < t$  and  $C \in \bar{\mathcal{F}}_s$  with  $\bar{P}(C) = 0$ . We claim that  $\bar{Q}(C) = 0$ .

Notice that also  $\tilde{P}(C) = 0$ . Now choose a set  $D \in \mathcal{F}_t \otimes \mathcal{E}$  such that  $C$  is the inverse image of  $D$  under the map  $(\omega, \omega') \mapsto (\omega, G(\omega'))$ . Then  $\int 1_D(\omega, x) Q_t(\omega, dx) = 0$  for  $P$ -a.a.  $\omega$ . With assumption (A') we conclude that  $\int 1_D(\omega, x) d\eta(x) = 0$  for  $P$ -a.a.  $\omega$ , and hence

$$\begin{aligned} \bar{Q}(C) &= \int \int 1_C(\omega, \omega') dP(\omega') dP(\omega) \\ &= \int \int 1_D(\omega, x) d\eta(x) dP(\omega) \\ &= 0. \end{aligned}$$

Thus we have shown the result.

Jacod does not use Girsanov's theorem in his paper [Jac85]. However, he points out that his results could also be deduced by applying it to the conditional measures  $P^x = P(\cdot | G = x)$ ,  $x \in E$ . Condition (A') implies that the conditional measures  $P^x$  are absolutely continuous with respect to  $P$ . Hence, by Girsanov, for a given  $(\mathcal{F}_t, P)$ -local martingale there is a drift  $A^x$  such that  $M - A^x$  is a  $(\mathcal{F}_t, P^x)$ -local martingale. By combining the processes  $A^x$  we obtain that

$$M - A^G$$

is a  $(\mathcal{G}_t, P)$ -local martingale. The main work consists in proving that the processes  $A^x$  can be combined in a meaningful way. As far as we know, Jacod's sketch has never been worked out rigorously.

In our approach we embed every local martingale into the product space  $\bar{\Omega}$ . We apply Girsanov's theorem on the product space and then translate our results back into the original space. One of the advantages of our approach is that we do not have to assume regular conditional distributions to exist. And we do not need to show how processes can be combined. Instead we have to show how one can transfer objects from  $\Omega$  to  $\bar{\Omega}$  and vice versa. Moreover we are not restricted to initial enlargements, but only to enlargements of the form

$$\mathcal{G}_t = \bigcap_{s>t} (\mathcal{F}_s \vee \mathcal{H}_s), \quad t \in [0, T].$$

Starting with Jacod's results one can obtain decompositions for filtrations of this kind by using predictable projections. For this suppose  $A$  to be a bounded variation process such that  $M - A$  is a local martingale with respect to the initially enlarged filtration  $(\mathcal{F}_t \vee \mathcal{H}_\infty)$ . If  $B$  is the predictable projection of  $A$  onto  $(\mathcal{G}_t)$ , then  $M - B$  is a  $(\mathcal{G}_t)$ -local martingale.

## 2 Estimates for the drift

Suppose  $M$  is a continuous  $(\mathcal{F}_t^P, P)$ -local martingale with  $M_0 = 0$ . Under the assumptions of the previous section we know that there is a  $(\mathcal{G}_t^P)$ -predictable process  $\alpha$  such that  $M - \alpha \cdot [M, M]$  is  $(\mathcal{G}_t^P, P)$ -local martingale. Moreover, the information drift  $\alpha$  satisfies

$$(\alpha^2 \cdot [M, M])_\infty \leq [L, L]_\infty^c. \quad (4)$$

In this section we provide bounds for

$$E [(\alpha^2 \cdot [M, M])_\infty^p]$$

for various moments  $p \geq 1$  based on inequality (4).

Throughout this section we suppose the assumptions of the previous section and maintain the notation. More precisely, we assume that  $\bar{P} \ll \bar{Q}$ , denote by  $\bar{Z}_t = \frac{d\bar{P}}{d\bar{Q}} \Big|_{\mathcal{F}_t^{\bar{Q}}}$  the density process, and by  $\bar{L}$  the stochastic logarithm of  $\bar{Z}$ . We use again the decomposition of  $\bar{L}$  into a continuous part  $\bar{L}^c$  and a part  $\bar{L}^d$  consisting of compensated jumps. As before we denote by  $Z, L$  and  $L^c$  the corresponding  $(\mathcal{G}_t)$ -adapted processes obtained by a right side application of  $\psi$ .

### 2.1 Moment $p = 1$

Recall that the relative entropy of two probability measures  $P$  and  $Q$  on some  $\sigma$ -algebra  $\mathcal{M}$  is defined by

$$H_{\mathcal{M}}(P \| Q) = \begin{cases} E^P \left( \log \frac{dP}{dQ} \Big|_{\mathcal{M}} \right), & \text{if } P \ll Q \text{ on } \mathcal{M} \\ \infty, & \text{if not } P \ll Q \text{ on } \mathcal{M}. \end{cases}$$

In our situation, the relative entropy  $H_{\mathcal{F}_\infty}(\bar{P} \| \bar{Q})$  provides an upper bound for the first moment of  $[L, L]_\infty^c$ :

**Lemma 7.**

$$\frac{1}{2} E^P [L, L]_\infty^c \leq H_{\mathcal{F}_\infty}(\bar{P} \| \bar{Q}).$$

If  $(\bar{Z}_t)_{t \geq 0}$  is continuous and  $\bar{Z}_0 = 1$ , then one even has

$$\frac{1}{2} E^P [L, L]_\infty = H_{\mathcal{F}_\infty}(\bar{P} \| \bar{Q}).$$

*Remark 3.* If the  $\sigma$ -field  $\mathcal{F}_0$  is trivial, then the measures  $\bar{P}$  and  $\bar{Q}$  coincide on  $\mathcal{F}_0 \otimes \mathcal{H}_0$ . As will be shown in Lemma 14 we have  $\bar{\mathcal{F}}_0^{\bar{Q}} = (\mathcal{F}_0 \otimes \mathcal{H}_0)^{\bar{Q}}$ , and hence in this case  $\bar{Z}_0 = 1$ .

*Proof.* To prove the first statement, we decompose  $\bar{L}$  into its continuous and discontinuous part  $\bar{L} = \bar{L}^c + \bar{L}^d$  and let  $\bar{Z}_t^c = \mathcal{E}(\bar{L}^c)_t$  and  $\bar{Z}_t^d = \bar{Z}_0 \mathcal{E}(\bar{L}^d)_t$ . Then  $\bar{Z}_t = \bar{Z}_t^c \bar{Z}_t^d$  on  $[0, \bar{S}[$ . The following results are only valid when stopping all processes at a stopping time  $\bar{T} = \inf\{t > 0 : \bar{Z}_t < \varepsilon\}$  for some  $\varepsilon > 0$ . To simplify notation, we omit the stopping times in the following computations. One has

$$\begin{aligned} \log \bar{Z}_t &= \log(\mathcal{E}(\bar{L}^c))_t + \log(\bar{Z}_t^d) \\ &= (\bar{L}_t^c - [\bar{L}, \bar{L}]_t^c) + \frac{1}{2}[\bar{L}, \bar{L}]_t^c + \log \bar{Z}_t^d, \end{aligned}$$

where the term in the first brackets is a  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -local martingale due to Girsanov's theorem. Consider the function  $\xi(x) = x \log x$  ( $x \in [0, \infty)$ ) and denote  $\bar{A}_t = \bar{Z}_t \log \bar{Z}_t^d = \bar{Z}_t^c \xi(\bar{Z}_t^d)$ . Then Itô's formula yields

$$\begin{aligned} \bar{A}_t &= \xi(\bar{Z}_0) + \int_{0+}^t \xi(\bar{Z}_{s-}^d) d\bar{Z}_s^c + \int_{0+}^t \bar{Z}_{s-}^c \xi'(\bar{Z}_{s-}^d) d\bar{Z}_s^d \\ &\quad + \sum_{0 < s \leq t} \bar{Z}_{s-}^c (\xi(\bar{Z}_s^d) - \xi(\bar{Z}_{s-}^d) - \xi'(\bar{Z}_{s-}^d) \Delta \bar{Z}_s^d), \end{aligned}$$

where all summands in the previous line are non-negative due to the convexity of  $\xi$ .

Let now  $(\bar{T}^n)$  denote a  $\bar{P}$ -localizing sequence of bounded stopping times such that the integrals in the stopped processes  $(\bar{A}_t^{\bar{T}^n})$  are  $\bar{Q}$ -submartingales and the stopped processes  $(\bar{L}^c - [\bar{L}, \bar{L}]_t^c)^{\bar{T}^n}$  are  $\bar{P}$ -martingales. Then

$$E^{\bar{Q}}[\bar{A}_{\bar{T}^n}] \geq E^{\bar{Q}}[\xi(\bar{Z}_0)] \geq 0,$$

which leads to

$$\begin{aligned} E^{\bar{Q}}[\xi(\bar{Z}_{\bar{T}^n})] &\geq E^{\bar{Q}}[\bar{Z}_{\bar{T}^n} \log \bar{Z}_{\bar{T}^n}^c] = E^{\bar{P}}[\log \bar{Z}_{\bar{T}^n}^c] \\ &= \frac{1}{2} E^{\bar{P}}[\bar{L}, \bar{L}]_{\bar{T}^n}^c. \end{aligned} \tag{5}$$

Note that by Jensen's inequality for all stopping times  $T$

$$E^{\bar{Q}}[\xi(\bar{Z}_\infty) | \bar{\mathcal{F}}_T] \geq \xi \left( E^{\bar{Q}}[\bar{Z}_\infty | \bar{\mathcal{F}}_T] \right) = \xi(\bar{Z}_T),$$

from which we deduce that  $E^{\bar{Q}}[\xi(\bar{Z}_{\bar{T}^n})] \leq E^{\bar{Q}}[\xi(\bar{Z}_\infty)] = H_{\bar{\mathcal{F}}_\infty}(\bar{P} \| \bar{Q})$ . With (5) we arrive at

$$\frac{1}{2} E^{\bar{P}}[\bar{L}, \bar{L}]_{\bar{T}^n}^c \leq H_{\bar{\mathcal{F}}_\infty}(\bar{P} \| \bar{Q})$$

and monotone convergence implies that  $\frac{1}{2} E^{\bar{P}}[\bar{L}, \bar{L}]_\infty^c \leq H_{\bar{\mathcal{F}}_\infty}(\bar{P} \| \bar{Q})$ .

It remains to show that in case  $(\bar{Z}_t)$  is continuous with  $\bar{Z}_0 = 1$  and  $E^{\bar{P}}[\bar{L}, \bar{L}]_\infty < \infty$ , we have  $\frac{1}{2} E^{\bar{P}}[\bar{L}, \bar{L}]_\infty \geq H_{\bar{\mathcal{F}}_\infty}(\bar{P} \| \bar{Q})$ . Indeed, then  $(\bar{L}_t - [\bar{L}, \bar{L}]_t)$  is an  $L^2$ -bounded  $(\bar{\mathcal{F}}_t^{\bar{P}}, \bar{P})$ -martingale and one has

$$\begin{aligned} \frac{1}{2}E^{\bar{P}}[\bar{L}, \bar{L}]_t &= E^{\bar{P}}\left[\bar{L}_t - [\bar{L}, \bar{L}]_t + \frac{1}{2}[\bar{L}, \bar{L}]_t\right] \\ &= E^{\bar{P}}\log \bar{Z}_t = E^{\bar{Q}}[\bar{Z}_t \log \bar{Z}_t]. \end{aligned}$$

The left hand side converges to  $\frac{1}{2}E^{\bar{P}}[\bar{L}, \bar{L}]_\infty$  as  $t \rightarrow \infty$ . On the other hand,  $\xi(\bar{Z}_t)$  converges to  $\xi(\bar{Z}_\infty)$  so that by Fatou's lemma

$$\liminf_{t \rightarrow \infty} E^{\bar{Q}}[\bar{Z}_t \log \bar{Z}_t] \geq E^{\bar{Q}}\xi(\bar{Z}_\infty) = H_{\mathcal{F}_\infty}(\bar{P} \parallel \bar{Q}).$$

This completes the proof.

## 2.2 Moments $p > 1$

Now we consider moments of order  $p > 1$ . In this case the  $p$ -th moment of  $[L, L]_\infty$  can be compared to some generalized relative entropy. See [Imk96] for elementary versions of the inequalities to be derived.

Our analysis requires some additional assumption. We suppose that  $(\mathcal{G}_t)$  is an initial enlargement of  $(\mathcal{F}_t)$ , i.e.

$$\mathcal{G}_t = \bigcap_{s>t} (\mathcal{F}_s \vee \mathcal{A}), \quad t \geq 0,$$

where  $\mathcal{A}$  is some fixed sub- $\sigma$ -algebra of  $\mathcal{F}$ . Moreover, we assume that  $\mathcal{F}_0$  is trivial. Additionally, we need to make the following assumption.

**Assumption 2 (C).** *Every  $(\mathcal{F}_t^P, P)$ -martingale has a continuous modification.*

We shall see that under this condition  $\bar{L}$  is a continuous  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale.

We begin with the definition of the generalized relative entropy.

**Definition 1.** *For  $p > 1$ , and probability measures  $P \ll Q$  on a  $\sigma$ -algebra  $\mathcal{M}$ , let*

$$H_{\mathcal{M}}^p(P \parallel Q) := E^P\left(\log_+ \frac{dP}{dQ} \Big|_{\mathcal{M}}\right)^p.$$

We provide now an upper bound of  $E[L, L]_\infty^p$  with the help of the generalized entropy of  $\bar{P}$  with respect to  $\bar{Q}$  on the set  $\bar{\mathcal{F}}_\infty$ . To simplify notations, we omit the  $\sigma$ -algebra  $\bar{\mathcal{F}}_\infty$ , and write only  $H^p(\bar{P} \parallel \bar{Q})$  and  $H(\bar{P} \parallel \bar{Q})$ . The aim of this section is to prove

**Theorem 6.** *For any  $p \geq 1$  there exists a universal constant  $C = C(p) < \infty$  such that under the above assumptions one has*

$$E[L, L]_\infty^p \leq C [H(\bar{P} \parallel \bar{Q}) + H^p(\bar{P} \parallel \bar{Q})].$$

For the proof we need some auxiliary results. We start by showing that there exists a continuous modification for  $\bar{Z}$ .

**Lemma 8.** *Let  $\bar{M}$  be a uniformly integrable  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale. If assumption (C) is satisfied, then for  $P$ -a.a.  $\omega'$  the process  $\bar{M}^{\omega'} = \bar{M}(\cdot, \omega')$  is a  $(\mathcal{F}_t^P)$ -local martingale.*

*Proof.* Choose a modification such that every path of  $\bar{M}$  is cadlag. Now let  $\hat{M}$  be an  $\mathcal{A} \otimes \mathcal{O}(\mathcal{F})$ -measurable process such that for all  $\omega'$  and  $s \geq 0$

$$\hat{M}_s^{\omega'} = E^P[\bar{M}_\infty^{\omega'} | \mathcal{F}_s].$$

For the existence of such a process we refer to [SY78], Proposition 3. Put  $C_t = \{\hat{M}_t > \bar{M}_t\}$ . Clearly  $C_t \in \bar{\mathcal{F}}_t^{\bar{Q}}$  and  $C_t(\cdot, \omega') \in \mathcal{F}_t^P$  for all  $P$ -a.a.  $\omega'$  (recall that  $(\mathcal{F}_t)$  is right-continuous). Moreover for  $t \geq 0$

$$\begin{aligned} & \int \int 1_{C_t}(\omega, \omega') (\hat{M}_t^{\omega'} - \bar{M}_t^{\omega'}) dP(\omega) dP(\omega') \\ &= E^{\bar{Q}}[1_{C_t}(\hat{M}_t - \bar{M}_t)] \\ &= E^{\bar{Q}}[1_{C_t}(\hat{M}_t - \bar{M}_\infty)] \\ &= \int \int 1_{C_t}(\omega, \omega') (\hat{M}_t^{\omega'} - \bar{M}_\infty^{\omega'}) dP(\omega) dP(\omega') \\ &= \int 0 dP(\omega') = 0, \end{aligned}$$

A similar result holds true on the set  $\{\hat{M}_t < \bar{M}_t\}$ , and as a consequence we have for  $P$ -a.a.  $\omega'$

$$\hat{M}_t(\cdot, \omega') = \bar{M}_t(\cdot, \omega'), \quad P\text{-a.s.}$$

Hence for  $P$ -a.a.  $\omega'$  the process  $(\bar{M}_q^{\omega'})_{q \in \mathbb{Q}^+}$  is a  $(\mathcal{F}_t^P)$ -martingale. Since  $\bar{M}_t$  is cadlag and uniformly integrable we obtain that also

$$(\bar{M}_t^{\omega'})_{t \geq 0}$$

is a  $(\mathcal{F}_t^P)$ -martingale for  $P$ -a.a.  $\omega'$ .

**Lemma 9.** *If (C) is satisfied, then every uniformly integrable  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale has a continuous modification.*

*Proof.* Let  $\bar{M}$  be a  $(\bar{\mathcal{F}}_t^{\bar{Q}}, \bar{Q})$ -local martingale. We may suppose that  $\bar{M}$  is cadlag everywhere, and hence, the set

$$N = \{(\omega, \omega') : t \mapsto \bar{M}_t(\omega, \omega') \text{ is not continuous}\}$$

is measurable. Fix  $\omega'$  and suppose that  $\bar{M}^{\omega'}$  is a  $(\mathcal{F}_t^P)$ -martingale. Then assumption (C) implies that for  $P$ -a.a.  $\omega$  the paths  $t \mapsto \bar{M}_t^{\omega'}(\omega)$  are continuous, i.e.  $P(N^{\omega'}) = 0$ . Now Fubini's Theorem yields with Lemma 8

$$\begin{aligned} E^{\bar{Q}}(N) &= \int \int 1_{N^{\omega'}}(\omega) dP(\omega) dP(\omega') \\ &= \int 0 dP(\omega') = 0, \end{aligned}$$

and hence the result.

For the rest of the section we will suppose that  $\bar{Z}$  is a continuous modification of our density process  $\left. \frac{d\bar{P}}{d\bar{Q}} \right|_{\mathcal{F}_t^{\bar{Q}}}$ . Similarly,  $\bar{L}$  will be assumed to be continuous.

*Proof ( of Theorem 6).* We assume that  $H(\bar{P}||\bar{Q})$  and  $H^p(\bar{P}||\bar{Q})$  are finite. Then  $X_t := \bar{L}_t - [\bar{L}, \bar{L}]_t$  is a continuous  $L^2$ -bounded  $\bar{P}$ -martingale by Lemma 7 and we write  $\log \bar{Z}_t = X_t + \frac{1}{2}A_t$  with  $A_t := [\bar{L}, \bar{L}]_t = [X, X]_t$ . Next, observe that

$$\begin{aligned} H^p(\bar{P}||\bar{Q})^{1/p} &= E^{\bar{P}} \left[ (X_\infty + \frac{1}{2}A_\infty)_+^p \right]^{1/p} \\ &\geq E^{\bar{P}} \left[ \left( \frac{1}{2}A_\infty - (|X_\infty| \wedge \frac{1}{2}A_\infty) \right)^p \right]^{1/p} \\ &\geq \frac{1}{2} E^{\bar{P}} [A_\infty^p]^{1/p} - E^{\bar{P}} [|X_\infty|^p]^{1/p} \\ &\geq \frac{1}{2} E^{\bar{P}} [A_\infty^p]^{1/p} - C E^{\bar{P}} [A_\infty^{p/2}]^{1/p}, \end{aligned} \tag{6}$$

where the last inequality holds for some constant  $C > 0$  due to the Burkholder-Davis-Gundy inequality. Now choose  $\xi > 0$  such that for all  $x \geq 0$

$$C^p x^{p/2} \leq \xi^p x + \frac{1}{4^p} x^p.$$

This leads to

$$C^p E^{\bar{P}} A_\infty^{p/2} \leq \xi^p E^{\bar{P}} A_\infty + \frac{1}{4^p} E^{\bar{P}} A_\infty^p$$

and hence to

$$C E^{\bar{P}} [A_\infty^{p/2}]^{1/p} \leq \xi E^{\bar{P}} [A_\infty]^{1/p} + \frac{1}{4} E^{\bar{P}} [A_\infty^p]^{1/p}.$$

With (6) we conclude that

$$H^p(\bar{P}||\bar{Q})^{1/p} \geq \frac{1}{4} E^{\bar{P}} [A_\infty^p]^{1/p} - \xi E^{\bar{P}} [A_\infty]^{1/p} = \frac{1}{4} E^{\bar{P}} [A_\infty^p]^{1/p} - \xi H(\bar{P}||\bar{Q})^{1/p}.$$

Consequently,

$$\begin{aligned} E^{\bar{P}} [A_\infty^p]^{1/p} &\leq 4\xi H(\bar{P}||\bar{Q})^{1/p} + 4H^p(\bar{P}||\bar{Q})^{1/p} \\ &\leq 8(\xi^p H(\bar{P}||\bar{Q}) + H^p(\bar{P}||\bar{Q}))^{1/p}, \end{aligned}$$

where the last step follows from the elementary inequality  $a+b \leq 2(a^p+b^p)^{1/p}$ ,  $a, b \geq 0$ .

*Remark 4.* The above proof is based on the fact that there exists a constant  $C_p$  such that for any continuous  $L^2$ -bounded  $\bar{P}$ -martingale  $(X_t)$  with  $X_0 = 0$  and quadratic variation process  $(A_t)$  one has

$$E^{\bar{P}} A_\infty^p \leq C_p E^{\bar{P}} \left[ X_\infty + \frac{1}{2}A_\infty + (X_\infty + \frac{1}{2}A_\infty)_+^p \right].$$

Improving the estimate to

$$E^{\bar{P}} A_{\infty}^p \leq C_p E^{\bar{P}} \left( X_{\infty} + \frac{1}{2} A_{\infty} \right)_+^p \quad (7)$$

would lead to the better estimate  $E^{\bar{P}} [L, L]^p \leq C_p H^p(\bar{P} \parallel \bar{Q})$ . However, an estimate stating (7) is not valid, as the following example shows.

*Example 1.* Let  $W$  be a Wiener process and for fixed  $\varepsilon > 0$ , let  $T$  denote the first hitting time of the slope  $t \mapsto \varepsilon - t/2$ . We consider  $X_t := W_t^T$  and  $A_t := [X, X]_t$ . Then by the Lévy-Bachelier formula the law of  $T = A_{\infty}$  has density

$$1_{(0, \infty)}(t) \frac{\varepsilon}{t^{3/2}} \phi\left(\frac{\varepsilon - t/2}{\sqrt{t}}\right),$$

where  $\phi$  is the density of the standard normal law. Hence,

$$E[A_{\infty}^p] = \varepsilon \int_0^{\infty} t^{p-3/2} \phi\left(\frac{\varepsilon - t/2}{\sqrt{t}}\right) dt.$$

In particular, for  $\varepsilon \downarrow 0$ , one has  $E[A_{\infty}^p] \approx \varepsilon$ . On the other hand,

$$E\left[\left(X_{\infty} + \frac{1}{2} A_{\infty}\right)_+^p\right] = E[(W_T + T/2)^p] = \varepsilon^p$$

such that one can always find a sufficiently small  $\varepsilon > 0$  for which the inequality (7) is not valid.

We next show a result which in a sense contains the inverse statement to Theorem 6.

**Lemma 10.** *For  $p \geq 1$  there exists a universal constant  $C = C(p) < \infty$  such that*

$$H^p(\bar{P} \parallel \bar{Q}) \leq C [E^{\bar{P}}[\bar{L}, \bar{L}]_{\infty}^p + 1].$$

*In particular finiteness of  $E^{\bar{P}}([\bar{L}, \bar{L}]_{\infty}^p)$  implies finiteness of the entropy  $H^p(\bar{P} \parallel \bar{Q})$ .*

*Proof.* We have, by Burkholder-Davis-Gundy, with a universal constant  $C_1$

$$\begin{aligned} H^p(\bar{P} \parallel \bar{Q})^{1/p} &\leq E\left(|\bar{L}_{\infty} - \frac{1}{2}[\bar{L}, \bar{L}]_{\infty}|^p\right)^{1/p} \\ &\leq E(|\bar{L}_{\infty}|^p)^{1/p} + E\left(\frac{1}{2}[\bar{L}, \bar{L}]_{\infty}^p\right)^{1/p} \\ &\leq C_1 E([\bar{L}, \bar{L}]_{\infty}^{p/2})^{1/p} + E\left(\frac{1}{2}[\bar{L}, \bar{L}]_{\infty}^p\right)^{1/p} \\ &\leq C_1 (1 + E[\bar{L}, \bar{L}]_{\infty}^p)^{1/p} + E\left(\frac{1}{2}[\bar{L}, \bar{L}]_{\infty}^p\right)^{1/p} \\ &\leq C_2 (1 + E([\bar{L}, \bar{L}]_{\infty}^p))^{1/p}, \end{aligned}$$

and thus the result.



Suppose now that the enlargement  $\mathcal{A}$  is induced by some discrete random variable  $G$ , i.e.  $\mathcal{A} = \sigma(G)$ . In that case one can estimate the moments of  $[L, L]_\infty$  against some generalized *absolute* entropy of  $G$ .

**Definition 2.** Let  $(q_g)$  denote the probability weights of  $G$ . We denote by

$$H^p(G) = \sum_g q_g (\log 1/q_g)^p.$$

the generalized absolute entropy of order  $p$ .

**Lemma 11.** One has

$$H^p(\bar{P} \parallel \bar{Q}) \leq H^p(G),$$

and if  $G$  is  $\mathcal{F}_\infty$ -measurable, then

$$H^p(\bar{P} \parallel \bar{Q}) = H^p(G).$$

*Proof.* For the proof we need a monotonicity property of  $f$ -divergences. Due to Corollary 1.29 in [LV87] one has

$$\begin{aligned} H^p(\bar{P} \parallel \bar{Q}) &= H^p(P_{\text{id}_{\mathcal{F}_\infty}, \text{id}_{\mathcal{A}}} \parallel P_{\text{id}_{\mathcal{F}_\infty}} \otimes P_{\text{id}_{\mathcal{A}}}) \\ &\leq H^p(P_{\text{id}_{\mathcal{F}_\infty}, G, \text{id}_{\mathcal{A}}} \parallel P_{\text{id}_{\mathcal{F}_\infty}, G} \otimes P_{\text{id}_{\mathcal{A}}}). \end{aligned}$$

Moreover, if  $G$  is  $\mathcal{F}_\infty$ -measurable, then one even has equality in the previous line. We denote by  $(q_g)$  the probability weights of  $G$ . One easily verifies that

$$\frac{dP_{\text{id}_{\mathcal{F}_\infty}, G, \text{id}_{\mathcal{A}}}}{dP_{\text{id}_{\mathcal{F}_\infty}, G} \otimes P_{\text{id}_{\mathcal{A}}}}(\omega, g, \omega') = 1_{\{g=G(\omega')\}} \frac{1}{q_g}.$$

Set  $f(g, g') = 1_{\{g=g'\}} \frac{1}{q_g}$ . Then

$$\begin{aligned} &H^p(P_{\text{id}_{\mathcal{F}_\infty}, G, \text{id}_{\mathcal{A}}} \parallel P_{\text{id}_{\mathcal{F}_\infty}, G} \otimes P_{\text{id}_{\mathcal{A}}}) \\ &= \int f(g, G(\omega')) (\log_+ f(g, G(\omega')))^p d(P_{\text{id}_{\mathcal{F}_\infty}, G} \otimes P_{\text{id}_{\mathcal{A}}})(\omega, g, \omega') \\ &= \int_{\{(g, \omega') : g=G(\omega')\}} \frac{1}{q_g} \left( \log_+ \frac{1}{q_g} \right)^p d(P_G \otimes P_{\text{id}_{\mathcal{A}}})(g, \omega'), \end{aligned}$$

since  $f(g, G(\omega')) = 0$  if  $g \neq G(\omega')$  and the integrand does not depend on  $\omega$ . Altogether, we arrive at

$$H^p(\bar{P} \parallel \bar{Q}) \leq \sum_g q_g \left( \log \frac{1}{q_g} \right)^p = H^p(G)$$

and equality holds if  $G$  is  $\mathcal{F}_\infty$ -measurable.

*Example 2.* Let  $M_t = W_t$  denote a Wiener process and consider the completed filtration  $(\mathcal{F}_t) = (\mathcal{F}_t^W)$  generated by the Wiener process. We now consider an initial enlargement of the filtration  $(\mathcal{F}_t)$  by some arbitrary  $\sigma$ -field  $\mathcal{A}$ , i.e.  $\mathcal{G}_t = \bigcap_{s>t} (\mathcal{F}_s \wedge \mathcal{A})$ . Supposing that  $\bar{P} \ll \bar{Q}$ , the Doob-Meyer decomposition for  $W$  with respect to  $(\mathcal{G}_t)$  is of the form

$$W_t = \tilde{W}_t + \int_0^t \alpha_s ds,$$

where  $\tilde{W}$  is a  $(\mathcal{G}_t)$ -Wiener process and  $\alpha$  is a  $(\mathcal{G}_t)$ -adapted process. In fact,  $\tilde{W}$  is continuous with quadratic variation process  $[\tilde{W}, \tilde{W}]_t = t$ . Moreover, since  $\mathcal{F}_0$  is trivial and all  $(\mathcal{F}_t)$ -martingales have continuous modifications, the results of this section lead to the estimate

$$E\left(\int_0^t \alpha_s^2 ds\right)^p \leq C_p [H(\bar{P}||\bar{Q}) + H^p(\bar{P}||\bar{Q})].$$

If in addition  $\mathcal{A} = \sigma(G)$  is generated by some discrete random variable  $G$ , then

$$E\left(\int_0^t \alpha_s^2 ds\right)^p \leq C_p [H(G) + H^p(G)].$$

### 3 Continuity of initial enlargements

In section 1 we have seen that every  $(\mathcal{F}_t^P)$ -semimartingale is also a semimartingale relative to a bigger filtration  $(\mathcal{G}_t^P)$  if the measure  $\bar{P}$  is absolutely continuous with respect to  $\bar{Q}$ . In this section we analyze to which extent this embedding of  $(\mathcal{F}_t^P)$ -semimartingales into some space of  $(\mathcal{G}_t^P)$ -semimartingales is continuous. For simplicity we restrict to initial enlargements. It turns out that the embedding is continuous if and only if some generalized entropy of the measures  $\bar{P}$  and  $\bar{Q}$  is finite.

Let  $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$  be a filtered probability space as in the previous section. Throughout this section we assume that  $\mathcal{F}_0$  is trivial and we let

$$\mathcal{G}_t = \bigcap_{s>t} (\mathcal{F}_s \vee \mathcal{A}), \quad t \geq 0,$$

where  $\mathcal{A}$  is some fixed sub- $\sigma$ -algebra of  $\mathcal{F}$ . The measures  $\bar{P}$  and  $\bar{Q}$  are defined as in the previous section and we assume again that  $\bar{P}$  is absolutely continuous with respect to  $\bar{Q}$ . As before we will abbreviate  $\bar{Z}_t = \frac{d\bar{P}}{d\bar{Q}}|_{\mathcal{F}_t}$ ,  $t \geq 0$ . For a treatment of basic questions and ideas of this section in the setting of initial enlargements by random variables see [Imk96].

#### 3.1 Preliminaries

We now recall the definition of some basic norms on the set of semimartingales. For this let  $X$  be a  $(\mathcal{F}_t^P)$ -semimartingale. Given a decomposition  $X = M + A$  we define for all  $1 \leq p < \infty$ ,

$$j_p(M, A) = \left\| [M, M]_\infty^{\frac{1}{2}} + \int_{[0, \infty[} |dA_s| \right\|_{L^p}$$

and

$$\|X\|_{\mathcal{S}^p} = \inf_{X=M+A} j_p(M, A).$$

We denote by  $\mathcal{S}^p$  the set of all  $(\mathcal{F}_t^P)$ -semimartingales  $X$  such that  $\|X\|_{\mathcal{S}^p} < \infty$ . If we want to emphasize the filtration we are referring to we write  $\mathcal{S}^p(\mathcal{F}_t)$ . The space  $\mathcal{S}^p$  is a Banach space with the following properties (see e.g. [DM82]):

- Any  $X \in \mathcal{S}^p$  is a special semimartingale.
- Let  $X \in \mathcal{S}^p$  and  $X = \bar{M} + \bar{A}$  be the unique decomposition such that  $\bar{A}$  is predictable and  $\bar{A}_0 = 0$ . There is a constant  $c > 0$ , depending only on  $p$ , such that  $j_p(\bar{M}, \bar{A}) \leq c\|X\|_{\mathcal{S}^p}$ .
- The space of all martingales in  $\mathcal{S}^p$ , denoted by  $\mathcal{H}^p$ , is a closed subspace.
- The set of all continuous semimartingales in  $\mathcal{S}^p$ , denoted by  $\mathcal{S}_c^p$ , and the set of all continuous martingales in  $\mathcal{S}^p$ , denoted by  $\mathcal{H}_c^p$ , are closed subspaces.
- The set of all predictable processes with integrable variation, vanishing in 0 and with norm  $A \mapsto \|\int |dA_s|\|_{L^p}$  is a closed subspace of  $\mathcal{S}^p$ .

We will see that under suitable conditions every semimartingale in  $\mathcal{S}^2(\mathcal{F}_t)$  belongs to  $\mathcal{S}^1(\mathcal{G}_t)$ .

### 3.2 Continuity and relative entropy

We are now in a position to prove the first main result.

**Theorem 7.** *Suppose  $H_{\mathcal{F}_\infty}(\bar{P}||\bar{Q}) = C < \infty$ . Then the embedding*

$$\mathcal{H}_c^2(\mathcal{F}_t) \rightarrow \mathcal{S}^1(\mathcal{G}_t), X \mapsto X,$$

*is a continuous linear mapping with norm  $\leq 1 + \sqrt{2C}$ .*

*Proof.* Let  $M \in \mathcal{H}^2(\mathcal{F}_t)$ . By Theorem 4,  $(M - [M, L]) + [M, L]$  is a decomposition relative to  $(\mathcal{G}_t)$ . The Kunita-Watanabe inequality implies

$$\left\| \int_0^\infty |d[M, L]_t| \right\|_1 \leq \|[L, L]_\infty^{\frac{1}{2}}\|_2 \|[M, M]_\infty^{\frac{1}{2}}\|_2.$$

Hence by Lemma 7

$$\begin{aligned} \|M\|_{\mathcal{S}^1(\mathcal{G}_t)} &\leq \left\| [M, M]_\infty^{\frac{1}{2}} + \int_0^\infty |d[M, L]_t| \right\|_1 \\ &\leq \left(1 + \|[L, L]_\infty^{\frac{1}{2}}\|_2\right) \|[M, M]_\infty^{\frac{1}{2}}\|_2 \\ &\leq \left(1 + (E[L, L]_\infty)^{\frac{1}{2}}\right) \|M\|_{\mathcal{H}^2(\mathcal{F}_t)} \\ &\leq (1 + \sqrt{2C}) \|M\|_{\mathcal{H}^2(\mathcal{F}_t)}, \end{aligned}$$

and the proof is complete.

As an immediate consequence we get the following

**Corollary 1.** *Suppose  $H_{\mathcal{F}_\infty}(\bar{P}||\bar{Q}) < \infty$ . Then the embedding*

$$\mathcal{S}_c^2(\mathcal{F}_t) \rightarrow \mathcal{S}^1(\mathcal{G}_t), X \mapsto X,$$

*is a continuous linear mapping.*

### 3.3 Continuity and generalized entropy

We aim at generalizing Theorem 7 and Corollary 1. Starting from the Banach space  $\mathcal{S}^r(\mathcal{F}_t)$  with  $r > 1$ , what are sufficient criteria for the embedding into the space of  $(\mathcal{G}_t)$ -semimartingales to be continuous?

Throughout this section we assume assumption (C). In other words, we will assume that  $\mathcal{H}_c^p(\mathcal{F}_t) = \mathcal{H}^p(\mathcal{F}_t)$  for  $p > 1$ .

We begin by stating a result obtained by Yor.

**Lemma 12.** (see Lemme 2 in [Yor85]) *Let  $r \geq 1$  and  $p, q > 0$  such that  $\frac{1}{r} = \frac{1}{2p} + \frac{1}{q}$ . Then the following conditions are equivalent:*

- 1) *There is a constant  $C > 0$  such that every continuous  $(\mathcal{G}_t)$ -local martingale satisfies*

$$\left\| \int_0^\infty |d[M, L]_t| \right\|_r \leq C \| [M, M]_\infty^{\frac{1}{2}} \|_q.$$

- 2)  $E[[L, L]_\infty^p] < \infty$ .

We are now ready to state the main theorem.

**Theorem 8.** *Suppose assumption (C) is satisfied and let  $p \geq 1$  and  $q, r \geq 0$  such that  $\frac{1}{r} = \frac{1}{2p} + \frac{1}{q}$ . The generalized entropy  $H^p(\bar{P}||\bar{Q})$  is finite if and only if the embedding*

$$\mathcal{S}^q(\mathcal{F}_t) \rightarrow \mathcal{S}^r(\mathcal{G}_t), X \mapsto X,$$

*is a continuous linear mapping.*

*Proof.* Suppose  $H^p(\bar{P}||\bar{Q}) < \infty$ . Theorem 6 implies that  $[L, L]_\infty$  is  $L^p$ -integrable. Thus, by Lemma 12, there is a constant  $C > 0$  such that for all continuous  $(\mathcal{G}_t)$ -local martingales we have

$$\left\| \int_0^\infty |d[M, L]_s| \right\|_{L^r} \leq C \| [M, M]_\infty^{\frac{1}{2}} \|_{L^q}.$$

Hence, for a martingale  $M$  in  $\mathcal{S}^q(\mathcal{F}_t)$  with decomposition  $M = (M - [M, L]) + [M, L]$  relative to  $(\mathcal{G}_t)$ , we have

$$\begin{aligned} \|M\|_{\mathcal{S}^r(\mathcal{G}_t)} &= \left\| [M, M]_\infty^{\frac{1}{2}} + \int_0^\infty |d[M, L]_s| \right\|_{L^r} \\ &\leq \| [M, M]_\infty^{\frac{1}{2}} \|_{L^r} + \left\| \int_0^\infty |d[M, L]_s| \right\|_{L^r} \\ &\leq \| [M, M]_\infty^{\frac{1}{2}} \|_{L^r} + C \| [M, M]_\infty^{\frac{1}{2}} \|_{L^q} \\ &\leq (1 + C) \| [M, M]_\infty^{\frac{1}{2}} \|_{L^q} \\ &\leq (1 + C) \|M\|_{\mathcal{S}^q(\mathcal{F}_t)}. \end{aligned}$$

Therefore the map  $\mathcal{S}^q(\mathcal{F}_t) \rightarrow \mathcal{S}^r(\mathcal{G}_t), X \mapsto X$ , is continuous.

Now suppose the embedding to be continuous. Then Lemma 12 implies

$$E[[\bar{L}, \bar{L}]_\infty^p] < \infty.$$

So by Lemma 10 the proof is complete.

*Example 3.* Suppose  $\mathcal{A}$  is generated by a countable partition  $\mathcal{P} = \{A_1, A_2, \dots\}$  of  $\Omega$  into  $\mathcal{F}_\infty$ -measurable sets. Then the corresponding initial enlargement can be viewed as enlargement by the discrete random variable  $G(\omega) := \sum_n n 1_{A_n}(\omega)$ . Hence, for  $p \geq 1$ , we have by Lemma 11

$$H_{\mathcal{F}_\infty}^p(\bar{P} \parallel \bar{Q}) = \sum_{i \geq 1} P(A_i) \left( \log \frac{1}{P(A_i)} \right)^p.$$

Now let  $q, r \geq 0$  such that  $\frac{1}{r} = \frac{1}{2p} + \frac{1}{q}$ . Theorem 8 implies that the embedding  $\mathcal{S}^q(\mathcal{F}_t) \rightarrow \mathcal{S}^r(\mathcal{G}_t), X \mapsto X$ , is a continuous if and only if

$$\sum_{i \geq 1} P(A_i) \left( \log \frac{1}{P(A_i)} \right)^p < \infty.$$

This result was already shown by Marc Yor, using different arguments (see Théorème 2 in [Yor85]).

### 3.4 Continuity and Shannon Information

If the filtration  $(\mathcal{F}_t)$  is generated by a fixed martingale  $M$  with cadlag paths, then the relative entropy of  $\bar{P}$  with respect to  $\bar{Q}$  is equal to the so-called mutual information between  $M$  and the enlarging  $\sigma$ -algebra  $\mathcal{A}$ . We recall this notion.

**Definition 3.** Let  $X$  and  $Y$  be two random variables with values in the measure spaces  $(M, \mathcal{M})$  and  $(K, \mathcal{K})$  respectively. The mutual information between  $X$  and  $Y$  is defined by

$$I(X, Y) = H_{\mathcal{M} \otimes \mathcal{K}}(P_{(X, Y)} \parallel P_X \otimes P_Y).$$

Similarly, one can define the generalized mutual information to be

$$I^p(X, Y) = H_{\mathcal{M} \otimes \mathcal{K}}^p(P_{(X, Y)} \parallel P_X \otimes P_Y), \quad p > 1.$$

For a given  $\sigma$ -algebra  $\mathcal{J} \subset \mathcal{F}$  let  $\text{id}_{\mathcal{J}}$  denote the map  $(\Omega, \mathcal{F}) \rightarrow (\Omega, \mathcal{J}), \omega \mapsto \omega$ . The mutual information between  $X$  and  $\mathcal{J}$  is defined by

$$I(X, \mathcal{J}) = I(X, \text{id}_{\mathcal{J}}).$$

We start with the following observation.

**Lemma 13.** If  $(\mathcal{F}_t)$  equals the filtration generated by  $M$ , then

$$I(M, \mathcal{A}) = H_{\mathcal{F}_\infty}(\bar{P} \parallel \bar{Q}),$$

and for  $p > 1$ ,

$$I^p(M, \mathcal{A}) = H_{\mathcal{F}_\infty}^p(\bar{P} \parallel \bar{Q}).$$

*Proof.* First observe that  $\bar{\mathcal{F}}_\infty = \mathcal{F}_\infty \otimes \mathcal{A}$ , because

$$\bar{\mathcal{F}}_\infty = \bigvee_t \bar{\mathcal{F}}_t \subset \bigvee_t (\mathcal{F}_t \otimes \mathcal{A}) \subset \mathcal{F}_\infty \otimes \mathcal{A} \subset \bar{\mathcal{F}}_\infty.$$

Now let  $\mathbb{D}$  denote the Skorokhod space. We define a map  $\phi$  by

$$\Omega \times \Omega \rightarrow \mathbb{D} \times \Omega, (\omega, \omega') \mapsto (M.(\omega), \omega').$$

Since  $\mathcal{F}_\infty$  is generated by  $M$ , we have

$$\phi^{-1}(\mathcal{B}(\mathbb{D}) \otimes \mathcal{A}) = M^{-1}(\mathcal{B}(\mathbb{D})) \otimes \mathcal{A} = \mathcal{F}_\infty \otimes \mathcal{A},$$

and hence

$$H_{\bar{\mathcal{F}}_\infty}(\bar{P} \parallel \bar{Q}) = H_{\mathcal{B}(\mathbb{D}) \otimes \mathcal{A}}(\bar{P}_\phi \parallel \bar{Q}_\phi).$$

Now observe

$$\bar{P}_\phi = P_{\phi \circ \psi} = P_{(M, \text{id}_{\mathcal{A}})}$$

and

$$\bar{Q}_\phi = P_M \otimes P_{\text{id}_{\mathcal{A}}},$$

which yields the first claim. The second follows by similar arguments.

As a consequence we obtain the following.

**Theorem 9.** *Suppose assumption (C) is satisfied and let  $p \geq 1$  and  $q, r \geq 0$  such that  $\frac{1}{r} = \frac{1}{2p} + \frac{1}{q}$ . If  $(\mathcal{F}_t)$  equals the filtration generated by  $M$ , then the generalized mutual information  $I^p(M, \mathcal{A})$  is finite if and only if the embedding*

$$\mathcal{S}^q(\mathcal{F}_t) \rightarrow \mathcal{S}^r(\mathcal{G}_t), X \mapsto X,$$

*is a continuous linear mapping.*

*Proof.* This follows by combining Theorem 8 with Lemma 13.

*Example 4.* Let  $W$  be the standard Wiener process and  $(\mathcal{F}_t)$  the filtration generated by  $W$  and completed by the negligible sets relative to the Wiener measure. Moreover, let  $V$  be a Gaussian element independent of  $\mathcal{F}_\infty$ , with zero mean and variance  $w > 0$ . Suppose the enlarging  $\sigma$ -algebra  $\mathcal{A}$  is generated by the random variable

$$W_1 + V.$$

One can easily verify that three random variables  $X, Y$  and  $Z$  satisfy

$$I^p(X, (Y, Z)) \leq I^p(X, Z) + I^p(X, Y|Z) \quad (p \geq 1).$$

Consequently, we obtain for the mutual information between  $\text{id}_{\mathcal{A}}$  and  $W$

$$\begin{aligned} I^p(W, \text{id}_{\mathcal{A}}) &= I^p(W_1 + V, (W_1, (W_t)_{0 \leq t < 1})) \\ &\leq I^p(W_1 + V, W_1) + I^p(W_1 + V, (W_t)_{0 \leq t < 1} | W_1) \\ &= I^p(W_1, W_1 + V) \\ &< \infty. \end{aligned}$$

Thus, for all  $p \geq 1$  and  $q, r \geq 0$  such that  $\frac{1}{r} = \frac{1}{2p} + \frac{1}{q}$ , the mapping  $\mathcal{S}^q(\mathcal{F}_t) \rightarrow \mathcal{S}^r(\mathcal{G}_t), X \mapsto X$ , is continuous.

## A Appendix: Product filtrations satisfying the usual conditions

Let  $(\Omega, \mathcal{F}, P)$  be a probability space with right-continuous filtrations  $(\mathcal{F}_t)_{t \geq 0}$  and  $(\mathcal{H}_t)_{t \geq 0}$ . In section 1 the filtration  $(\mathcal{G}_t)$  has been defined as the smallest right-continuous filtration containing

$$\mathcal{G}_t^0 = \mathcal{F}_t \vee \mathcal{H}_t, \quad t \geq 0.$$

We will see that under suitable conditions the filtration  $(\mathcal{G}_t^0)$  itself is already right-continuous.

We start by analyzing the filtration  $(\bar{\mathcal{F}}_t)$  on our product space  $\bar{\Omega}$ .

**Lemma 14.** *The filtration*

$$\mathcal{F}_t \otimes \mathcal{H}_t, \quad t \geq 0,$$

*completed by the  $\bar{Q}$ -negligible sets, is right-continuous. In particular,  $\bar{\mathcal{F}}_t^{\bar{Q}} = (\mathcal{F}_t \otimes \mathcal{H}_t)^{\bar{Q}}$  for  $t \geq 0$ .*

*Proof.* The filtrations  $(\mathcal{F}_t \otimes \{\emptyset, \Omega\})$  and  $(\{\emptyset, \Omega\} \otimes \mathcal{H}_t)$  are right-continuous and independent with respect to  $\bar{Q}$ . According to a result by Wu and Wang, the filtration

$$(\mathcal{F}_t \otimes \{\emptyset, \Omega\}) \vee (\{\emptyset, \Omega\} \otimes \mathcal{H}_t), \quad t \geq 0,$$

completed by the  $\bar{Q}$ -negligible sets, is right-continuous (see Theorem 1 in [HW82]).

With the help of the preceding lemma we can easily derive a sufficient criterion for the filtration  $(\mathcal{G}_t^0)$  to satisfy the usual conditions.

**Theorem 10.** *Suppose that  $(\mathcal{F}_t)_{t \geq 0}$  and  $(\mathcal{H}_t)_{t \geq 0}$  are completed by the  $P$ -negligible sets, and hence satisfy the usual conditions. If  $\bar{P} \ll \bar{Q}$ , then also  $(\mathcal{G}_t^0)$  satisfies the usual conditions.*

*Proof.* Let  $\mathcal{N}^{\bar{Q}}$  denote the set of  $\bar{Q}$ -negligible sets. By Lemma 14, the filtration

$$(\mathcal{F}_t \otimes \mathcal{H}_t) \vee \mathcal{N}^{\bar{Q}}, \quad t \geq 0,$$

is right-continuous. Consequently,

$$\begin{aligned} \psi^{-1} \left( (\mathcal{F}_t \otimes \mathcal{H}_t) \vee \mathcal{N}^{\bar{Q}} \right) &= \mathcal{F}_t \vee \mathcal{H}_t \vee \psi^{-1}(\mathcal{N}^{\bar{Q}}) \\ &= \mathcal{G}_t^0 \vee \psi^{-1}(\mathcal{N}^{\bar{Q}}) \end{aligned}$$

is right-continuous, too. Since  $\bar{P} \ll \bar{Q}$ , we have  $\psi^{-1}(\mathcal{N}^{\bar{Q}}) \subset \mathcal{N}^P$ . This implies that  $(\mathcal{G}_t^0 \vee \psi^{-1}(\mathcal{N}^{\bar{Q}}))$  is equal to  $(\mathcal{G}_t^0)$ , and hence the result.

## References

- [ADI04] S. Ankirchner, S. Dereich, and P. Imkeller. The Shannon information of filtrations and the additional logarithmic utility of insiders. Preprint, 2004.
- [AIS98] J. Amendinger, P. Imkeller, and M. Schweizer. Additional logarithmic utility of an insider. *Stochastic Process. Appl.*, 75(2):263–286, 1998.
- [DM82] C. Dellacherie and P.A. Meyer. *Probabilities and Potential*. North Holland, Amsterdam, 1982.
- [FI93] H. Föllmer and P. Imkeller. Anticipation cancelled by a Girsanov transformation: a paradox on Wiener space. *Ann. Inst. H. Poincaré Probab. Statist.*, 29(4):569–586, 1993.
- [GP98] A. Ghorud and M. Pontier. Insider trading in a continuous time market model. *International Journal of Theoretical and Applied Finance*, 1:331–347, 1998.
- [GV03] D. Gasbarra and E. Valkeila. Initial enlargement: a Bayesian approach. Preprint, 2003.
- [HW82] S. W. He and J. G. Wang. The property of predictable representation of the sum of independent semimartingales. *Z. Wahrsch. Verw. Gebiete*, 61(1):141–152, 1982.
- [Imk96] P. Imkeller. Enlargement of the Wiener filtration by an absolutely continuous random variable via Malliavin’s calculus. *Probab. Theory Related Fields*, 106(1):105–135, 1996.
- [Imk03] P. Imkeller. Malliavin’s calculus in insider models: additional utility and free lunches. *Math. Finance*, 13(1):153–169, 2003. Conference on Applications of Malliavin Calculus in Finance (Rocquencourt, 2001).
- [IPW01] P. Imkeller, M. Pontier, and F. Weisz. Free lunch and arbitrage possibilities in a financial market model with an insider. *Stochastic Process. Appl.*, 92(1):103–130, 2001.
- [Jac79] J. Jacod. *Calcul stochastique et problèmes de martingales*, volume 714 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [Jac85] J. Jacod. Grossissement initial, hypothèse (H’), et théorème de Girsanov. In Th. Jeulin and M. Yor, editors, *Grossissements de filtrations: exemples et applications*, pages 15–35. Springer-Verlag, 1985.
- [JY85] Th. Jeulin and M. Yor, editors. *Grossissements de filtrations: exemples et applications*, volume 1118 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985. Papers from the seminar on stochastic calculus held at the Université de Paris VI, Paris, 1982/1983.
- [Len77] E. Lenglart. Transformation des martingales locales par changement absolument continu de probabilités. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 39(1):65–70, 1977.
- [LV87] F. Liese and I. Vajda. *Convex statistical distances*, volume 95 of *Teubner-Texte zur Mathematik [Teubner Texts in Mathematics]*. BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1987. With German, French and Russian summaries.
- [Pro04] Ph. Protter. *Stochastic Integration and Differential Equations*. Springer-Verlag, New York, second edition, 2004.
- [SY78] C. Stricker and M. Yor. Calcul stochastique dépendant d’un paramètre. *Z. Wahrsch. Verw. Gebiete*, 45(2):109–133, 1978.
- [Yoe85] C. Yoeurp. Grossissement de filtration et théorème de girsanov généralisé. In Th. Jeulin and M. Yor, editors, *Grossissements de filtrations: exemples et applications*. Springer-Verlag, 1985.



- [Yor85] M. Yor. Entropie d'une partition, et grossissement initial d'une filtration. In Th. Jeulin and M. Yor, editors, *Grossissements de filtrations: exemples et applications*, pages 45–58. Springer-Verlag, 1985.