

GENERALISATION OF THE PERRON-FROBENIUS THEORY TO MATRIX PENCILS

VOLKER MEHRMANN , REINHARD NABBEN , AND ELENA VIRNIK *

Dedicated to Hans Schneider on the occasion of his 80th birthday

Abstract. We present a new extension of the well-known Perron-Frobenius theorem to regular matrix pairs (E, A) . The new extension is based on projector chains and is motivated from the solution of positive differential-algebraic systems or descriptor systems. We present several examples where the new condition holds, whereas conditions in previous literature are not satisfied.

Key words. Perron-Frobenius theorem; matrix pencils; differential-algebraic equation, descriptor system, projector chain, tractability index, differentiation index.

AMS subject classifications. 15A18; 15A22; 15A48;

1. Introduction. The well-known Perron-Frobenius Theorem states that for an elementwise nonnegative matrix, the spectral radius, i.e., the largest modulus of an eigenvalue is itself an eigenvalue and has a nonnegative eigenvector. This result has many applications in all areas of science and engineering, in particular in economics and population dynamics see e.g. [2]. If the dynamics of the system, however, is described by an implicit differential or difference equation, usually called differential-algebraic equation (DAE) or descriptor system such as

$$E\dot{x} = Ax + f, \quad x(t_0) = x_0,$$

or as a discrete time system

$$Ex_{k+1} = Ax_k + f_k, \quad x_0 \text{ given,}$$

where E, A are real $n \times n$ matrices, then the dynamics is described by the eigenvalues and eigenvectors associated with the matrix pencil $\lambda E - A$, or just the matrix pair (E, A) .

Due to the many applications, several approaches have been presented in the literature to generalise the classical Perron-Frobenius theory to matrix pencils or further to matrix polynomials. In [11] a direct generalisation of the nonnegativity condition for $A, y \geq 0 \Rightarrow Ay \geq 0$, is given as a sufficient condition, $E^T y \geq 0 \Rightarrow A^T y \geq 0$, for the existence of a positive eigenvalue and a corresponding nonnegative eigenvector. In [1] a sufficient condition, $(E - A)^{-1}A \geq 0$, for the existence of a positive eigenvalue in $(0, 1)$ and a corresponding positive eigenvector if $(E - A)^{-1}A$ is irreducible, is proved. The relationship of the two ideas from [1] and [11] is studied in [15]. In [14], the condition from [1] is imposed by requiring $E - A$ to be a nonsingular M -matrix and $A \geq 0$. Here, the structure of nonnegative eigenvectors is studied from the combinatorial point of view. In [16] the Perron-Frobenius theory was extended to matrix polynomials, where the coefficient matrices are entrywise nonnegative. Other extensions concerning matrix polynomials are given in [6].

*Institut für Mathematik, MA 4-5, TU Berlin, D-10623 Berlin, Germany. E-mail: mehrmann@math.tu-berlin.de, nabben@math.tu-berlin.de, virnik@math.tu-berlin.de. *The research of these authors is supported by the DFG Research Center MATHEON in Berlin.*

The main drawbacks of the generalisation in [1] is that on the one hand it is a restrictive condition, since $E - A$ is not necessarily invertible, and on the other hand it does not have the classical Perron-Frobenius theory as a special case, where $E = I$. Furthermore, only the existence of a nonnegative real eigenvalue is guaranteed instead of the spectral radius being an eigenvalue. The condition in [11] has the classical Perron-Frobenius as a special case but the condition is not easy to verify.

In this paper we propose a new approach to extend the classical Perron-Frobenius theory to matrix pairs (E, A) , where a sufficient condition guarantees that the finite spectral radius of (E, A) is an eigenvalue with a corresponding nonnegative eigenvector. As mentioned before, our approach is motivated by the study of positive systems of differential-algebraic equations, see, e.g., [3],[9]. It is based on the construction of projector chains as they were introduced in the context in [8]. For the special case $E = I$ our new approach reduces to the classical Perron-Frobenius theorem for matrices. We present several examples where the new condition holds, whereas previous conditions in the literature are not satisfied.

2. Preliminaries.

2.1. The generalised eigenvalue problem. Let $E, A \in \mathbb{R}^{n \times m}$. A matrix pair (E, A) , or matrix pencil $\lambda E - A$, is called *regular* if E and A are square ($n = m$) and $\det(\lambda E - A) \neq 0$ for some $\lambda \in \mathbb{R}$. It is called *singular* otherwise. In this paper we only consider square and regular pencils.

A scalar $\lambda \in \mathbb{C}$ is said to be a (*generalised*) *eigenvalue* of the matrix pair (E, A) if $\det(\lambda E - A) = 0$. A vector $x \in \mathbb{C}^n \setminus \{0\}$ such that $(\lambda E - A)x = 0$ is called (*generalised*) *eigenvector* of (E, A) corresponding to λ .

If E is singular and $v \in \mathbb{C}^n \setminus \{0\}$, such that $Ev = 0$ holds, then v is called *eigenvector* of (E, A) corresponding to the eigenvalue ∞ . The equation

$$\lambda Ev = Av, \quad (2.1)$$

is called *generalised eigenvalue problem*. The set of all finite eigenvalues is called *finite spectrum* of (E, A) and is denoted by $\sigma_f(E, A)$. The set of all eigenvalues is called *spectrum* of (E, A) and is defined by

$$\sigma(E, A) := \begin{cases} \sigma_f(E, A), & \text{if } E \text{ is invertible,} \\ \sigma_f(E, A) \cup \{\infty\}, & \text{if } E \text{ is singular.} \end{cases}$$

In the case that E is invertible, we denote by $\rho(E, A)$ the spectral radius of (E, A) defined by

$$\rho(E, A) = \max_{\lambda \in \sigma(E, A)} |\lambda|.$$

If E is singular, then we denote by $\rho_f(E, A) = \max_{\lambda \in \sigma_f(E, A)} |\lambda|$ the *finite spectral radius* of (E, A) . Vectors v_1, \dots, v_k form a *Jordan chain* of the matrix pair (E, A) corresponding to an eigenvalue λ if $(\lambda E - A)v_i = -Ev_{i-1}$ for all $1 \leq i \leq k$ and $v_0 = 0$. A k -dimensional subspace $S_\lambda^{def} \subset \mathbb{C}^n$ is called *right deflating subspace* of (E, A) corresponding to λ , if there exists a k -dimensional subspace $\mathcal{W} \subset \mathbb{C}^n$ such that $ES_\lambda^{def} \subset \mathcal{W}$ and $AS_\lambda^{def} \subset \mathcal{W}$. Let $\lambda_1, \dots, \lambda_p$ be the pairwise distinct finite eigenvalues of (E, A) and let $S_{\lambda_i}^{def}$, $i = 1, \dots, p$ be the corresponding deflating subspaces associated with these eigenvalues. We call the subspace defined by

$$S_f^{def} := S_{\lambda_1}^{def} \oplus \dots \oplus S_{\lambda_p}^{def} \quad (2.2)$$

the *right finite deflating subspace* of (E, A) .

Two matrix pairs (E, A) and (\tilde{E}, \tilde{A}) are called equivalent if there exist regular matrices W, T such that

$$E = W\tilde{E}T, \quad A = W\tilde{A}T. \quad (2.3)$$

In this case we write $(E, A) \sim (\tilde{E}, \tilde{A})$.

THEOREM 2.1 (Weierstraß canonical form). *Let (E, A) be a regular matrix pair. Then, we have*

$$(E, A) \sim \left(\begin{bmatrix} I & 0 \\ 0 & N \end{bmatrix}, \begin{bmatrix} J & 0 \\ 0 & I \end{bmatrix} \right), \quad (2.4)$$

where J is a matrix in Jordan canonical form and N is a nilpotent matrix also in Jordan canonical form.

Proof. See, e.g., [7]. \square

LEMMA 2.2. *Let (E, A) be a regular matrix pair. Let $\hat{\lambda}$ be chosen such that $\hat{\lambda}E - A$ is non-singular. Then, the matrices*

$$\hat{E} = (\hat{\lambda}E - A)^{-1}E \quad \text{and} \quad \hat{A} = (\hat{\lambda}E - A)^{-1}A$$

commute.

Proof. See, e.g., [5], [10]. \square

2.2. Projectors and index of (E, A) . A matrix Q is called *projector* if $Q^2 = Q$. A projector Q is called projector *onto* a subspace $S \subseteq \mathbb{R}^n$ if $\text{im } Q = S$. It is called projector *along* a subspace $S \subseteq \mathbb{R}^n$ if $\ker Q = S$. We will make use of the following well-known property of projectors.

LEMMA 2.3.

1. P_1, P_2 are two projectors onto a subspace S if and only if $P_1 = P_2P_1$ and $P_2 = P_1P_2$.
2. P_1, P_2 are two projectors along a subspace S if and only if $P_1 = P_1P_2$ and $P_2 = P_2P_1$.

Proof.

1. “ \Rightarrow ” For all $x \in \mathbb{R}^n$ we have $P_2x \in \text{im } P_2 = S = \text{im } P_1$ and, hence, $P_2x = P_1P_2x$. Analogously, we show $P_1 = P_2P_1$.
 “ \Leftarrow ” From $P_1 = P_2P_1$, we have $\text{im } P_1 \subset \text{im } P_2$ and from $P_2 = P_1P_2$, we have $\text{im } P_2 \subset \text{im } P_1$ and, hence, P_1, P_2 project onto the same subspace.
2. “ \Rightarrow ” Let $Q_1 = I - P_1$ and $Q_2 = I - P_2$. Then,

$$P_1P_2 = (I - Q_1)(I - Q_2) = I - Q_1 - Q_2 + Q_1Q_2 = I - Q_1 = P_1,$$

since, by 1. we have $Q_1Q_2 = Q_2$. Analogously, we show $P_2 = P_2P_1$.

“ \Leftarrow ” Let $Q_1 = I - P_1$ and $Q_2 = I - P_2$. Then, from $P_1 = P_1P_2$, we have $Q_2 = Q_1Q_2$ and from $P_2 = P_2P_1$, we have $Q_1 = Q_2Q_1$. By 1. we have $\text{im } Q_1 = \text{im } Q_2$ and, hence, $\ker P_1 = \ker P_2$.

\square

Let (E, A) be a regular matrix pair. As introduced in [8] we define a matrix chain by setting

$$E_0 := E, \quad A_0 := A \quad \text{and} \quad (2.5a)$$

$$E_{i+1} := E_i - A_i\tilde{Q}_i, \quad A_{i+1} := A_i\tilde{P}_i, \quad \text{for } i \geq 0, \quad (2.5b)$$

where \tilde{Q}_i are projectors onto $\ker E_i$ and $\tilde{P}_i = I - \tilde{Q}_i$. Since we have assumed (E, A) to be regular, there exists an index ν such that E_ν is nonsingular and all E_i are singular for $i < \nu$, [12]. Note, that ν is independent of a special choice of the projectors Q_i . We say that the matrix pair (E, A) has (*tractability*) *index* ν and denote it by $\text{ind}(E, A) = \nu$. It is well known that for regular pairs (E, A) the tractability index is equal to the differentiation index, see, e.g., [4], and it can be determined as the size of the largest Jordan block associated with the eigenvalue infinity in the Weierstraß canonical form of the pair (E, A) , see [10, 12]. In the following we, therefore, only speak of the *index* of the pair (E, A) .

LEMMA 2.4 ([12]). *Let (E, A) be a matrix pair and define a matrix chain as in (2.5). Furthermore, define sets S_i by*

$$S_i := \{y \in \mathbb{R}^n : A_i y \in \text{im } E_i\}. \quad (2.6)$$

Then, if E_{i+1} is nonsingular, we have that

$$Q_i = -\tilde{Q}_i E_{i+1}^{-1} A_i$$

is a projector onto $\ker E$ along S_i .

Proof. The matrix $-\tilde{Q}_i E_{i+1}^{-1} A_i$ is a projector since

$$\begin{aligned} (-\tilde{Q}_i E_{i+1}^{-1} A_i)^2 &= \tilde{Q}_i E_{i+1}^{-1} (A_i \tilde{Q}_i) \tilde{Q}_i E_{i+1}^{-1} A_i = \tilde{Q}_i E_{i+1}^{-1} (E_i - E_{i+1}) \tilde{Q}_i E_{i+1}^{-1} A_i = \\ &= -\tilde{Q}_i E_{i+1}^{-1} A_i. \end{aligned}$$

For any $z \in \mathbb{R}^n$ the decomposition

$$z = (I + \tilde{Q}_i E_{i+1}^{-1} A_i)z - \tilde{Q}_i E_{i+1}^{-1} A_i z =: z_1 + z_2$$

is unique and we have that $z_2 \in \ker E_i$, since \tilde{Q}_i is a projector onto $\ker E_i$ and for z_1 we obtain

$$A_i z_1 = (I + A_i \tilde{Q}_i E_{i+1}^{-1}) A_i z = (I + (E_i - E_{i+1}) E_{i+1}^{-1}) A_i z = E_i E_{i+1}^{-1} A_i z \in \text{im } E_i,$$

i.e. $z_1 \in S_i$. Hence, $Q_i := -\tilde{Q}_i E_{i+1}^{-1} A_i$ is a projector onto $\ker E$ along S_i . \square

For the construction of special projectors in the higher index cases in Section 3.3, we will need the following properties.

LEMMA 2.5. *Let (E, A) be a regular matrix pair of $\text{ind}(E, A) = \nu$ and define a matrix chain as in (2.5), where the projectors \tilde{Q}_i are chosen such that $\tilde{Q}_j \tilde{Q}_i = 0$ holds for all $0 \leq i < j \leq \nu - 1$. For $0 \leq i \leq \nu - 1$ we define projectors Q_i onto $\ker E_i$ by setting $Q_i = -\tilde{Q}_i E_\nu^{-1} A_i$ and $P_i = I - Q_i$. Then, $Q_j Q_i = 0$ holds for all $0 \leq i < j \leq \nu - 1$.*

Proof. The matrix $-\tilde{Q}_i E_\nu^{-1} A_i$ is a projector for all $0 \leq i \leq \nu - 1$, since

$$\begin{aligned} (-\tilde{Q}_i E_\nu^{-1} A_i)^2 &= \tilde{Q}_i E_\nu^{-1} (E_i - E_{i+1}) \tilde{Q}_i E_\nu^{-1} A_i = \\ &= -\tilde{Q}_i E_\nu^{-1} E_{i+1} \tilde{Q}_i E_\nu^{-1} A_i = -Q_i E_\nu^{-1} A_i, \end{aligned}$$

where we have used that $E_\nu \tilde{Q}_i = (E_{i+1} - A_{i+1} \tilde{Q}_{i+1} - \dots - A_{\nu-1} Q_{\nu-1}) \tilde{Q}_i = E_{i+1} \tilde{Q}_i$.

To show that $Q_j Q_i = 0$ holds for all $0 \leq i < j \leq \nu - 1$, let i, j be arbitrarily chosen fixed indices $0 \leq i < j \leq \nu - 1$. Then, we have that

$$\begin{aligned} Q_j Q_i &= \tilde{Q}_j E_\nu^{-1} A_j \tilde{Q}_i E_\nu^{-1} A_i = \tilde{Q}_j E_\nu^{-1} A_i \tilde{P}_i \dots \tilde{P}_{j-1} \tilde{Q}_i E_\nu^{-1} A_i = \\ &= \tilde{Q}_j E_\nu^{-1} A_i (I - \tilde{Q}_i) \dots (I - \tilde{Q}_{j-1}) \tilde{Q}_i E_\nu^{-1} A_i = \tilde{Q}_j E_\nu^{-1} A_i (\tilde{Q}_i - \tilde{Q}_i) = \\ &= 0. \end{aligned}$$

□

Note that by definition and Lemma 2.4, we have that $Q_{\nu-1}$ is a projector onto $\ker E_{\nu-1}$ along $S_{\nu-1}$. This projector $Q_{\nu-1}$ is called *canonical* in [13].

2.3. Nonnegative matrices. A matrix $T \in \mathbb{R}^{n \times n}$, $T = [t_{ij}]_{i,j=1}^n$ is called *nonnegative* and we write $T \geq 0$ if all entries t_{ij} are nonnegative. The classical Perron-Frobenius Theorem, see, e.g., [2, pp. 26/27], states as follows.

THEOREM 2.6 (Perron-Frobenius Theorem). *Let $T \geq 0$ have the spectral radius $\rho(T)$. Then $\rho(T)$ is an eigenvalue of T and T has a nonnegative eigenvector v corresponding to $\rho(T)$. If, in addition, T is irreducible, then $\rho(T)$ is simple and T has a positive eigenvector v corresponding to $\rho(T)$. Furthermore, if $w > 0$ is an eigenvector of T , then $w = \alpha v$, $\alpha \in \mathbb{R}_+$.*

3. Perron-Frobenius theory for matrix pencils. Several generalisations of the classical Perron-Frobenius Theorem 2.6 have been presented in the literature. In [1] the condition $(E - A)^{-1}A \geq 0$ is shown to be sufficient for the existence of an eigenvalue $\lambda \in (0, 1)$ and a corresponding nonnegative eigenvector. In [11] a direct generalisation of the nonnegativity condition $y \geq 0 \Rightarrow Ay \geq 0$ of A is given as a sufficient condition $E^T y \geq 0 \Rightarrow A^T y \geq 0$ for the existence of a positive eigenvalue and a corresponding nonnegative eigenvector.

In the following two subsections we present a different, projector-based extension of the Perron-Frobenius theory to regular matrix pairs (E, A) that has a number of advantages over the existing conditions in the literature. In Section 3.1, for the case of index one pencils, we prove an easily computable sufficient condition in Theorem 3.1 that guarantees that the finite spectral radius of (E, A) is an eigenvalue with a corresponding nonnegative eigenvector. We present several examples where the new condition holds, whereas the conditions in [1] and [11] are not satisfied. In the general case (where the index may be arbitrary) presented in Section 3.2, we have to impose an additional condition on the projectors, see Lemma 3.5, that is satisfied naturally in the index one case. The general sufficient condition that we then prove in Theorem 3.6 is in the index one case the same as in Theorem 3.1 and also guarantees that the finite spectral radius of (E, A) is an eigenvalue with a corresponding nonnegative eigenvector. In Corollary 3.7, we prove two further conditions that are equivalent to the condition in Theorem 3.6. All conditions have the classical Perron-Frobenius theory as a special case when $E = I$.

3.1. Regular matrix pairs of index at most one. In this subsection we study regular pairs (E, A) of index at most one.

THEOREM 3.1. *Let (E, A) , with $E, A \in \mathbb{R}^{n \times n}$, be a regular matrix pair with $\text{ind}(E, A) \leq 1$. Let Q_0 be a projector onto $\ker E$ along the subspace S_0 defined as in (2.6) for $i = 0$, i.e.,*

$$S_0 := \{y \in \mathbb{R}^n : Ay \in \text{im } E\}, \tag{3.1}$$

let $P_0 = I - Q_0$, and $A_1 = AP_0$. Then $E_1 := E - AQ_0$ is nonsingular and if we have that

$$E_1^{-1}A_1 \geq 0, \tag{3.2}$$

then the finite spectral radius $\rho_f(E, A)$ is an eigenvalue of the matrix pair (E, A) and there exists a nonnegative eigenvector v corresponding to $\rho_f(E, A)$. If $E_1^{-1}A_1$, in

addition, is irreducible, then $\rho_f(E, A)$ is simple and $v > 0$ is unique up to a scalar multiple.

Note, that if $\text{ind}(E, A) = 0$, meaning that E is regular, then we have that $Q_0 = 0$ and $E_1 = E$ and the condition in Theorem 3.1 reduces to the one of the classical Perron-Frobenius theorem for $E^{-1}A$.

Proof. Consider the generalised eigenvalue problem (2.1). Since (E, A) is regular of $\text{ind}(E, A) \leq 1$, we have that E_1 as defined in (2.5) is nonsingular, see [12], and we can premultiply equation (2.1) by E_1^{-1} . By also using that $P_0 + Q_0 = I$, we obtain

$$\begin{aligned} E_1^{-1}(\lambda E - A)(P_0 + Q_0)v &= 0 \\ \Leftrightarrow (\lambda E_1^{-1}E - E_1^{-1}AP_0 - E_1^{-1}AQ_0)v &= 0. \end{aligned}$$

Furthermore, we have $E_1^{-1}E = P_0$ since $E_1P_0 = (E - AQ_0)P_0 = E$ and $-E_1^{-1}AQ_0 = -Q_0$ since $E_1Q_0 = (E - AQ_0)Q_0 = -AQ_0$. Hence, we obtain

$$[(\lambda I - E_1^{-1}A)P_0 + Q_0]v = 0,$$

which after multiplication by P_0 and Q_0 from the left is equivalent to the system of two equations

$$\begin{cases} P_0[(\lambda I - E_1^{-1}A)P_0 + Q_0]v = 0, \\ Q_0[(\lambda I - E_1^{-1}A)P_0 + Q_0]v = 0. \end{cases}$$

We have that Q_0 is a projector onto $\ker E$ along S_0 and by Lemma 2.4 we conclude that $-Q_0E_1^{-1}A$ is also a projector onto $\ker E$ along S_0 . Hence, by part 2. of Lemma 2.3 or by Lemma 2.4, we have that $(-Q_0E_1^{-1}A)P_0 = Q_0P_0 = 0$. Therefore, by writing $P_0 = I - Q_0$ in the first equation, the two equations reduce to

$$\begin{cases} (\lambda I - E_1^{-1}A)P_0v = 0, \\ Q_0v = 0. \end{cases}$$

Since $P_0 = P_0P_0$, this is equivalent to

$$\begin{cases} (\lambda I - E_1^{-1}AP_0)P_0v = 0, \\ Q_0v = 0. \end{cases} \quad (3.3)$$

Setting $x = P_0v$, $y = Q_0v$ and $v = P_0v + Q_0v = x + y$, we obtain a standard eigenvalue problem in the first equation and a linear system in the second equation. From the first equation we know from the Perron-Frobenius Theorem that if $E_1^{-1}AP_0 \geq 0$, then the spectral radius of $E_1^{-1}AP_0$ is an eigenvalue with a corresponding nonnegative eigenvector. If $E_1^{-1}AP_0$ is in addition irreducible, we have that $\rho(E_1^{-1}AP_0)$ is a simple eigenvalue and there is a corresponding positive eigenvector that is unique up to a multiple. Set $\hat{\lambda} := \rho(E_1^{-1}AP_0)$ and let $\hat{x} = P_0v$ be the corresponding nonnegative (positive) eigenvector. Then, we obtain

$$\begin{aligned} E_1^{-1}AP_0\hat{x} &= \hat{\lambda}\hat{x} \\ \Leftrightarrow AP_0\hat{x} &= \hat{\lambda}E_1\hat{x} \\ \Leftrightarrow AP_0\hat{x} &= \hat{\lambda}(E - AQ_0)\hat{x} \\ \Leftrightarrow AP_0P_0v &= \hat{\lambda}EP_0v - AQ_0P_0v \\ \Leftrightarrow A(P_0v + Q_0v) &= \hat{\lambda}Ev \\ \Leftrightarrow Av &= \hat{\lambda}Ev, \end{aligned}$$

which is the generalised eigenvalue problem (2.1). Hence, $\rho(E_1^{-1}AP_0) = \rho_f(E, A)$ and the proof is complete. \square

REMARK 3.2.

1. Considering the case $E = I$ in Theorem 3.1, we have $P_0 = I$, and the condition $E_1^{-1}A_1 \geq 0$ reduces to the condition $A \geq 0$ of the classical Perron-Frobenius theorem.
2. Condition $E_1^{-1}A_1 \geq 0$, written out, reads as

$$(E - A(I - P_0))^{-1}AP_0 \geq 0,$$

which, without the projectors, would be the condition in [1]:

$$(E - A)^{-1}A \geq 0.$$

Yet, whereas $(E - A(I - P_0))$ is nonsingular by construction, the matrix $E - A$ is not necessarily invertible. Hence, the new condition finds a much broader applicability.

EXAMPLE 3.3. Consider the pair (E, A) given by

$$E = \begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

We have $\text{ind}(E, A) = 1$ and the pair has only one finite eigenvalue $\lambda = 0.5$ with eigenvector $v = \begin{bmatrix} v_1 \\ 0 \end{bmatrix}$, where we may normalise the eigenvector so that $v_1 > 0$.

To check the sufficient condition (3.2) of Theorem 3.1, we choose a projector \tilde{Q}_0 onto $\ker E_0$, e.g.,

$$\tilde{Q}_0 = \begin{bmatrix} 1 & 0 \\ -1 & 0 \end{bmatrix},$$

and get

$$\tilde{E}_1 = E_0 - A_0\tilde{Q}_0 = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}.$$

For the inverse we obtain

$$\tilde{E}_1^{-1} = \begin{bmatrix} 0 & 1 \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix},$$

and a projector Q_0 onto $\ker E$ along S_0 is given by

$$Q_0 = -\tilde{Q}_0\tilde{E}_1^{-1}A_0 = \begin{bmatrix} 0 & -1 \\ 0 & 1 \end{bmatrix}.$$

We then have

$$E_1 = E_0 - A_0Q_0 = \begin{bmatrix} 2 & 3 \\ 0 & -1 \end{bmatrix}, \quad E_1^{-1} = \begin{bmatrix} \frac{1}{2} & \frac{3}{2} \\ 0 & -1 \end{bmatrix},$$

and we set $P_0 = I - Q_0$. Condition (3.2) then reads

$$E_1^{-1}A_1 = E_1^{-1}AP_0 = \begin{bmatrix} \frac{1}{2} & \frac{3}{2} \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{bmatrix} \geq 0,$$

and we can apply Theorem 3.1.

For this example, the theories in [1] and [11] cannot be applied, since $(E-A)^{-1}A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix} \not\geq 0$ and there exists a vector, e.g., $y = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ such that $E^T y \geq 0$ but $A^T y \not\geq 0$.

EXAMPLE 3.4. Consider a pair (E, A) with $\text{ind}(E, A) = 1$ and $E = \begin{bmatrix} E_{11} & 0 \\ 0 & 0 \end{bmatrix}$, where E_{11} is nonsingular and $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$ is partitioned accordingly. For a pencil in this form, $\text{ind}(E, A) = 1$ is equivalent to A_{22} being nonsingular, see, e.g., [10]. We choose any projector \tilde{Q}_0 onto $\ker E$, e.g.

$$\tilde{Q}_0 = \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix},$$

and compute \tilde{E}_1 and \tilde{E}_1^{-1} . We obtain

$$\tilde{E}_1 = E - A\tilde{Q}_0 = \begin{bmatrix} E_{11} & -A_{12} \\ 0 & -A_{22} \end{bmatrix}, \quad \tilde{E}_1^{-1} = \begin{bmatrix} E_{11}^{-1} & -E_{11}^{-1}A_{12}A_{22}^{-1} \\ 0 & -A_{22}^{-1} \end{bmatrix}.$$

Then, we determine a projector Q_0 onto $\ker E$ along $S_0 = \{y \in \mathbb{R}^n : Ay \in \text{im } E\}$ as

$$Q_0 = -\tilde{Q}_0 E_1^{-1} A = \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} E_{11}^{-1} & -E_{11}^{-1}A_{12}A_{22}^{-1} \\ 0 & -A_{22}^{-1} \end{bmatrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ A_{22}^{-1}A_{21} & I \end{bmatrix}.$$

Furthermore, we get $P_0 = \begin{bmatrix} I & 0 \\ -A_{22}^{-1}A_{21} & 0 \end{bmatrix}$ and then compute E_1 and E_1^{-1} . We obtain

$$E_1 = E - AQ_0 = \begin{bmatrix} E_{11} - A_{12}A_{22}^{-1}A_{21} & -A_{12} \\ -A_{21} & -A_{22} \end{bmatrix},$$

$$E_1^{-1} = \begin{bmatrix} E_{11}^{-1} & -E_{11}^{-1}A_{12}A_{22}^{-1} \\ -A_{22}^{-1}A_{21}E_{11}^{-1} & A_{22}^{-1}A_{21}E_{11}^{-1}A_{12}A_{22}^{-1} - A_{22}^{-1} \end{bmatrix}.$$

Condition (3.2) then reads as

$$E_1^{-1}AP_0 = \begin{bmatrix} E_{11}^{-1} & -E_{11}^{-1}A_{12}A_{22}^{-1} \\ -A_{22}^{-1}A_{21}E_{11}^{-1} & A_{22}^{-1}A_{21}E_{11}^{-1}A_{12}A_{22}^{-1} - A_{22}^{-1} \end{bmatrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} I & 0 \\ -A_{22}^{-1}A_{21} & 0 \end{bmatrix} =$$

$$= \begin{bmatrix} E_{11}^{-1}A_S & 0 \\ -A_{22}^{-1}A_{21}E_{11}^{-1}A_S & 0 \end{bmatrix} \geq 0,$$

where $A_S = A_{11} - A_{12}A_{22}^{-1}A_{21}$.

Consider again the eigenvalue problem

$$(\lambda E - A)v = 0.$$

In our case we obtain

$$\begin{bmatrix} \lambda E_{11} - A_{11} & -A_{12} \\ -A_{21} & -A_{22} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = 0.$$

Since E_{11} is nonsingular, we can rewrite this system as

$$\begin{cases} (\lambda I - E_{11}^{-1}A_{11})v_1 - E_{11}^{-1}A_{12}v_2 &= 0, \\ -A_{21}v_1 - A_{22}v_2 &= 0, \end{cases}$$

which is equivalent to

$$\begin{cases} (\lambda I - E_{11}^{-1}A_S)v_1 &= 0, \\ v_2 &= -A_{22}^{-1}A_{21}v_1, \end{cases} \quad (3.4)$$

where $A_S = A_{11} - A_{12}A_{22}^{-1}A_{21}$. Condition (3.2) gives $E_{11}^{-1}A_S \geq 0$ and, by the Perron-Frobenius Theorem, we obtain from the first equation of (3.4) that $\rho(E_{11}^{-1}A_S) =: \hat{\lambda}$ is an eigenvalue and with a corresponding eigenvector $v_1 \geq 0$. Using this, from the second equation of (3.4) we obtain

$$v_2 = -A_{22}^{-1}A_{21}v_1 = -\lambda^{-1}A_{22}^{-1}A_{21}\lambda v_1 = -\lambda^{-1}A_{22}^{-1}A_{21}E_{11}^{-1}A_S v_1 \geq 0,$$

since $-A_{22}^{-1}A_{21}E_{11}^{-1}A_S \geq 0$ by (3.2) and we have $\hat{\lambda} \geq 0$ and $v_1 \geq 0$ from the first equation of (3.4).

3.2. Regular matrix pairs of general index. In this section we consider a regular matrix pair (E, A) of $\text{ind}(E, A) = \nu$. For $\nu > 1$ we need to define the matrix chain in (2.5) with special projectors. The following Lemma 3.5 guarantees the existence of projectors with the required property. The canonical projectors as defined in [13] fulfil the condition of Lemma 3.5. An alternative way to construct these projectors along with some examples is presented in Section 3.3.

LEMMA 3.5. *Let (E, A) , with $E, A \in \mathbb{R}^{n \times n}$, be a regular matrix pair of $\text{ind}(E, A) = \nu$. Then, a matrix chain as in (2.5) can be constructed with special projectors Q_i, P_i such that $Q_i v = 0$ holds for all $v \in S_f^{def}$ and for all $0 \leq i < \nu$.*

Proof. From [12] we know that for a regular matrix pair (E, A) , we have that

$$\ker E_i \cap \ker A_i = \{0\} \quad (3.5)$$

holds for all $0 \leq i < \nu$. Furthermore, from (3.5) or from [12] we obtain that

$$\ker E_i \cap \ker E_{i+1} = \{0\} \quad (3.6)$$

for all $0 \leq i < \nu - 1$. We now show by induction that we can construct projectors Q_i such that $Q_i v = 0$ holds for all $v \in S_f^{def}$ and for all $0 \leq i < \nu$. For the existence of such a Q_0 , we have to show that $\ker E_0 \cap S_f^{def} = \{0\}$. Suppose that $x \in \ker E_0 \cap S_f^{def}$. Then from $E_0 x = 0$ we obtain that $x = 0$, since otherwise, by definition, x would be an eigenvector of (E, A) corresponding to the eigenvalue ∞ . Thus, we can choose the projector Q_0 onto $\ker E_0$ along some subspace M_0 that completely contains S_f^{def} . This ensures $Q_0 v = 0$ for all $v \in S_f^{def}$. Now, suppose that $Q_i v = 0$ holds for all $v \in S_f^{def}$ and for all $0 \leq i \leq k$ for some $k < \nu - 1$. Note that for the complementary projectors $P_i = I - Q_i$, this implies that $P_i v = v$ for all $v \in S_f^{def}$. To construct a projector Q_{k+1} onto $\ker E_{k+1}$ such that $Q_{k+1} v = 0$ holds for all $v \in S_f^{def}$, we have to show that $\ker E_{k+1} \cap S_f^{def} = \{0\}$. For this, suppose that $0 \neq x \in \ker E_{k+1} \cap S_f^{def}$. Then, by using the assumption, we obtain

$$0 = E_{k+1}x = (E_0 - A_0Q_0 - \dots - A_kQ_k)x = E_0x,$$

from which we again conclude that $x = 0$, since otherwise, by definition, x would be an eigenvector of (E, A) corresponding to the eigenvalue ∞ . Thus, we can choose the projector Q_{k+1} onto $\ker E_{k+1}$ along some subspace M_{k+1} that completely contains S_f^{def} . This ensures $Q_{k+1}v = 0$ for all $v \in S_f^{def}$ and completes the proof. \square

Note that for $\nu = 1$, condition $Q_0v = 0$ holds naturally for all $v \in S_f^{def}$ and in particular for all eigenvectors, see (3.3).

The following theorem states the main result of the present work. We prove a new, broadly applicable Perron-Frobenius-type condition for matrix pairs (E, A) in the general index case.

THEOREM 3.6. *Let (E, A) , with $E, A \in \mathbb{R}^{n \times n}$, be a regular matrix pair of $\text{ind}(E, A) = \nu$. Let a matrix chain as in (2.5) be constructed with projectors Q_i, P_i as in Lemma 3.5. If*

$$E_\nu^{-1}A_\nu \geq 0, \quad (3.7)$$

holds, then the finite spectral radius $\rho_f(E, A)$ is an eigenvalue and there exists a corresponding nonnegative eigenvector $v \geq 0$. If $E_\nu^{-1}A_\nu$ is, in addition, irreducible, then we have that $\rho_f(E, A)$ is simple and $v > 0$ is unique up to a scalar multiple.

Proof. Consider the generalised eigenvalue problem (2.1)

$$(\lambda E - A)v = 0.$$

If v is an eigenvector corresponding to a finite eigenvalue, i.e., $v \in S_f^{def}$, then we have $Q_i v = 0$ for all $0 \leq i < \nu$ and we can equivalently express (2.1) as

$$\begin{aligned} (\lambda(E - A_0Q_0 - A_1Q_1 - \dots - A_{\nu-1}Q_{\nu-1}) - A)v &= 0 \\ \Leftrightarrow (\lambda E_\nu - A)v &= 0 \\ \Leftrightarrow (\lambda I - E_\nu^{-1}A)v &= 0. \end{aligned} \quad (3.8)$$

By construction, we have that $\hat{v} = P_0 \cdots P_{\nu-1} \hat{v}$ and we obtain that (3.8) is equivalent to

$$\begin{aligned} (\hat{\lambda}I - E_\nu^{-1}A)P_0 \cdots P_{\nu-1} \hat{v} &= 0 \\ \Leftrightarrow (\hat{\lambda}I - E_\nu^{-1}AP_0 \cdots P_{\nu-1})P_0 \cdots P_{\nu-1} \hat{v} &= 0 \\ \Leftrightarrow (\hat{\lambda}I - E_\nu^{-1}A_\nu) \hat{v} &= 0. \end{aligned}$$

By the Perron-Frobenius Theorem, we obtain that $\hat{\lambda} := \rho(E_\nu^{-1}A_\nu)$ is an eigenvalue and there exists a corresponding eigenvector $\hat{v} \geq 0$. Hence,

$$\begin{aligned} E_\nu^{-1}A_\nu \hat{v} &= \hat{\lambda} \hat{v} \\ \Leftrightarrow AP_0 \cdots P_{\nu-1} \hat{v} &= \hat{\lambda} E_\nu \hat{v} \\ \Leftrightarrow A \hat{v} &= \hat{\lambda} (E - A_0Q_0 - A_1Q_1 - \dots - A_{\nu-1}Q_{\nu-1}) \hat{v} \\ \Leftrightarrow A \hat{v} &= \hat{\lambda} E \hat{v}, \end{aligned}$$

which is the generalised eigenvalue problem (2.1). Hence, we have that $\rho(E_\nu^{-1}A_\nu) = \rho_f(E, A)$ is an eigenvalue and $\hat{v} \geq 0$ is a corresponding eigenvector. If, in addition, $E_\nu^{-1}A_\nu$ is irreducible, then we have that $\rho_f(E, A)$ is simple and $v > 0$ is unique up to a scalar multiple. \square

COROLLARY 3.7. *Let P_r be a projector onto the right finite deflating subspace S_f^{def} , let \hat{E} be defined as in Lemma 2.2 and let \hat{E}^D denote the Drazin inverse of \hat{E} . Under the assumptions of Theorem 3.6 the conditions*

$$E_\nu^{-1}AP_r \geq 0, \quad (3.9)$$

$$E_\nu^{-1}A\hat{E}^D\hat{E} \geq 0 \quad (3.10)$$

are equivalent to condition (3.7).

Proof. See [13] or Appendix A. \square

REMARK 3.8. Condition

$$E_\nu^{-1}A \geq 0, \quad (3.11)$$

can also be proved to be sufficient in Theorem 3.6, see Equation (3.8), yet there is no evidence for it to ever hold.

3.3. Construction of projectors. In Section 3.2, Lemma 3.5, we have proved the existence of special projectors for constructing the matrix chain in (2.5) in order to establish a sufficient condition in Theorem 3.6 for $\rho_f(E, A)$ to be an eigenvalue with a corresponding nonnegative eigenvector. In this section we show how these projectors can be constructed. A different way to construct (canonical) projectors that fulfil the condition of Lemma 3.5 is given in [13].

Consider a regular matrix pair (E, A) of $\text{ind}(E, A) = \nu$. We make the following observations:

1. For fixed projectors $Q_0, \dots, Q_{\nu-2}$, the projector $Q_{\nu-1}$ is uniquely defined by being a projector onto $\ker E_{\nu-1}$ along $S_{\nu-1}$, since if there were two projectors $Q_{\nu-1}, \tilde{Q}_{\nu-1}$ with these two properties, by Lemma 2.3, we would obtain $Q_{\nu-1} = Q_{\nu-1}\tilde{Q}_{\nu-1} = \tilde{Q}_{\nu-1}$.
2. Consider the sets S_i as defined in (2.6). We have that $S_f^{def} \subseteq S_0$, since for any $v \in S_f^{def}$ there exists a $w \in S_f^{def}$ such that $Av = Ew$, and hence, $Av \in \text{im } E$, i.e., $v \in S_0$. Furthermore, we have that $S_0 \subseteq S_1 \subseteq \dots \subseteq S_{\nu-1}$, see [12], and therefore, $S_f^{def} \subseteq S_{\nu-1}$. From this we conclude that $Q_{\nu-1}v = 0$ holds for all $v \in S_f^{def}$.

In the following, recursive constructions of matrix and projector chains, we denote by $E_j^{(i)}, A_j^{(i)}, Q_j^{(i)}, P_j^{(i)}$ the i -th iterate of E_j, A_j, Q_j, P_j in the recursive construction.

With the chain constructed in (2.5), we set $E_j^{(1)} = E_j, A_j^{(1)} = A_j, Q_j^{(1)} = Q_j$, and $P_j^{(1)} = P_j$.

If the index is $\text{ind}(E, A) = \nu = 1$, then we take any projector $Q_0^{(1)}$ onto $\ker E$ and having computed $E_1^{(1)}$ we set $Q_0^{(2)} = -Q_0^{(1)}(E_1^{(1)})^{-1}A$. By using $Q_0^{(2)}$, we compute $E_1^{(2)}$ and its inverse and use it to check the sufficient condition in Theorems 3.1 (or Theorem 3.6) that now reads $(E_1^{(2)})^{-1}AP_0^{(2)} \geq 0$.

For $\nu = 2$ we already have to construct the projectors recursively. We choose any projectors $Q_0^{(1)}, Q_1^{(1)}$ onto $\ker E_0^{(1)}, \ker E_1^{(1)}$, respectively. We then determine $E_2^{(1)}$ and set $Q_1^{(2)} = -Q_1^{(1)}(E_2^{(1)})^{-1}A_1^{(1)}$. Then we have $Q_1^{(2)}v = 0$ for all $v \in S_f^{def}$. By using $Q_1^{(2)}$ compute $E_2^{(2)}$. We proceed by setting $Q_0^{(2)} = -Q_0^{(1)}(E_2^{(2)})^{-1}A_0^{(1)}$, which is

a projector by Lemma 2.5. For any $v \in S_f^{def}$ we have $w \in S_f^{def}$ such that

$$Q_0^{(2)}v = -Q_0^{(1)}(E_2^{(2)})^{-1}Av = -Q_0^{(1)}(E_2^{(2)})^{-1}Ew = -Q_0^{(1)}(I - Q_0^{(1)} - Q_1^{(2)})w = 0,$$

since $Q_1^{(2)}w = 0$. Here we have used the properties $(E_2^{(2)})^{-1}A_i^{(1)}Q_i^{(1)} = -Q_i^{(1)}$ for $i = 0, 1$ and

$$\begin{aligned} E_2^{(2)} &= E_0^{(1)} - A_0^{(1)}Q_0^{(1)} - A_1^{(1)}Q_1^{(2)} \\ \Leftrightarrow I &= (E_2^{(2)})^{-1}E_0^{(1)} - (E_2^{(2)})^{-1}A_0^{(1)}Q_0^{(1)} - (E_2^{(2)})^{-1}A_1^{(1)}Q_1^{(2)} \\ \Leftrightarrow I &= (E_2^{(2)})^{-1}E_0^{(1)} + Q_0^{(1)} + Q_1^{(2)}. \end{aligned}$$

By using $Q_0^{(2)}$ we compute $E_1^{(2)}$ and $A_1^{(2)}$. Now, we proceed as in the case $\nu = 1$ to define $Q_1^{(3)}$ as a projector onto $\ker E_1^{(2)}$. To ensure that it projects along S_1 we again compute $E_2^{(3)}$, set $Q_1^{(4)} = -Q_1^{(3)}(E_2^{(3)})^{-1}A_1^{(2)}$ and obtain that $Q_1^{(4)}v = 0$ for all $v \in S_f^{def}$. Finally, we compute $E_2^{(4)}$. The sufficient condition of Theorem 3.6 is then checked with $E_2^{(4)}$ instead of E_2 and reads $E_2^{(4)}AP_0^{(2)}P_1^{(4)} \geq 0$. For an illustration of the recursive construction of the projectors in the index 2 case with the properties required in Theorem 3.6, see Figure 3.1.




FIGURE 3.1. Illustration of the recursive construction of projectors in the index 2 case. Top down, we have the chain matrices in increasing order. From left to right, we have the successive calculation of these.

Before proceeding to the calculation of projectors in the general index case, we present two index 2 examples, where condition (3.7) of Theorem 3.6 holds, whereas the conditions in [1] and [11] do not hold.

EXAMPLE 3.9. Consider the matrix pair (E, A) with

$$E = \left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{array} \right] \quad \text{and} \quad A = \left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{array} \right].$$

We have that (E, A) is regular with $\text{ind}(E, A) = 2$ and there is one finite eigenvalue $\rho_f(E, A) = 2$ and a corresponding eigenvector $[0 \ 0 \ v_3]^T$, which can be chosen so that $v_3 > 0$.

We compute the matrix chain by setting, e.g.,

$$Q_0^{(1)} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad E_1^{(1)} = E - AQ_0^{(1)} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A_1^{(1)} = A_0P_0^{(1)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

We choose, e.g.,

$$Q_1^{(1)} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \text{and} \quad P_1^{(1)} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

and compute

$$E_2^{(1)} = E_1^{(1)} - A_1^{(1)}Q_1^{(1)} = \begin{bmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad (E_2^{(1)})^{-1} = \begin{bmatrix} 0 & -1 & 0 \\ 1 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Then, we compute the projector onto $\ker E_1^{(1)}$ along S_1 by setting

$$Q_1^{(2)} = -Q_1^{(1)}(E_2^{(1)})^{-1}A_1^{(1)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

and determine

$$E_2^{(2)} = E_1^{(1)} - A_1^{(1)}Q_1^{(2)} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad (E_2^{(2)})^{-1} = \begin{bmatrix} -1 & -1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We set

$$Q_0^{(2)} = -Q_0^{(1)}(E_2^{(2)})^{-1}A_0 = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad P_0^{(2)} = \begin{bmatrix} 0 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

and compute

$$E_1^{(2)} = E_0 - A_0Q_0^{(2)} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad A_1^{(2)} = A_0P_0^{(2)} = \begin{bmatrix} 0 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

Choosing $Q_1^{(3)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, we determine

$$E_2^{(3)} = E_1^{(2)} - A_1^{(2)}Q_1^{(3)} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = E_2^{(2)} \quad \text{and} \quad (E_2^{(3)})^{-1} = (E_2^{(2)})^{-1},$$

and verify that $Q_1^{(4)} = -Q_1^{(3)}(E_2^{(3)})^{-1}A_1^{(2)} = Q_1^{(3)}$. We finally set $P^{(4)} = I - Q_1^{(4)}$. The sufficient condition (3.7) of Theorem 3.6 then holds, since

$$(E_2^{(4)})^{-1}AP_0^{(2)}P_1^{(4)} = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 2 \end{bmatrix} \geq 0.$$

The condition in [1], however, is not satisfied, since

$$(E - A)^{-1}A = \begin{bmatrix} -1 & -1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} -1 & -1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \not\geq 0.$$

Also the condition in [11] does not hold, since, e.g., for $y = [-1 \ 1 \ 1]^T$ we have $Ey \geq 0$ but $Ay \not\geq 0$. Note, that we have $P_r = P_0^{(2)}P_1^{(4)}$, yet, condition (3.11) does not hold, since $(E_2^{(4)})^{-1}A \not\geq 0$.

EXAMPLE 3.10. Consider the regular matrix pair (E, A) of $\text{ind}(E, A) = 2$, where

$$E = \begin{bmatrix} E_{11} & E_{12} & 0 & 0 \\ E_{21} & E_{22} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad A = \begin{bmatrix} 0 & 0 & 0 & A_{14} \\ 0 & A_{22} & 0 & 0 \\ 0 & 0 & A_{33} & 0 \\ A_{41} & 0 & 0 & 0 \end{bmatrix}.$$

Note, that every regular matrix pair of index 2 can be equivalently transformed into such a form, where $A_{14}, A_{41}, A_{33}, E_{22}$ are square regular matrices, see [10]. We choose

$$Q_0^{(1)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & 0 & I \end{bmatrix} \quad \text{and compute}$$

$$P_0^{(1)} = \begin{bmatrix} I & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad E_1^{(1)} = \begin{bmatrix} E_{11} & E_{12} & 0 & -A_{14} \\ E_{21} & E_{22} & 0 & 0 \\ 0 & 0 & -A_{33} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Choosing

$$Q_1^{(1)} = \begin{bmatrix} I & 0 & 0 & 0 \\ -E_{22}^{-1}E_{21} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ A_{14}^{-1}\tilde{E}_{11} & 0 & 0 & 0 \end{bmatrix},$$

where $\tilde{E}_{11} = E_{11} - E_{12}E_{22}^{-1}E_{21}$, we obtain

$$P_1^{(1)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ E_{22}^{-1}E_{21} & I & 0 & 0 \\ 0 & 0 & I & 0 \\ -A_{14}^{-1}\tilde{E}_{11} & 0 & 0 & I \end{bmatrix}, \quad A_1^{(1)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & A_{22} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ A_{41} & 0 & 0 & 0 \end{bmatrix}, \quad \text{and}$$

$$E_2^{(1)} = \begin{bmatrix} E_{11} & E_{12} & 0 & -A_{14} \\ E_{21} + A_{22}E_{22}^{-1}E_{21} & E_{22} & 0 & 0 \\ 0 & 0 & -A_{33} & 0 \\ -A_{41} & 0 & 0 & 0 \end{bmatrix},$$

$$(E_2^{(1)})^{-1} = \begin{bmatrix} 0 & 0 & 0 & -A_{41}^{-1} \\ 0 & E_{22}^{-1} & 0 & E_{22}^{-1}(E_{21} + A_{22}E_{22}^{-1}E_{21})A_{41}^{-1} \\ 0 & 0 & -A_{33}^{-1} & 0 \\ -A_{14}^{-1} & A_{14}^{-1}E_{12}E_{22}^{-1} & 0 & -A_{14}^{-1}(\tilde{E}_{11} - E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}E_{21})A_{41}^{-1} \end{bmatrix}.$$

We verify that $Q_1^{(2)} = -Q_1^{(1)}(E_2^{(1)})^{-1}A_1^{(1)} = Q_1^{(1)}$ and, hence, $P_1^{(2)} = P_1^{(1)}$, $A_1^{(2)} = A_1^{(1)}$, $E_2^{(2)} = E_2^{(1)}$ and $(E_2^{(2)})^{-1} = (E_2^{(1)})^{-1}$. Setting

$$Q_0^{(2)} = -Q_0^{(1)}(E_2^{(2)})^{-1}A_0 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & I \\ A_{14}(\tilde{E}_{11} - E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}E_{21}) & -A_{14}E_{12}E_{22}^{-1}A_{22} & 0 & I \end{bmatrix},$$

we compute

$$P_0^{(2)} = \begin{bmatrix} I & 0 & 0 & 0 \\ 0 & I & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -A_{14}(\tilde{E}_{11} - E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}E_{21}) & A_{14}E_{12}E_{22}^{-1}A_{22} & 0 & 0 \end{bmatrix},$$

$$E_1^{(2)} = E - AQ_0^{(2)} = \begin{bmatrix} E_{12}(I + E_{22}^{-1}A_{22})E_{22}^{-1}E_{21} & E_{12}(I + E_{22}^{-1}A_{22}) & 0 & -A_{14} \\ E_{21} & E_{22} & 0 & 0 \\ 0 & 0 & -A_{33} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$A_1^{(2)} = AP_0^{(2)} = \begin{bmatrix} -E_{11} + E_{12}(I + E_{22}^{-1}A_{22})E_{22}^{-1}E_{21} & E_{12}E_{22}^{-1}A_{22} & 0 & 0 \\ 0 & A_{22} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ A_{41} & 0 & 0 & 0 \end{bmatrix}.$$

Choosing

$$Q_1^{(3)} = \begin{bmatrix} I & 0 & 0 & 0 \\ -E_{22}^{-1}E_{21} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

we determine

$$P_1^{(3)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ E_{22}^{-1}E_{21} & I & 0 & 0 \\ 0 & 0 & I & 0 \\ 0 & 0 & 0 & I \end{bmatrix},$$

$$E_2^{(3)} = \begin{bmatrix} E_{11} + E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}E_{21} & E_{12} + E_{12}E_{22}^{-1}A_{22} & 0 & -A_{14} \\ E_{21} + A_{22}E_{22}^{-1}E_{21} & E_{22} & 0 & 0 \\ 0 & 0 & -A_{33} & 0 \\ -A_{41} & 0 & 0 & 0 \end{bmatrix},$$

$$(E_2^{(3)})^{-1} = \begin{bmatrix} 0 & 0 & 0 & -A_{41}^{-1} \\ 0 & E_{22}^{-1} & 0 & E_{22}^{-1}(E_{21} + A_{22}E_{22}^{-1}E_{21})A_{41}^{-1} \\ 0 & 0 & -A_{33}^{-1} & 0 \\ -A_{14}^{-1} & A_{14}^{-1}E_{12}(I + E_{22}^{-1}A_{22})E_{22}^{-1} & 0 & -A_{14}^{-1}(\tilde{E}_{11} - E_{12}E_{22}^{-1}A_{22}(I + E_{22}^{-1}A_{22})E_{22}^{-1}E_{21})A_{41}^{-1} \end{bmatrix}.$$

We verify that $Q_1^{(4)} = -Q_1^{(3)}(E_2^{(3)})^{-1}A_1^{(2)} = Q_1^{(3)}$ and, hence, $P_1^{(4)} = P_1^{(3)}$, $E_2^{(4)} = E_2^{(3)}$ and $(E_2^{(4)})^{-1} = (E_2^{(3)})^{-1}$. The sufficient condition (3.7) of Theorem 3.6 then reads as

$$(E_2^{(4)})^{-1}A_1^{(2)}P_1^{(4)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ E_{22}^{-1}A_{22}E_{22}^{-1}E_{21} & E_{22}^{-1}A_{22} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ A_{14}^{-1}E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}E_{21} & A_{14}^{-1}E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}A_{22} & 0 & 0 \end{bmatrix} \geq 0.$$

Consider again the eigenvalue problem

$$(\lambda E - A)v = 0.$$

For the given E and A , we obtain

$$\begin{bmatrix} \lambda E_{11} & \lambda E_{12} & 0 & -A_{14} \\ \lambda E_{21} & \lambda E_{22} - A_{22} & 0 & 0 \\ 0 & 0 & -A_{33} & 0 \\ -A_{14} & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix} = 0.$$

Since A_{41} and A_{33} are nonsingular, we obtain $v_1 = v_3 = 0$ and the following system of equations

$$\begin{cases} \lambda E_{12}v_2 - A_{14}v_4 = 0, \\ (\lambda E_{22} - A_{22})v_2 = 0, \end{cases}$$

which is equivalent to

$$\begin{cases} (\lambda I - E_{22}^{-1}A_{22})v_2 = 0, \\ v_4 = \lambda A_{14}^{-1}E_{12}v_2. \end{cases}$$

Condition (3.7) gives $E_{22}^{-1}A_{22} \geq 0$ and, hence, we obtain from the first equation that $\rho(E_{22}^{-1}A_{22}) =: \lambda$ is an eigenvalue and there exists a corresponding eigenvector $v_2 \geq 0$. By using this, we obtain from the second equation that

$$v_4 = \lambda A_{14}^{-1}E_{12}v_2 = A_{14}^{-1}E_{12}E_{22}^{-1}A_{22}v_2 = -\lambda^{-1}A_{14}^{-1}E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}A_{22}v_2 \geq 0,$$

since $A_{14}^{-1}E_{12}E_{22}^{-1}A_{22}E_{22}^{-1}A_{22} \geq 0$ by (3.7) and $\lambda \geq 0$, $v_2 \geq 0$ from the first equation.

The condition in [1], however, is not necessarily applicable, since $E - A$ may not be invertible if $E_{22} - A_{22}$ is not. Also the condition in [11] will not hold in most cases, since we may choose y_1, y_2 in $y = [y_1 \ y_2 \ y_3 \ y_4]^T$ such that $Ey \geq 0$ and choose y_3, y_4 such that $Ay \not\geq 0$.

We proceed with the construction of the special projectors as required in Theorem 3.6 in the case that $\nu > 2$. To begin with, we choose the projectors $Q_0^{(1)}, \dots, Q_{\nu-1}^{(1)}$ as in [12] such that $Q_j^{(1)}Q_i^{(1)} = 0$ for all $j > i$. Then, we define the projector $Q_{\nu-1}^{(2)}$ onto $\ker E_{\nu-1}^{(1)}$ along $S_{\nu-1}$ by setting $Q_{\nu-1}^{(2)} = -Q_{\nu-1}^{(1)}(E_{\nu-1}^{(1)})^{-1}A_{\nu-1}^{(1)}$ as in Lemma 2.4 and setting $P_{\nu-1}^{(2)} = I - Q_{\nu-1}^{(2)}$ and $E_{\nu-1}^{(2)} = E_{\nu-1}^{(1)} - A_{\nu-1}^{(1)}Q_{\nu-1}^{(2)}$.

Next, we set $Q_{\nu-2}^{(2)} = -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-2}^{(1)}$. This is a projector by Lemma 2.5 and we obtain $Q_{\nu-2}^{(2)}v = 0$ for all $v \in S_f^{def}$. This follows, since for all $v \in S_f^{def}$ there exists $w \in S_f^{def}$ with $Av = Ew$ and thus

$$\begin{aligned} Q_{\nu-2}^{(2)}v &= -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-2}^{(1)}v = -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-3}^{(1)}P_{\nu-3}^{(1)}v = \\ &= -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-3}^{(1)}(I - Q_{\nu-3}^{(1)})v = -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-3}^{(1)}v - Q_{\nu-2}^{(1)}Q_{\nu-3}^{(1)}v = \\ &= -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_{\nu-4}^{(1)}P_{\nu-4}^{(1)}v = \dots = -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}A_0v = \\ &= -Q_{\nu-2}^{(1)}(E_{\nu-2}^{(1)})^{-1}E_0w = -Q_{\nu-2}^{(1)}(I - Q_0^{(1)} - Q_1^{(1)} - \dots - Q_{\nu-2}^{(1)} - Q_{\nu-1}^{(1)})w = \\ &= Q_{\nu-2}^{(1)}Q_{\nu-1}^{(1)}w = 0. \end{aligned}$$

By using the new projector $Q_{\nu-2}^{(2)}$ we redefine the last part of the chain by setting $P_{\nu-2}^{(2)} = I - Q_{\nu-2}^{(2)}$, $E_{\nu-1}^{(2)} = E_{\nu-2}^{(1)} - A_{\nu-2}^{(1)}Q_{\nu-2}^{(2)}$, $A_{\nu-1}^{(2)} = A_{\nu-2}^{(1)}P_{\nu-2}^{(2)}$ and $E_{\nu-1}^{(3)} = E_{\nu-1}^{(2)}$ -

$A_{\nu-1}^{(2)}Q_{\nu-1}^{(2)}$. Since we have a new $E_{\nu-1}^{(2)}$ we may need to redefine $Q_{\nu-1}^{(3)}$ and $Q_{\nu-1}^{(4)}$ as in the index 2 case.

Then we proceed by setting $Q_{\nu-3}^{(2)} = -Q_{\nu-3}^{(1)}(E_{\nu-3}^{(3)})^{-1}A_{\nu-3}^{(1)}$ and show in the same manner that $Q_{\nu-3}^{(2)}v = 0$ for all $v \in S_f^{def}$. With the same procedure we redefine all projectors down to $Q_0^{(2)}$. We redefine the matrix chain using the final projector $Q_0^{(2)}$ and proceed by applying the same procedure as if we had a chain of length $\nu - 1$. Eventually, we obtain a matrix chain with projectors that satisfy the condition $Q_i^{(2^{i+1})}v = 0$ for all $v \in S_f^{def}$ and for all $i = 0, \dots, \nu - 1$. In total, we have to make $\sum_{i=1}^{\nu} 2^i = 2^{\nu} - 1$ updates of the projectors Q_i . The sufficient condition of Theorem 3.6 is then checked with $E_{\nu-1}^{(2^{\nu})}$ instead of $E_{\nu-1}$ and reads

$$\begin{aligned} E_{\nu-1}^{(2^{\nu})}AP_0^{(2)}P_1^{(4)} \dots P_{\nu-1}^{(2^{\nu})} &\geq 0 \\ \Leftrightarrow E_{\nu-1}^{(2^{\nu})}A_{\nu-1}^{(2^{\nu-1})}P_{\nu-1}^{2^{\nu}} &\geq 0. \end{aligned}$$

4. Conclusions. We have presented a new generalisation of the Perron-Frobenius Theorem to regular matrix pairs (E, A) of arbitrary index. We have constructed projector chains that allow to derive a new sufficient condition. The new generalisation reduces to the classical Perron-Frobenius Theorem if $E = I$ and is different from previous such generalisations. We have demonstrated the broad applicability of the new generalisation by several examples.

Appendix A. Proof of Corollary 3.7.

Proof. In [13], the canonical projectors $\tilde{P}_0 \dots \tilde{P}_{\nu-1}$ are constructed such that the condition $\tilde{Q}_i = -\tilde{Q}_i \tilde{P}_{i+1} \dots \tilde{P}_{\nu-1} \tilde{E}_{\nu-1}^{-1} \tilde{A}_i$ holds for all i . For the canonical projectors, conditions (3.7), (3.9), (3.10) are equivalent, since $\tilde{P}_0 \dots \tilde{P}_{\nu-1} = P_r = \hat{E}^D \hat{E}$, see [13]. Now, we show that for projectors $P_0, \dots, P_{\nu-1}$ as constructed in Section 3.3, we have $P_0 \dots P_{\nu-1} = \tilde{P}_0 \dots \tilde{P}_{\nu-1}$, where $\tilde{P}_0, \dots, \tilde{P}_{\nu-1}$ are canonical projectors as introduced in [13]. For a matrix pair of $\text{ind}(E, A) = \nu$, we have the projector decomposition $I = P_0 \dots P_{\nu-1} + P_0 \dots P_{\nu-2}Q_{\nu-1} + \dots + P_0Q_1 + Q_0$. Furthermore, we have the property, see [13],

$$\text{im}(P_0 \dots P_{\nu-2}Q_{\nu-1} + \dots + P_0Q_1 + Q_0) = \text{im } Q_0 + \dots + \text{im } Q_{\nu-1}.$$

For the canonical projectors \tilde{Q}_i as constructed in [13] and for the projectors Q_i as constructed in Section 3.3, we have $\text{im } Q_i = \text{im } \tilde{Q}_i$ for all i . Hence,

$$\text{im}(P_0 \dots P_{\nu-2}Q_{\nu-1} + \dots + P_0Q_1 + Q_0) = \text{im}(\tilde{P}_0 \dots \tilde{P}_{\nu-2}\tilde{Q}_{\nu-1} + \dots + \tilde{P}_0\tilde{Q}_1 + \tilde{Q}_0).$$

Therefore, we conclude that $\ker P_0 \dots P_{\nu-1} = \ker \tilde{P}_0 \dots \tilde{P}_{\nu-1}$ since $P_0 \dots P_{\nu-2}Q_{\nu-1} + \dots + P_0Q_1 + Q_0 = I - P_0 \dots P_{\nu-1}$ is a projector and

$$\dim(\text{im } P_0 \dots P_{\nu-1}) = \dim(\text{im } \tilde{P}_0 \dots \tilde{P}_{\nu-1}) = n - \dim(\ker P_0 \dots P_{\nu-1}).$$

On the other hand, we know that $\text{im } P_r \subset \text{im } P_0 \dots P_{\nu-1}$, since for any $x \in \mathbb{R}^n$ we get $P_r x \in S_f^{def}$ and $P_0 \dots P_{\nu-1}P_r x = P_r x$. This together with the dimension argument gives us $P_r = \tilde{P}_0 \dots \tilde{P}_{\nu-1} = P_0 \dots P_{\nu-1}$.

Condition (3.10) is sufficient, since $\hat{E}^D \hat{E} = P_r$. This can be verified by transforming $\hat{E}^D \hat{E}$ into Weierstraß canonical form. If

$$\begin{aligned} \hat{E} &= (sE - A)^{-1}E = \left(sW \begin{bmatrix} I & 0 \\ 0 & N \end{bmatrix} T - W \begin{bmatrix} J & 0 \\ 0 & I \end{bmatrix} T \right)^{-1} W \begin{bmatrix} I & 0 \\ 0 & N \end{bmatrix} T = \\ &= T^{-1} \begin{bmatrix} (sI - J)^{-1} & 0 \\ 0 & (sN - I)^{-1}N \end{bmatrix} T, \end{aligned}$$

then

$$\hat{E}^D = T^{-1} \begin{bmatrix} sI - J & 0 \\ 0 & 0 \end{bmatrix} T,$$

and

$$\begin{aligned} \hat{E}^D \hat{E} &= T^{-1} \begin{bmatrix} sI - J & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} (sI - J)^{-1} & 0 \\ 0 & (sN - I)^{-1}N \end{bmatrix} T = \\ &= T^{-1} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} T = P_r. \end{aligned}$$

□

REFERENCES

- [1] R. B. BAPAT, D. D. OLESKY, AND P. V. DEN DRIESSCHE, *Perron-Frobenius theory for a generalized eigenproblem*, Linear and Multilinear Algebra, 40 (1995), pp. 141–152.
- [2] A. BERMAN AND R. J. PLEMMONS, *Nonnegative Matrices in the Mathematical Sciences*, vol. 9 of Classics in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, 1994. Second edition.
- [3] R. BRU, C. COLL, S. ROMERO-VIVO, AND E. SÁNCHEZ, *Some problems about structural properties of positive descriptor systems*, in Positive systems (Rome, 2003), vol. 294 of Lecture Notes in Control and Inform. Sci., Springer, Berlin, 2003, pp. 233–240.
- [4] S. L. CAMPBELL, *A general form for solvable linear time varying singular systems of differential equations*, SIAM J. Math. Anal., 18 (1987), pp. 1101–1115.
- [5] S. L. CAMPBELL AND C. D. MEYER, JR., *Generalized inverses of linear transformations*, Dover Publications Inc., New York, 1991. Corrected reprint of the 1979 original.
- [6] K.-H. FÖRSTER AND B. NAGY, *Spectral properties of operator polynomials with nonnegative coefficients*, in Operator theory and indefinite inner product spaces, vol. 163 of Oper. Theory Adv. Appl., Birkhäuser, Basel, 2006, pp. 147–162.
- [7] F. R. GANTMACHER, *The theory of matrices. Vols. 1, 2*, Translated by K. A. Hirsch, Chelsea Publishing Co., New York, 1959.
- [8] E. GRIEPENTROG AND R. MÄRZ, *Differential-algebraic equations and their numerical treatment*, vol. 88 of Teubner-Texte zur Mathematik [Teubner Texts in Mathematics], BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1986.
- [9] T. KACZOREK, *Positive 1D and 2D Systems*, Springer-Verlag, London, 2002.
- [10] P. KUNKEL AND V. MEHRMANN, *Differential-algebraic equations. Analysis and numerical solution*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2006.
- [11] O. L. MANGASARIAN, *Perron-Frobenius properties of $Ax - \lambda Bx$* , J. Math. Anal. Appl., 36 (1971), pp. 86–102.
- [12] R. MÄRZ, *Numerical methods for differential algebraic equations*, in Acta numerica, 1992, Cambridge Univ. Press, Cambridge, 1992, pp. 141–198.
- [13] ———, *Canonical projectors for linear differential algebraic equations*, Comput. Math. Appl., 31 (1996), pp. 121–135.
- [14] J. J. McDONALD, D. D. OLESKY, H. SCHNEIDER, M. J. TSATSOMEROS, AND P. VAN DEN DRIESSCHE, *Z-pencils*, Electron. J. Linear Algebra, 4 (1998), pp. 32–38 (electronic).
- [15] V. MEHRMANN, D. OLESKY, T. PHAN, AND P. VAN DEN DRIESSCHE, *Relations between Perron-Frobenius results for matrix pencils*, Linear Algebra Appl., 287 (1998), pp. 257–269.
- [16] P. J. PSARRAKOS AND M. J. TSATSOMEROS, *A primer of Perron-Frobenius theory for matrix polynomials*, Linear Algebra Appl., 393 (2004), pp. 333–351.