

Utility maximization in incomplete markets*

Ying Hu

IRMAR

Campus de Beaulieu

Université de Rennes 1

F-35042 Rennes Cedex

France

Peter Imkeller

Institut für Mathematik

Humboldt-Universität zu Berlin

Unter den Linden 6

D-10099 Berlin

Germany

Matthias Müller

Institut für Mathematik

Humboldt-Universität zu Berlin

Unter den Linden 6

D-10099 Berlin

Germany

November 25, 2003

Abstract

We consider the problem of utility maximization for small traders on incomplete financial markets. As opposed to most of the papers dealing with this subject, the investors' trading strategies we allow underly constraints described by closed, but not necessarily convex, sets. The final wealths obtained by trading under these constraints are identified as stochastic processes which usually are supermartingales, and even martingales for particular strategies. These strategies are seen to be optimal, and the corresponding value functions determined simply by the initial values of the supermartingales. We separately treat the cases of exponential, power and logarithmic utility.

2000 AMS subject classifications: primary 60 H 10, 91 B 28; secondary 60 G 44, 91 B 70, 91 B 16, 60 H 20, 93 E 20.

Key words and phrases: financial market; incomplete market; maximal utility; exponential utility; power utility; logarithmic utility; supermartingale; stochastic differential equation; backwards stochastic differential equation.

*This work was partially supported by the DFG research center 'Mathematics for key technologies' (FZT 86) in Berlin.

Introduction

In this paper we consider a small trader on an incomplete financial market who can trade in a finite time interval $[0, T]$ by investing in risky stocks and a riskless bond. He aims at maximizing the utility he draws from his final wealth measured by some utility function. The trading strategies he may choose to attain his wealth underly some restriction formalized by a constraint. For example, he may be forced not to have a negative number of shares or that his investment in risky stocks is not allowed to exceed a certain threshold. We will be interested not only in describing the trader's optimal utility, but also the strategies which he may follow to reach this goal. As opposed to most of the papers dealing so far with the maximization of expected utility under constraints we essentially relax the hypotheses to be fulfilled by them. They are formulated as usual by the requirement that the strategies take their values in some set, which is supposed to simply be closed instead of convex. We consider three types of utility functions. In the second section we carry out the calculation of the value function and an optimal strategy for exponential utility. In this case, the investor is allowed to have an additional liability, and maximizes the utility of its sum with terminal wealth. In section 3 we consider power utility, and in the final section the simplest one: logarithmic utility.

The method that we apply in order to obtain value function and optimal strategy is simple. We propose to construct a stochastic process R^ρ depending on the investor's trading strategy ρ , and such that its terminal value equals the utility of the trader's terminal wealth. As mentioned above, to model the constraint, trading strategies are supposed to take their values in a closed set. In our market, the absence of completeness is not explicitly described by a set of martingale measures equivalent to the historical probability. Instead, we assume more directly that for every trading strategy ρ , R^ρ is a supermartingale. Moreover, we assume that there exists at least one particular trading strategy ρ^* such that R^{ρ^*} is a martingale. Hereby, the initial value is supposed not to depend on the strategy. Evidently, the strategy ρ^* related to the martingale has to be the optimal one. Then the value function of the optimization problem is just given by the initial value of R^{ρ^*} .

Since we work on a Wiener filtration, the powerful tool of backward stochastic differential equations (BSDE) is available. It allows the construction of the stochastic control process ρ^* , and thus the description of the value function in terms of the solution of a BSDE.

In a related paper, El Karoui and Rouge [ER] compute the value function and the optimal strategy for exponential utility by means of BSDE, assuming more restrictively that the strategies be confined to a convex cone. Sekine [Sek] relies on a duality result obtained by Cvitanic and Karatzas [CK], also describing constraints through convex cones. He studies the maximization problem for the exponential and power utility functions, and uses an attainability condition which solves the primal and dual problems, finally writing this condition as a BSDE. In contrast to these papers, we do not use duality, and directly characterize the solution of the primal problem. This allows us to pass from convex to closed constraints.

Utility maximization is one of the most frequent problems in financial mathematics

and has been considered by numerous authors. Here are some of the milestones viewed from our perspective of maximization under constraints using the tools of BSDEs. For a complete market, utility maximization has been considered in [KLS]. Cvitanic and Karatzas [CK] prove existence and uniqueness of the solution for the utility maximization problem in a Brownian filtration constraining strategies to convex sets. There are numerous papers considering general semimartingales as stock price processes. Delbaen et al. [DGR] give a duality result between the optimal strategy for the maximization of the exponential utility and the martingale measure minimizing the relative entropy with respect to the real world measure P . This duality can be used to characterize the utility indifference price for an option. Also relying upon duality theory, Kramkov and Schachermayer [KS] and Cvitanic et al. [CSW] give a fairly complete solution of the utility optimization problem on incomplete markets for a class of general utility functions not containing the exponential one. See also the review paper by Schachermayer [Sch] for a more complete account and further references.

The powerful tool of BSDE has been introduced to stochastic control theory by Bismut [B]. Its mathematical treatment in terms of stochastic analysis was initiated by Pardoux and Peng [PP], and its particular significance for the field of utility maximization in financial stochastics clarified in El Karoui, Peng and Quenez [EPQ].

1 Preliminaries and the market model

A probability space (Ω, \mathcal{F}, P) carrying an m -dimensional Brownian motion $(W_t)_{t \in [0, T]}$ is given. The filtration \mathbb{F} is the completion of the filtration generated by W .

Let us briefly explain some special notation that will be used in the paper. For $q \geq 1$, L^q denotes the set of \mathcal{F}_T -measurable random variables F such that $E[F^q] < \infty$, for $k \in \mathbb{N}$, $\mathcal{H}^k(\mathbb{R}^d)$ the set of all \mathbb{R}^d -valued stochastic processes ϑ which are predictable with respect to \mathbb{F} and satisfy $E[\int_0^T \|\vartheta_t\|^k dt] < \infty$. $\mathcal{H}^\infty(\mathbb{R}^d)$ is the set of all \mathbb{F} -predictable \mathbb{R}^d -valued processes that are $\lambda \otimes P$ -a.e. bounded on $[0, T] \times \Omega$. Note here that we write λ for the Lebesgue measure on $[0, T]$ or \mathbb{R} .

Let M denote a continuous semimartingale. The stochastic exponential $\mathcal{E}(M)$ is given by

$$\mathcal{E}(M)_t = \exp\left(M_t - \frac{1}{2}\langle M \rangle_t\right), \quad t \in [0, T],$$

where the quadratic variation is denoted by $\langle M \rangle$. Let C denote a closed subset of \mathbb{R}^m and $a \in \mathbb{R}^m$. If $|\cdot|$ stands for the Euclidean norm in \mathbb{R}^m , the distance between a and C is defined as

$$\text{dist}_C(a) = \min_{b \in C} |a - b|.$$

The set $\Pi_C(a)$ consists of those elements of C at which the minimum is obtained:

$$\Pi_C(a) = \{b \in C : |a - b| = \text{dist}_C(a)\}.$$

This set is not empty and evidently may contain more than one point.

The financial market consists of one bond with interest rate zero and $d \leq m$ stocks. In case $d < m$ we face an incomplete market. The price process of stock i evolves according to the equation

$$\frac{dS_t^i}{S_t^i} = b_t^i dt + \sigma_t^i dW_t, \quad i = 1, \dots, d, \quad (1)$$

where b^i (resp. σ^i) is an \mathbb{R} -valued (resp. $\mathbb{R}^{1 \times m}$ -valued) predictable uniformly bounded stochastic process. The lines of the $d \times m$ -matrix σ are given by the vector σ_t^i , $i = 1, \dots, d$. The volatility matrix $\sigma = (\sigma^i)_{i=1, \dots, d}$ has full rank and we assume that $\sigma \sigma^{tr}$ is uniformly elliptic, i.e. $KI_d \geq \sigma \sigma^{tr} \geq \varepsilon I_d$, P -a.s. for constants $K > \varepsilon > 0$. The predictable \mathbb{R}^m -valued process

$$\theta_t = \sigma_t^{tr} (\sigma_t \sigma_t^{tr})^{-1} b_t, \quad t \in [0, T],$$

is then also uniformly bounded.

There exists a nonempty set \mathbb{P} of probability measures equivalent to P such that S is a Q -martingale for all $Q \in \mathbb{P}$. A probability measure $Q^0 \in \mathbb{P}$ is determined by an exponential density with respect to a market price of risk process θ in the form

$$\frac{dQ^0}{dP} = \mathcal{E}\left(-\int_0^\cdot \theta dW\right)_T.$$

Let us now fix a market price of risk θ and the corresponding probability measure Q^0 . A d -dimensional \mathbb{F} -predictable process $\pi = (\pi_t)_{0 \leq t \leq T}$ is called trading strategy if $\int \pi \frac{dS}{S}$ is well defined, e.g. $\int_0^T \|\pi_t \sigma_t\|^2 dt < \infty$ P -a.s. and Q^0 -a.s. For $1 \leq i \leq d$, the process π_t^i describes the amount of money invested in stock i at time t . The number of shares is $\frac{\pi_t^i}{S_t^i}$. The wealth process X^π of a trading strategy π with initial capital x satisfies the equation

$$X_t^\pi = x + \sum_{i=1}^d \int_0^t \frac{\pi_{i,u}}{S_{i,u}} dS_{i,u} = x + \int_0^t \pi_u \sigma_u (dW_u + \theta_u du), \quad t \in [0, T].$$

In this notation π has to be taken as a vector in $\mathbb{R}^{1 \times d}$. Trading strategies are self-financing. The investor uses his initial capital and during the trading interval $[0, T]$ there is no extra money flow out of or into his portfolio. Gains or losses are only obtained by trading with the stock.

The conditions on the trading strategies of the following definition guarantee that there is no arbitrage. In addition, we allow constraints on the trading strategies. Formally, they are supposed to take their values in a closed set, i.e. $\pi_t(\omega) \in \tilde{C}$, with $\tilde{C} \subseteq \mathbb{R}^{1 \times d}$. We emphasize that \tilde{C} is not assumed to be convex. For technical reasons we impose some further integrability assumptions.

Definition 1 (Admissible Strategies with constraints) *Let \tilde{C} be a closed set in $\mathbb{R}^{1 \times d}$. The set of admissible trading strategies $\tilde{\mathcal{A}}$ consists of all d -dimensional predictable processes $\pi = (\pi_t)_{0 \leq t \leq T}$ which satisfy $\pi_t \in \tilde{C}$ $\lambda \otimes P$ -a.s., $\int_0^\cdot \pi_s \sigma_s dW_s$ is a P -BMO-martingale, and $E(U(X_T^\pi)) > -\infty$.*

The boundedness of θ and Theorem 3.6 in [Kaz] imply that the wealth process X^π is a BMO–martingale under the equivalent probability measure Q^0 . Therefore the set $\tilde{\mathcal{A}}$ is free of arbitrage, i.e. in this set there is no trading strategy π with initial capital $X_0^\pi = 0$, terminal wealth $X_T^\pi \geq 0$ P -a.s. and $P[X_T^\pi > 0] > 0$.

For $t \in [0, T], \omega \in \Omega$ define the set $C_t(\omega) \subseteq \mathbb{R}^m$ by

$$C_t(\omega) = \tilde{C}\sigma_t(\omega). \quad (2)$$

The entries of the matrix–valued process σ are uniformly bounded. Therefore we get

$$\min\{|a| : a \in C_t(\omega)\} \leq k_1 \quad \text{for} \quad \lambda \otimes P - \text{a.e. } (t, \omega) \quad (3)$$

with a constant $k_1 \geq 0$.

Remark 2 *Writing*

$$p_t = \pi_t \sigma_t, \quad t \in [0, T],$$

the set of admissible trading strategies $\tilde{\mathcal{A}}$ is equivalent to a set \mathcal{A} of $\mathbb{R}^{1 \times m}$ -valued predictable stochastic processes p with $p \in \mathcal{A}$ iff $p_t(\omega) \in C_t(\omega)$ P -a.s. and $\int_0^\cdot p_s dW_s$ is a P -BMO–martingale. Such a process $p \in \mathcal{A}$ will also be named strategy, and $X^{(p)}$ denotes its wealth process.

Suppose our investor has a liability F at time T . This random variable F is assumed to be \mathcal{F}_{T-} -measurable and bounded, but not necessarily positive. He tries to find a trading strategy that is optimal in a certain sense in presence of this liability F .

2 Exponential Utility

In this section, we specify this sense by stipulating that the investor wants to maximize his expected utility with respect to the exponential utility from his total wealth $X_T^p - F$. Let us recall that for $\alpha > 0$ the exponential utility function is defined as

$$U(x) = -\exp(-\alpha x), \quad x \in \mathbb{R}.$$

So the investor wants to solve the maximization problem

$$V(x) := \sup_{\pi \in \tilde{\mathcal{A}}} E \left[-\exp \left(-\alpha \left(x + \int_0^T \pi_t \frac{dS_t}{S_t} - F \right) \right) \right],$$

where x is the initial wealth. V is called value function. Losses, i.e. realizations with $X^\pi - F < 0$, are punished very strongly. Large gains or realizations with $X^\pi - F > 0$ are weakly valued. The maximization problem is evidently equivalent to

$$V(x) = \sup_{p \in \mathcal{A}} E \left[-\exp \left(-\alpha \left(x + \int_0^T p_t (dW_t + \theta_t dt) - F \right) \right) \right]. \quad (4)$$

In order to find the value function and an optimal strategy we construct a family of stochastic processes $R^{(p)}$ with the following properties:

- $R_T^{(p)} = -\exp(-\alpha(X_T^p - F))$ for all $p \in \mathcal{A}$,
- $R_0^{(p)} = R_0$ is constant for all $p \in \mathcal{A}$,
- $R^{(p)}$ is a supermartingale for all $p \in \mathcal{A}$ and there exists a $p^* \in \mathcal{A}$ such that $R^{(p^*)}$ is a martingale.

The process $R^{(p)}$ and its initial value R_0 depend of course on the initial capital x . Given processes possessing these properties we can compare the expected utilities of the strategies $p \in \mathcal{A}$ and $p^* \in \mathcal{A}$ by

$$E[-\exp(-\alpha(X_T^p - F))] \leq R_0(x) = E[-\exp(-\alpha(X_T^{p^*} - F))] = V(x), \quad (5)$$

whence p^* is the desired optimal strategy. To construct this family, we set

$$R_t^{(p)} := -\exp(-\alpha(X_t^{(p)} - Y_t)), \quad t \in [0, T], \quad p \in \mathcal{A},$$

where (Y, Z) is a solution of the BSDE

$$Y_t = F - \int_t^T Z_s dW_s - \int_t^T f(s, Z_s) ds, \quad t \in [0, T].$$

In these terms we are bound to choose a function f for which $R^{(p)}$ is a supermartingale for all $p \in \mathcal{A}$ and there exists a $p^* \in \mathcal{A}$ such that $R^{(p^*)}$ is a martingale. This function f also depends on the constraint set (C_t) where (p_t) takes its values (see (2)). We get

$$V(x) = R_0^{(p,x)} = -\exp(-\alpha(x - Y_0)), \quad \text{for all } p \in \mathcal{A}.$$

In order to calculate f , we write R as the product of a (local) martingale $M^{(p)}$ and a (not strictly) decreasing process $\tilde{A}^{(p)}$ that is constant for some $p^* \in \mathcal{A}$. For $t \in [0, T]$ define

$$M_t^{(p)} = \exp(-\alpha(x - Y_0)) \exp\left(-\int_0^t \alpha(p_s - Z_s) dW_s - \frac{1}{2} \int_0^t \alpha^2 (p_s - Z_s)^2 ds\right)$$

Comparing $R^{(p)}$ and $M^{(p)} \tilde{A}^{(p)}$ yields

$$\tilde{A}_t^{(p)} = -\exp\left(\int_0^t v(s, p_s, Z_s) ds\right), \quad t \in [0, T],$$

with

$$v(t, p, z) = -\alpha p \theta_t + \alpha f(t, z) + \frac{1}{2} \alpha^2 |p - z|^2.$$

In order to obtain a decreasing process $\tilde{A}^{(p)}$ evidently f has to satisfy

$$v(t, p_t, Z_t) \geq 0 \quad \text{for all } p \in \mathcal{A}$$

and

$$v(t, p_t^*, Z_t) = 0$$

for some particular $p^* \in \mathcal{A}$. For $t \in [0, T]$ we have

$$\begin{aligned} \frac{1}{\alpha}v(t, p_t, Z_t) &= \frac{\alpha}{2}|p_t|^2 - \alpha p_t \left(Z_t + \frac{1}{\alpha}\theta_t \right) + \frac{\alpha}{2}|Z_t|^2 + f(t, Z_t) \\ &= \frac{\alpha}{2}|p_t - (Z_t + \frac{1}{\alpha}\theta_t)|^2 - \frac{\alpha}{2}|Z_t + \frac{1}{\alpha}\theta_t|^2 + \frac{\alpha}{2}Z_t^2 + f(t, Z_t) \\ &= \frac{\alpha}{2}|p_t - (Z_t + \frac{1}{\alpha}\theta_t)|^2 - Z_t\theta_t - \frac{1}{2\alpha}|\theta_t|^2 + f(t, Z_t). \end{aligned}$$

Now set

$$f(t, z) = -\frac{\alpha}{2}\text{dist}^2\left(z + \frac{1}{\alpha}\theta_t, C_t(\omega)\right) + z\theta_t + \frac{1}{2\alpha}|\theta_t|^2.$$

The function f is well defined because it only depends on the distance between a point and a closed set.

For this choice we get $v(t, p, z) \geq 0$ and for

$$p_t^* \in \Pi_{C_t(\omega)}\left(Z_t + \frac{1}{\alpha}\theta_t\right), \quad t \in [0, T],$$

we obtain $v(\cdot, p^*, Z) = 0$.

Theorem 3 *The value function of the optimization problem (4) is given by*

$$V(x) = -\exp(-\alpha(x - Y_0)),$$

where Y_0 is defined by a solution (Y, Z) of the BSDE

$$Y_t = F - \int_t^T Z_s dW_s - \int_t^T f(s, Z_s) ds, \quad t \in [0, T], \quad (6)$$

with

$$f(\cdot, z) = -\frac{\alpha}{2}\text{dist}^2\left(z + \frac{1}{\alpha}\theta, C\right) + z\theta + \frac{1}{2\alpha}|\theta|^2.$$

There exists an optimal trading strategy $p^* \in \mathcal{A}$ with

$$p_t^* \in \Pi_{C_t(\omega)}\left(Z_t + \frac{1}{\alpha}\theta_t\right), \quad t \in [0, T], \quad P - a.s. \quad (7)$$

Proof In order to get the existence of solutions of the BSDE (6) we apply Theorem 2.3 of [Kob]. According to Lemma 6 below, for fixed $z \in \mathbb{R}^m$, $(f(t, z))_{t \in [0, T]}$ defines a predictable process. A sufficient condition for the existence of a solution is condition (H1) in [Kob]: there are constants c_0, c_1 such that

$$|f(t, z)| \leq c_0 + c_1|z|^2 \quad \text{for all } z \in \mathbb{R}^n \quad P - a.s. \quad (8)$$

By means of (3) we get for $z \in \mathbb{R}^m, t \in [0, T]$

$$\text{dist}^2\left(z + \frac{1}{\alpha}\theta_t, C_t\right) \leq 2|z|^2 + 2\left(\frac{1}{\alpha}|\theta_t| + k_1\right)^2.$$

So (8) follows from the boundedness of θ . Theorem 2.3 in [Kob] states that the BSDE (6) possesses at least one solution $(Y, Z) \in \mathcal{H}^\infty(\mathbb{R}) \times \mathcal{H}^2(\mathbb{R}^m)$.

To prove uniqueness, suppose that solutions $(Y^1, Z^1), (Y^2, Z^2)$ of the BSDE are given. Then we have

$$Y^1 - Y^2 = - \int_{\cdot}^T (Z^1 - Z^2) dW - \int_{\cdot}^T (f(s, Z_s^1) - f(s, Z_s^2)) ds.$$

Now note that for $s \in [0, T], z^1, z^2 \in \mathbb{R}^m$ we may write

$$f(s, z^1) - f(s, z^2) = -\frac{\alpha}{2} [\text{dist}^2(z^1 + \frac{1}{\alpha}\theta_s, C_s) - \text{dist}^2(z^2 + \frac{1}{\alpha}\theta_s, C_s) + (z^1 - z^2)\theta_s].$$

Using the Lipschitz property of the distance function from a closed set we obtain the estimate

$$\begin{aligned} |f(s, z^1) - f(s, z^2)| &\leq c_1 |z^1 - z^2| + c_2 (|z^1| + |z^2|) (|z^1 - z^2|) \\ &\leq c_3 (1 + |z^1| + |z^2|) |z^1 - z^2|. \end{aligned}$$

Let us set

$$\beta(t) = \begin{cases} \frac{f(t, Z_t^1) - f(t, Z_t^2)}{Z_t^1 - Z_t^2}, & \text{if } Z_t^1 - Z_t^2 \neq 0, \\ 0, & \text{if } Z_t^1 - Z_t^2 = 0. \end{cases}$$

Then we obtain from the preceding estimate

$$|\beta(t)| \leq c(1 + |Z_t^1| + |Z_t^2|), \quad t \in [0, T].$$

Moreover, from the boundedness of F , the P -BMO property of $\int_0^{\cdot} Z^i(s) dW_s, i = 1, 2$, follows. This in turn entails that $\int_0^{\cdot} \beta(s) dW_s$ is a P -BMO martingale. But this allows us to give an alternative description of the difference of solutions in

$$\begin{aligned} Y^1 - Y^2 &= - \int_{\cdot}^T (Z_s^1 - Z_s^2) dW_s - \int_{\cdot}^T \beta(s) (Z_s^1 - Z_s^2) ds \\ &= - \int_{\cdot}^T (Z_s^1 - Z_s^2) [dW_s + \beta(s) ds]. \end{aligned}$$

This process is a martingale under the equivalent probability measure Q which has density

$$\mathcal{E}(- \int_0^T \beta(t) dW_t)$$

with respect to P (see Lemma 7). Since $Y_T^1 = F = Y_T^2$ we therefore conclude $Y^1 = Y^2$ and $Z^1 = Z^2$, and uniqueness is established.

To find the value function of our optimization problem, we proceed with a solution $(Y, Z) \in \mathcal{H}^\infty(\mathbb{R}) \times \mathcal{H}^2(\mathbb{R}^n)$ of (6). Let p^* denote the predictable process constructed in Lemma 6 for $a = Z + \frac{1}{\alpha}\theta$. By Lemma 7 below, $\int_0^{\cdot} p_s^* dW_s$ is a P -BMO-martingale and therefore $p^* \in \mathcal{A}$. Furthermore, $\tilde{A}_t^{(p^*)} = -1$ for $\lambda \otimes P$ almost all (t, ω) and with Lemma

7 and Theorem 2.3 in [Kaz], $\mathcal{E}(-\alpha \int (p_s^* - Z_s) dW_s)$ is a martingale with expectation 1. Hence $R^{(p^*, x)}$ is a martingale and

$$\begin{aligned} R_0^{(p^*)} &= E \left[-\exp \left(-\alpha \left(x + \int_0^T p_s^* (dW_s + \theta_s ds) - F \right) \right) \right] \\ &= -\exp(-\alpha(x - Y_0)). \end{aligned}$$

For $p \in \mathcal{A}$, $\mathcal{E}(-\alpha \int (p_s - Z_s) dW_s)$ is a martingale and $\tilde{A}^{(p)}$ is decreasing. Therefore

$$\begin{aligned} R_0^{(p)} &= -\exp(-\alpha(x - Y_0)) \\ &\geq E \left[-\exp \left(-\alpha \left(x + \int_0^T p_s (dW_s + \theta_s ds) - F \right) \right) \right]. \end{aligned}$$

The supremum in (4) is uniquely defined, hence the unique solution of (6) has the same initial value Y_0 . □

We can show that the strategy p^* chosen above is optimal in a wider sense. In fact, an investor who has chosen at time 0 the strategy p^* will stick to this decision if he starts solving the optimization problem at some later time between 0 and T . For this purpose, let us formulate the optimization problem more generally for a stopping time $\tau \leq T$ and an \mathcal{F}_τ -measurable random variable which describes the capital at time τ , i.e. $X_\tau = X_\tau^p$ for some $p \in \mathcal{A}$. So we consider the maximization problem

$$V(\tau, X_\tau) = \text{ess sup}_{p \in \mathcal{A}} E \left[-\exp \left(-\alpha \left(X_\tau + \int_\tau^T p_s (dW_s + \theta_s ds) - F \right) \right) \right]. \quad (9)$$

Proposition 4 (Dynamic Principle) *The value function $x \mapsto -\exp(-\alpha(x - y))$ satisfies the dynamic programming principle, i.e.*

$$V(\tau, X_\tau) = -\exp(-\alpha(X_\tau - Y_\tau))$$

for all stopping times $\tau \leq T$ where Y_τ belongs to a solution of the BSDE (6). An optimal strategy that attains the essential supremum in (9) is given by p^* , the optimal strategy constructed in Theorem 3.

Proof For $t \in [0, T]$, set

$$R_t = -\exp(-\alpha(X_t - Y_t)) \mathcal{E} \left(-\int_t^T \alpha(p_s - Z_s) dW_s \right) \exp \left(\int_t^T v(s, p_s, Z_s) ds \right)$$

and apply the optional stopping theorem to the stochastic exponential. The claim follows as in Theorem 3. □

Remark 5 *If the constraint C on the strategies is a convex cone, the value function V and the optimal strategy p^* both constructed in Theorem 3 are equivalent to those determined in [Sek] and [ER].*

Sekine considers the utility function $x \mapsto -\frac{1}{\alpha} \exp(-\alpha x)$. He obtains the value function

$$V(x) = -\frac{1}{\alpha} \exp(-\alpha x + \bar{Y}_0)$$

starting with the BSDE

$$\bar{Y}_t = \alpha F - \int_t^T \bar{z}_s dW_s - \int_t^T \bar{f}(s, \theta_s, \bar{z}_s) ds, \quad t \in [0, T],$$

where

$$\bar{f}(t, \theta_t, \bar{z}) = \theta_t \Pi_{C_t}(\bar{z} + \theta_t) - \frac{1}{2} |\bar{z} - \Pi_{C_t}(\bar{z} + \theta_t)|^2.$$

We evidently have to show that $\bar{Y}_t = \alpha Y_t$ for $t \in [0, T]$ or equivalently $\alpha f(t, \theta_t, \frac{z}{\alpha}) = \bar{f}(t, \theta_t, z)$. Note that for a convex set C , the projection $\Pi_C(a)$ is unique. If C is a convex cone and $\beta > 0$, then $\beta \Pi_C(a) = \Pi_C(\beta a)$. The equality for the functions f and \bar{f} therefore follows. El Karoui and Rouge [ER] have obtained the same BSDE and value function before Sekine.

In the following Lemma we return to a technical point in the proof of Theorem 3. We show that it is possible to define a predictable process which satisfies (7). Instead of referring to a classical section theorem, see Dellacherie and Meyer [DM], we prefer to give a direct and constructive proof.

Lemma 6 (measurable selection) *Let $(a_t)_{t \in [0, T]}$, $(\sigma_t)_{t \in [0, T]}$ be $\mathbb{R}^{1 \times m}$ - resp. $\mathbb{R}^{d \times m}$ -valued predictable stochastic processes, $\tilde{C} \subset \mathbb{R}^d$ a closed set and $C_t = \tilde{C} \sigma_t$, $t \in [0, T]$.*

(a) *The process*

$$d = (\text{dist}(a_t, \tilde{C} \sigma_t))_{t \in [0, T]}$$

is predictable.

(b) *There exists a predictable process a^* with*

$$a_t^* \in \Pi_{C_t}(a_t) \quad \text{for all } t \in [0, T].$$

Proof In order to prove (a), observe that d is the composition of continuous mappings with predictable processes. For $k \in \mathbb{N}$ let H^k denote the space of compact subsets of \mathbb{R}^k equipped with the Hausdorff metric and $\mathcal{B}(H^k)$ the Borel sigma algebra with respect to this metric. The mapping $\text{dist} : \mathbb{R}^m \times H^m \rightarrow \mathbb{R}$ is jointly continuous hence $\mathcal{B}(\mathbb{R}^m) \otimes \mathcal{B}(H^m)$ - $\mathcal{B}(\mathbb{R})$ measurable. Now consider $j : \mathbb{R}^{d \times m} \times H^d \rightarrow H^m$ that maps a compact subset \tilde{C} in \mathbb{R}^d by applying an $d \times m$ -matrix $\tilde{\sigma}$ to a compact subset \tilde{K} of \mathbb{R}^m . More formally, j maps \tilde{C} to the following set:

$$\tilde{K} = \{b \in \mathbb{R}^m \mid \exists \tilde{c} \in \tilde{C} : b = \tilde{c} \tilde{\sigma}\}.$$

The mapping j is also jointly continuous and therefore $\mathcal{B}(\mathbb{R}^{m \times d}) \otimes \mathcal{B}(H^d)$ - $\mathcal{B}(H^m)$ -measurable. Hence (a) follows for compact \tilde{C} .

If more generally \tilde{C} is closed but not bounded, take $\tilde{C}_n = \tilde{C} \cap B_n$ where B_n is the closed ball with radius n centered at the origin. According to what has already

been shown, for $n \in \mathbb{N}$, $\text{dist}(a_t, \tilde{C}_n \sigma_t)$ defines a predictable process and $\text{dist}(a_t, \tilde{C}_n \sigma_t)$ converges to $\text{dist}(a_t, \tilde{C} \sigma_t)$, for $n \rightarrow \infty$. This proves the first claim.

In order to prove the second claim, we first concentrate on the case of compact \tilde{C} . We have to show that for $z \in \mathbb{R}^m$ and a compact set $\tilde{K} \subset \mathbb{R}^m$ there exists a $\mathcal{B}(\mathbb{R}^m) \otimes \mathcal{B}(H^m) - \mathcal{B}(\mathbb{R}^m)$ -measurable mapping $\xi(z, \tilde{K})$ with $\xi(z, \tilde{K}) \in \Pi_{\tilde{K}}(z)$. This is achieved by the definition of a sequence of mappings $\xi_n(z, \tilde{K})$ with a subsequence of randomly chosen index that converges to an element of $\Pi_{\tilde{K}}(z)$. The choice of the converging subsequence will depend in a measurable way on z and \tilde{K} .

For $n \in \mathbb{N}$, let $G_n = (x_i^n)_{i \in \mathbb{N}}$ be a dyadic grid with $\min_{x \in G_n} \text{dist}(\bar{z}, x) \leq \frac{1}{n}$ for all $\bar{z} \in \mathbb{R}^m$. Let the elements of the grid G_n be numbered by $G_n = \{g_i^n : i \in \mathbb{N}\}$. Let \tilde{K}_n be the elements of the grid with distance at most $\frac{1}{n}$ from G_n . Since we can describe the sets \tilde{K}_n as the intersections of the discrete set G_n with the closed set of all points in \mathbb{R}^m having distance at most $\frac{1}{n}$ from \tilde{K} , and this closed set depends continuously on \tilde{K} , \tilde{K}_n is measurable in \tilde{K} . For any $z \in \mathbb{R}^m$, let $\Pi_n(z, \tilde{K})$ be the set of all points in \tilde{K}_n with minimal distance from z . Since \tilde{K}_n is measurable in \tilde{K} , $\Pi_n(z, \tilde{K})$ is obviously measurable in (z, \tilde{K}) . To define $\xi_n(z, \tilde{K})$, we have to choose one point in $\Pi_n(z, \tilde{K})$. Let it be the one with minimal index in the enumeration of G_n . This choice preserves the measurability in (z, \tilde{K}) . Hence we obtain that $\xi_n(z, \tilde{K})$ is $\mathcal{B}(\mathbb{R}^m) \otimes \mathcal{B}(H^m) - \mathcal{B}(\mathbb{R}^m)$ -measurable.

We next have to choose the subsequence with random indices. This is done as in Lemma 1.55 in [FS]. Let us argue with respect to an arbitrary probability measure Q on the Borel sets of $\mathbb{R}^m \times H^m$. First note that $\eta = \liminf_{n \rightarrow \infty} |\xi_n| < \infty$. Let $\tau_1 = 1$, and for $n > 1$

$$\tau_n = \inf \{ m > \tau_{n-1} : |\xi_m - \eta| \leq \frac{1}{n} \}.$$

Then it is plain that τ_n is $\mathcal{B}(\mathbb{R}^m) \otimes \mathcal{B}(H^m) - \mathcal{B}(\mathbb{R}^m)$ -measurable. Now take

$$\xi = \liminf_{n \rightarrow \infty} \xi_{\tau_n},$$

which obviously inherits the measurability properties of the subsequence with random indices. But ξ is a selection. Indeed, for every $n \in \mathbb{N}$, $\text{dist}(\xi_{\tau_n}(z, \tilde{K}), \Pi_{\tilde{K}}(z)) \leq \frac{1}{\tau_n}$. Thus by construction, $\xi(z, \tilde{K}) \in \Pi_{\tilde{K}}(z)$ for all $(z, \tilde{K}) \in \mathbb{R}^m \times H^m$.

We may then choose

$$a^* = \xi(a, C\sigma)$$

to satisfy the requirements of the second part of the assertion in the compact case.

Finally, if C is only closed, we may proceed as in the proof for (a).

□

Lemma 7 *Let Z be the second component of a solution of the BSDE (6), and let p^* be given by Lemma 6 for $a = Z + \frac{1}{\alpha}\theta$. Then the processes*

$$\int_0^\cdot Z_s dW_s, \quad \int_0^\cdot p_s^* dW_s$$

are P -BMO martingales.

Proof Due to Corollary 2.2 of Kobylanski, the process Y corresponding to Z in the solution of the BSDE (6) is uniformly bounded P -a.s. Let k denote the upper bound. Applying Itô's formula to the non positive process $Y - k$, we obtain for stopping times $\tau \leq T$

$$\begin{aligned} E \left[\int_{\tau}^T Z_s^2 ds \middle| \mathcal{F}_{\tau} \right] &= E[(H - k)^2 | \mathcal{F}_{\tau}] - |Y_{\tau} - k|^2 \\ &\quad - 2E \left[\int_{\tau}^T (Y_s - k) f(s, Z_s) ds \middle| \mathcal{F}_{\tau} \right] \end{aligned}$$

The definition of f yields for all $(t, z) \in [0, T] \times \mathbb{R}^m$

$$f(t, z) \leq z\theta_t + \frac{1}{2\alpha}|\theta_t|^2.$$

Therefore there exist positive constants c_1, c_2 and \tilde{c}_1 such that

$$\begin{aligned} E \left[\int_{\tau}^T |Z_s|^2 ds \middle| \mathcal{F}_{\tau} \right] &\leq c_1 + c_2 E \left[\int_{\tau}^T |Z_s + 1| ds \middle| \mathcal{F}_{\tau} \right] \\ &\leq \tilde{c}_1 + \frac{1}{2} E \left[\int_{\tau}^T |Z_s|^2 ds \middle| \mathcal{F}_{\tau} \right]. \end{aligned}$$

Hence, $\int_0^{\cdot} Z_s dW_s$ is a BMO-martingale.

We next deal with the stochastic integral process of p^* . The triangle inequality implies

$$|p^*| \leq |Z + \frac{1}{\alpha}\theta| + |p^* - (Z + \frac{1}{\alpha}\theta)|.$$

The definition of p^* together with (3) yields for some constants k_1, k_2

$$|p_t^*| \leq 2|Z_t| + \frac{2}{\alpha}|\theta_t| + k_1 \leq 2|Z_t| + k_2, \quad t \in [0, T],$$

and thus for every stopping time $\tau \leq T$

$$E \left[\int_{\tau}^T |p_t^*|^2 dt \middle| \mathcal{F}_{\tau} \right] \leq E \left[\int_{\tau}^T 8|Z_t|^2 dt + 2Tk_2^2 \middle| \mathcal{F}_{\tau} \right].$$

This implies the P -BMO property of $\int_0^{\cdot} p_s^* dW_s$.

□

3 Power utility

In this section we calculate the value function and characterize the optimal strategy for the utility maximization problem with respect to

$$U_{\gamma}(x) = \frac{1}{\gamma}x^{\gamma}, \quad x \geq 0, \quad \gamma \in (0, 1).$$

This time, our investor maximizes the expected utility of his wealth at time T without an additional liability. The trading strategies are constraint to take values in a closed set $\tilde{C}_2 \subseteq \mathbb{R}^d$. In this section, we shall use a somewhat different notion of trading strategy: $\tilde{\rho} = (\tilde{\rho}^i)_{i=1,\dots,d}$ denotes the part of the wealth invested in stock i . The number of shares of stock i is given by $\frac{\tilde{\rho}_t^i X_t}{S_t^i}$. A d -dimensional \mathbb{F} -predictable process $\tilde{\rho} = (\tilde{\rho}_t)_{0 \leq t \leq T}$ is called trading strategy (part of wealth) if the following wealth process is well defined:

$$X_t^{(\tilde{\rho})} = x + \int_0^t \sum_{i=1}^d \frac{X_s^{(\tilde{\rho})} \tilde{\rho}_{i,s}}{S_{i,s}} dS_{i,s} = x + \int_0^t X_s^{(\tilde{\rho})} \tilde{\rho}_s \sigma_s (dW_s + \theta_s ds), \quad (10)$$

and the initial capital x is positive. The wealth process $X^{(\tilde{\rho})}$ can be written as:

$$X_t^{(\tilde{\rho})} = x \mathcal{E} \left(\int_t \tilde{\rho}_s \sigma_s (dW_s + \theta_s ds) \right), \quad t \in [0, T].$$

As before, it is more convenient to introduce

$$\rho_t = \tilde{\rho}_t \sigma_t, \quad t \in [0, T].$$

Accordingly, ρ is constraint to take its values in

$$C_t(\omega) = \tilde{C} \sigma_t(\omega) \quad t \in [0, T], \omega \in \Omega.$$

The sets C_t satisfy (3). In order to formulate the optimization problem we first define the set of admissible trading strategies.

Definition 8 *The set of admissible trading strategy $\tilde{\mathcal{A}}$ consists of all d -dimensional predictable processes $\rho = (\rho_t)_{0 \leq t \leq T}$ that satisfy $\rho_t \in C_t(\omega)$ P -a.s and $\int_0^\cdot \rho_s dW_s$ is a BMO-martingale.*

The investor faces the maximization problem

$$\bar{V}(x) = \sup_{\tilde{\rho} \in \tilde{\mathcal{A}}} E \left[U \left(X_T^{(\tilde{\rho})} \right) \right]. \quad (11)$$

In order to find the value function and an optimal strategy we apply the same method as for the exponential utility function. We therefore have to construct a stochastic process $\tilde{R}^{(\rho)}$ with terminal value

$$\tilde{R}_T^{(\rho)} = U \left(x + \int_0^T X_s \rho_s \frac{dS_s}{S_s} \right),$$

and an initial value $\tilde{R}_0^{(\rho)} = \tilde{R}_0^x$ that does not depend on ρ , $\tilde{R}^{(\rho)}$ is a supermartingale for all $\rho \in \tilde{\mathcal{A}}$ and a martingale for a $\rho^* \in \tilde{\mathcal{A}}$. Then ρ^* is the optimal strategy and the value function given by $\bar{V}(x) = \tilde{R}_0^x$. Applying the utility function to the wealth process yields

$$(X_t^{\rho^*,x})^\gamma = x^\gamma \exp \left(\int_0^t \gamma \rho_s dW_s + \int_0^t \gamma \rho_s \theta_s ds - \frac{1}{2} \int_0^t \gamma |\rho_s|^2 ds \right), \quad t \in [0, T].$$

This equation suggests the following choice:

$$\tilde{R}_t^{(\rho)} = x^\gamma \exp \left(\int_0^t \gamma \rho_s dW_s + \int_0^t \gamma \rho_s \theta_s ds - \frac{1}{2} \int_0^t \gamma |\rho_s|^2 ds + Y_t \right), \quad (12)$$

where (Y, Z) is a solution of the BSDE

$$Y_t = 0 - \int_t^T Z_s dW_s - \int_t^T f(s, Z_s) ds, \quad t \in [0, T].$$

In order to get the supermartingale property of $\tilde{R}^{(\rho)}$ we have to construct $f(t, z)$ such that for $t \in [0, T]$

$$\gamma \rho_t \theta_t - \frac{1}{2} \gamma |\rho_t|^2 + f(t, Z_t) \leq -\frac{1}{2} |\gamma \rho_t + Z_t|^2 \quad \text{for all } \rho \in \tilde{\mathcal{A}}. \quad (13)$$

$\tilde{R}^{(\rho^*)}$ will even be a martingale if equality holds for $\rho^* \in \tilde{\mathcal{A}}$. This is equivalent to

$$f(t, Z_t) \leq \frac{1}{2} \gamma (1 - \gamma) \left| \rho_t - \frac{1}{1 - \gamma} (Z_t + \theta_t) \right|^2 - \frac{1}{2} \frac{\gamma |Z_t + \theta_t|^2}{1 - \gamma} - \frac{1}{2} |Z_t|^2.$$

Hence the appropriate choice for f is

$$f(t, z) = \frac{\gamma(1 - \gamma)}{2} \text{dist}^2 \left(\frac{1}{1 - \gamma} (z + \theta_t), C_t \right) - \frac{\gamma |z + \theta_t|^2}{2(1 - \gamma)} - \frac{1}{2} |z|^2,$$

and a candidate for the optimal strategy must satisfy

$$\rho_t^* \in \Pi_{C_t(\omega)} \left(\frac{1}{1 - \gamma} (Z_t + \theta_t) \right), \quad t \in [0, T].$$

In the following Theorem both value function and optimal strategy are described.

Theorem 9 *The value function of the optimization problem is given by*

$$V(x) = x^\gamma \exp(Y_0) \quad \text{for } x > 0,$$

where Y_0 is defined by a solution (Y, Z) of the BSDE

$$Y_t = 0 - \int_t^T Z_s dW_s - \int_t^T f(s, Z_s) ds, \quad t \in [0, T], \quad (14)$$

with

$$f(t, z) = \frac{\gamma(1 - \gamma)}{2} \text{dist}^2 \left(\frac{1}{1 - \gamma} (z + \theta_t), C_t \right) - \frac{\gamma |z + \theta_t|^2}{2(1 - \gamma)} - \frac{1}{2} |z|^2.$$

There exists an optimal trading strategy $\rho^* \in \tilde{\mathcal{A}}$ with the property

$$\rho_t^* \in \Pi_{C_t(\omega)} \left(\frac{1}{1 - \gamma} (Z_t + \theta_t) \right). \quad (15)$$

Proof According to Lemma 6, $(f(t, z))_{t \in [0, T]}$ is a predictable stochastic process which also depends on σ . Due to (3) and the boundedness of θ , Condition (H1) for Theorem 2.3 in [Kob] is fulfilled. We obtain the existence of a solution $(Y, Z) \in \mathcal{H}^\infty(\mathbb{R}) \times \mathcal{H}^2(\mathbb{R}^m)$ for the BSDE (14). Uniqueness follows from the comparison arguments in the uniqueness part of the proof of Theorem 3.

Let ρ^* denote the predictable process constructed with Lemma 6 for $a = \frac{1}{1-\gamma}(Z + \theta)$. Lemma 12 below shows that $\rho^* \in \tilde{\mathcal{A}}$. By Theorem 2.3 in [Kaz], the process $\tilde{R}^{(\rho^*)}$ is a martingale with terminal value

$$\tilde{R}_T^{(\rho^*)} = x^\gamma \exp \left(\int_0^T \gamma \rho_s^* dW_s + \int_0^T \gamma \rho_s^* \theta_s ds - \frac{1}{2} \int_0^T \gamma |\rho_s^*|^2 ds \right).$$

This is the power utility from terminal wealth of the trading strategy ρ^* . Therefore the expected utility of ρ^* is equal to $\tilde{R}_0^{(\rho^*, x)} = x^\gamma \exp(Y_0)$.

To show that this provides the value function let $\rho \in \tilde{\mathcal{A}}$. (13) yields

$$\tilde{R}_t^{(\rho)} = x^\gamma \exp(Y_0) \mathcal{E} \left(\int (\gamma \rho_s + Z_s) dW_s \right)_t \exp \left(\int_0^t v_s ds \right), \quad t \in [0, T],$$

for a process v with $v_s \leq 0$ $\lambda \otimes P$ a.s. The process in the stochastic exponential is a BMO–martingale, whence the stochastic exponential is a martingale. Therefore $R^{(\rho, x)}$ is a supermartingale. The terminal value $R_T^{(\rho, x)}$ is the utility of the terminal wealth of the trading strategy ρ . Consequently

$$E[U(X_T^{(\rho, x)})] \leq R_0^{(x)} = x^\gamma \exp(Y_0) \quad \text{for all } \rho \in \mathcal{A}.$$

The supremum in the maximization problem (11) is a unique real number. Therefore the initial value Y_0 for every solution of the BSDE (14) must be the same. □

Again we can show that an investor starting to act at some stopping time in the trading interval $[0, T]$ will perceive the strategy ρ^* just constructed as optimal. Let $\tau \leq T$ denote a stopping time and X_τ an \mathcal{F}_τ –measurable random variable which describes the capital at time τ , i.e. $X_\tau = X_\tau^\rho$ for a $\rho \in \tilde{\mathcal{A}}$ and an initial capital $x > 0$. Consider the maximization problem

$$\bar{V}(\tau, X_\tau) = \text{ess sup}_{p \in \mathcal{A}_\tau} E \left[U \left(X_\tau + \int_\tau^T X_s \rho_s (dW_s + \theta_s ds) \right) \right]. \quad (16)$$

Proposition 10 (Dynamic Principle) *The value function $x^\gamma \exp(y)$ satisfies the dynamic programming principle, i.e.*

$$\bar{V}(\tau, X_\tau) = (X_\tau)^\gamma \exp(Y_\tau)$$

for all stopping times $\tau \leq T$, where Y_τ is given by a solution of the BSDE (14). An optimal strategy which attains the essential supremum in (16) is given by ρ^* constructed in Theorem 9.

Proof See Proposition 4.

Remark 11 Suppose that the constraint set C is a convex cone. Then the optimal strategy ρ^* constructed in Theorem 9 is the same as in [Sek].

Sekine uses the utility function $x \mapsto \frac{1}{\gamma}x^\gamma$ and obtains the value function

$$\tilde{V}(x) = \frac{1}{\gamma}x^\gamma \exp((1-\gamma)\tilde{Y}_0),$$

where \tilde{Y}_0 is defined by the solution of the BSDE

$$\tilde{Y}_t = 0 - \int_t^T \tilde{Z}_s dW_s - \int_t^T g(s, \tilde{Z}_s) ds, \quad t \in [0, T].$$

Here

$$g(t, \tilde{z}) = \frac{|\theta_t|^2}{2} - \frac{1}{2} \left| \theta_t - \Pi_{C_t} \left(\tilde{z} + \frac{\theta_t}{1-\gamma} \right) \right|^2 - \frac{1-\gamma}{2} \left| \tilde{z} - \Pi_{C_t} \left(\tilde{z} + \frac{\theta_t}{1-\gamma} \right) \right|^2.$$

As for the exponential utility function we have to show $(1-\gamma)\tilde{Y} = Y$ or equivalently $(1-\gamma)g(t, \frac{z}{1-\gamma}) = f(t, z)$. In fact, we have

$$\begin{aligned} (1-\gamma)g\left(t, \frac{z}{1-\gamma}\right) &= (1-\gamma) \left[\frac{|\theta_t|^2}{2} - \frac{1}{2} \left| \theta_t - \Pi_{C_t} \left(\frac{z+\theta_t}{1-\gamma} \right) \right|^2 \right] \\ &\quad - \frac{(1-\gamma)^2}{2} \left| \frac{z}{1-\gamma} - \Pi_{C_t} \left(\frac{z+\theta_t}{1-\gamma} \right) \right|^2 \\ &= \theta_t \Pi_{C_t}(z+\theta_t) - \frac{1}{2(1-\gamma)} |\Pi_{C_t}(z+\theta_t)|^2 \\ &\quad - \frac{1}{2} |z|^2 + z \Pi_{C_t}(z+\theta_t) - \frac{1}{2} |\Pi_{C_t}(z+\theta_t)|^2 \\ &= (z+\theta_t) \Pi_{C_t}(z+\theta_t) - \frac{2-\gamma}{2(1-\gamma)} |\Pi_{C_t}(z+\theta_t)|^2 - \frac{1}{2} |z|^2 \\ &= -\frac{\gamma}{2(1-\gamma)} |\Pi_{C_t}(z+\theta_t)|^2 - \frac{1}{2} |z|^2. \end{aligned}$$

To obtain the last equality, we use

$$(z+\theta_t) \Pi_{C_t}(z+\theta_t) = |\Pi_{C_t}(z+\theta_t)|^2$$

(see (17) below).

For the function f we obtain

$$\begin{aligned} f(t, z) &= \frac{\gamma(1-\gamma)}{2} \left| \frac{1}{1-\gamma}(z+\theta_t) - \Pi_{C_t} \left(\frac{1}{1-\gamma}(z+\theta_t) \right) \right|^2 \\ &\quad - \frac{\gamma}{2} \frac{(z+\theta_t)^2}{(1-\gamma)} - \frac{1}{2} |z|^2 \\ &= -\frac{\gamma}{1-\gamma} (z+\theta_t) \Pi_{C_t}(z+\theta_t) + \frac{\gamma}{2(1-\gamma)} |\Pi_{C_t}(z+\theta_t)|^2 - \frac{1}{2} |z|^2 \\ &= -\frac{\gamma}{2(1-\gamma)} |\Pi_{C_t}(z+\theta_t)|^2 - \frac{1}{2} |z|^2. \end{aligned}$$

For $t \in [0, T]$, $z \in \mathbb{R}^m$ we therefore have

$$(1 - \gamma)g(t, \frac{z}{1 - \gamma}) = f(t, z).$$

It remains to prove that for a convex cone C and $a \in \mathbb{R}^m$ the following equality holds:

$$\Pi_C(a)(a - \Pi_C(a)) = 0. \tag{17}$$

Let $1 \neq \lambda \geq 0$. For a convex cone C we have $\Pi_C(\lambda a) = \lambda \Pi_C(a)$ and

$$\begin{aligned} |a - \Pi_C(a)|^2 &\leq |a - \lambda \Pi_C(a)|^2, \\ a^2 - 2a\Pi_C(a) + |\Pi_C(a)|^2 &\leq a^2 - 2\lambda a\Pi_C(a) + \lambda^2|\Pi_C(a)|^2. \end{aligned}$$

Thus

$$2a(\lambda - 1)\Pi_C(a) - (\lambda + 1)(\lambda - 1)|\Pi_C(a)|^2 \leq 0,$$

and

$$(\lambda - 1)\Pi_C(a) [2a - (\lambda + 1)\Pi_C(a)] \leq 0.$$

For $\lambda > 1$ we obtain

$$\Pi_C(a)(2a - (\lambda + 1)\Pi_C(a)) \leq 0,$$

hence for $\lambda \rightarrow 1^+$

$$\Pi_C(a)(a - \Pi_C(a)) \leq 0.$$

For $\lambda < 1$ we get

$$\Pi_C(a)(2a - (\lambda + 1)\Pi_C(a)) \geq 0,$$

and $\lambda \rightarrow 1^-$ yields

$$\Pi_C(a)(a - \Pi_C(a)) \geq 0.$$

□

Lemma 12 *Let Z be the second component of a solution of the BSDE (14), and let ρ^* be given by (15). Then the processes*

$$\int_0^\cdot Z_s dW_s, \quad \int_0^\cdot \rho_s^* dW_s$$

are P -BMO martingales.

Proof We can use the same line of reasoning as in the proof of Lemma 7. The argument given there has to be slightly modified, however. We may take a lower bound k for Y , and apply Itô's formula to $|Y - k|^2$, to conclude in the same manner as before.

□

4 Log Utility

To complete the spectrum of important utility functions, in this section we shall consider logarithmic utility. As in the preceding section, the agent has no liability at time T . Trading strategies and wealth process have the same meaning as in section 3 (see Definition 8 and (10)). The trading strategies $\tilde{\rho}$ are constraint to take values in a closed set $\tilde{C}_2 \subset \mathbb{R}^d$. For $\rho_t = \tilde{\rho}_t \sigma_t$ the constraints are described by $C_t = \tilde{C}_2 \sigma_t, t \in [0, T]$.

For the logarithmic utility function

$$U(x) = \log(x), \quad x > 0,$$

we obtain a particularly simple BSDE that leads to the value function and the optimal strategy. The optimization problem is given by

$$V(x) = \sup_{\rho \in \tilde{\mathcal{A}}} E[\log(X_T^{(\rho)})] \quad (18)$$

$$= \log(x) + \sup_{\rho \in \tilde{\mathcal{A}}} E \left[\int_0^T \rho_s dW_s + \int_0^T \left(\rho_s \theta_s - \frac{1}{2} |\rho_s|^2 \right) ds \right], \quad (19)$$

where the initial capital x is positive again. As in section 2 we want to determine a process $R^{(\rho)}$ with $R_T^{(\rho)} = \log(X_T^{(\rho)})$, and an initial value that does not depend on ρ . Furthermore, $R^{(\rho)}$ is a supermartingale for all $\rho \in \tilde{\mathcal{A}}$, and there exists a $\rho^* \in \tilde{\mathcal{A}}$ such that $R^{(\rho^*)}$ is a martingale. The strategy ρ^* is the optimal strategy and R_0^* is the value function of the optimization problem (18).

We can choose for $t \in [0, T]$

$$R_t^{(\rho)} = \log x + Y_0 + \int_0^t (\rho_s + Z_s) dW_s + \int_0^t \left(-\frac{1}{2} |\rho_s - \theta_s|^2 + \frac{1}{2} \theta_s^2 + f(s) \right) ds,$$

where

$$f(t) = \frac{1}{2} \text{dist}^2(\theta_t, C_t) - \frac{1}{2} |\theta_t|^2, \quad t \in [0, T],$$

and (Y_t, Z_t) is the solution of the following BSDE:

$$Y_t = 0 - \int_t^T Z_s dW_s - \int_t^T f(s) ds, \quad t \in [0, T].$$

The initial value Y_0 satisfies

$$Y_0 = -E \left[\int_0^T f(s) ds \right].$$

Hence

$$V(x) = R_0^*(x) = \log(x) + E \left[- \int_0^T f(s) ds \right].$$

An optimal trading strategy can be constructed by means of Lemma 6. It guarantees the existence of an admissible trading strategy ρ^* which satisfies $\rho_t^* \in \Pi_{C_t}(\theta_t)$. In particular ρ^* only depends on θ and σ , and the set \tilde{C} describing the constraints on the trading strategies.

References

- [B] Bismut, J.-M. “Conjugate convex functions in optimal stochastic control.” *J. Math. Anal. Appl.* 44 (1973), 383-404.
- [CK] Cvitanic, J.; Karatzas, I. “Convex Duality in Constrained Portfolio Optimization”. *Ann. Appl. Prob.* 2(4), 767–818 (1992).
- [CSW] Cvitanic, J.; Schachermayer, W.; Wang, H. “Utility maximization in incomplete markets with random endowment” *Finance and Stochastics* 5, 259–272 (2001)
- [DGR] Delbaen, F.; Grandits, P.; Rheinländer, T.; Samperi, D.; Schweizer, M.; Stricker, C. “Exponential Hedging and Entropic Penalties”. *Mathematical Finance* 12, 99–123 (2002).
- [DM] Dellacherie, C., Meyer, P.-A. “Probabilités et potentiel. Ch. I-IV. Paris: Hermann 1975.
- [EPQ] El Karoui, N.; Peng, S.; Quenez, M.C. “Backward stochastic differential equations in finance.” *Math. Finance* 7 (1997), 1-71.
- [ER] El Karoui, N.; Rouge, R. “Pricing via utility maximization and entropy.” *Math. Fin.* 10(2), 259–276 (2000).
- [FS] Föllmer, H.; Schied, A. “Stochastic Finance. An Introduction in Discrete Time”. *Studies in Mathematics* 27, de Gruyter, Berlin, New York, 2002.
- [Kaz] Kazamaki, N. “Continuous Exponential Martingales and BMO”. *Lecture Notes in Mathematics* 1579, Springer, Berlin, 1994.
- [KLS] Karatzas, I.; Lehoczky, J. P.; Shreve, S. E. “Optimal portfolio and consumption decisions for a small investor on a finite horizon”. *SIAM J. Control Optimization* 25, 1557-1586 (1987).
- [KS] Kramkov, D., Schachermayer, W. “The asymptotic elasticity of utility functions and optimal investment in incomplete markets.” *Ann. Appl. Probability* 9 (1999), 904-950.
- [Kob] Kobylanski, M. 2000, “Backward Stochastic Differential Equations and Partial Differential Equations with Quadratic Growth”. *Ann. Prob.* 28(2) (2000), 558–602.
- [PP] Pardoux, E.; Peng, S.G. “Adapted solution of a backward stochastic differential equation.” *Syst. Control Lett.* 14 (1990), 55-61.
- [Sek] Sekine, J. “Exponential hedging by solving a backward stochastic differential equation: an alternative approach.” *Preprint, Osaka University* (2002).

[Sch] Schachermayer, W. "Optimal investment in incomplete financial markets."
Mathematical Finance Bachelier Congress 2000, eds.: Geman, H.; Madan,
D.; Pliska, St. R.; Vorst, T. Springer, 427–462 (2001)