

# On algebraic multilevel methods for non-symmetric systems - convergence results

C. Mense \*      R. Nabben<sup>†</sup>

February 28, 2008

## Abstract

Here we analyze algebraic multilevel methods applied to non-symmetric M-matrices. We consider two types of multilevel approximate block factorizations. The first one is related to the AMLI method. The second method is the multiplicative counterpart of the AMLI approach which we call multiplicative algebraic multilevel method, the MAMLI method. The MAMLI method is closely related to certain geometric and algebraic multigrid methods like the AMGr method. Although these multilevel methods work very well in practice for many problems, there is not that much known about theoretical convergence properties for non-symmetric problems. Here, we establish convergence results and comparison results between AMLI and MAMLI multilevel methods applied to non-symmetric M-matrices.

**Keywords.** Algebraic multilevel methods, multilevel approximate block factorizations, algebraic multigrid methods, AMLI method

**AMS subject classifications.** 65F10, 65F50, 65N22

## 1 Introduction

In many recent papers algebraic multigrid methods or multilevel methods were designed to solve large sparse linear systems by using only information on the matrix structure and the matrix entries. Among several algebraic methods the algebraic multigrid method (AMG) and the multilevel approximate block factorization are best-known. The pioneering work on algebraic multilevel methods was done by Brandt, McCormick and Ruge [12] and Ruge and Stüben [32, 34] by introducing the AMG method in the eighties, see also [35].

Recently, a theoretical comparison of different algebraic multigrid methods applied to symmetric positive definite systems was given by Notay in [29]. However, there are not that many theoretical results known for algebraic multigrid methods applied to non-symmetric matrices.

Here we analyze algebraic multilevel methods applied to non-symmetric M-matrices. Algebraic multilevel methods are often used as preconditioners for Krylov subspace methods. But here we focus on the convergence of these methods used as solvers.

---

\*Institut für Mathematik, MA 3-3, Technische Universität Berlin, Strasse des 17. Juni 136, D-10623 Berlin, Germany, phone: +49 30 314 29293, fax: +49 30 314 29621 ([mense@math.tu-berlin.de](mailto:mense@math.tu-berlin.de))

<sup>†</sup>Institut für Mathematik, MA 3-3, Technische Universität Berlin, Strasse des 17. Juni 136, D-10623 Berlin, Germany, phone: +49 30 314 29291, fax: +49 30 314 29621 ([nabben@math.tu-berlin.de](mailto:nabben@math.tu-berlin.de))

M-matrices occur in various fields of applied mathematics such as numerical analysis, probability, economics and operations research [9]. Moreover, Markov chain modeling became relevant in several applications from computer science such as information retrieval [20]. Iterative solvers, like algebraic multigrid methods are used to compute the steady state solution of a Markov chain, i.e. algebraic multigrid methods are used to find the solution of a system with non-symmetric M-matrix structure – a non-trivial task given the size of the Google matrix for example (see e.g. [20]).

Most of the algebraic multilevel methods start with a partitioning of the unknowns into fine and coarse grid unknowns. Related to this ordering, the  $n \times n$  system matrix  $A$  can be permuted in a block  $2 \times 2$  form

$$PAP^T = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}.$$

Here  $F$  denotes the set of fine grid unknowns, and  $C$  denotes the set of coarse grid unknowns with  $|F| = n_F$  and  $|C| = n_C$ . This process is called coarsing. There are a lot of different algorithms and strategies to perform the above partitioning, see e.g. [32, 22, 11, 39, 30]. The choice of the partitioning has a major influence on the convergence behavior of AMG methods. But here we assume that the coarsing is done already in an arbitrary way. Thus, we assume that  $A$  is given by

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}. \quad (1)$$

If the submatrix  $A_{FF}$  is nonsingular,  $A$  can be factorized as

$$A = \begin{bmatrix} I_F & 0 \\ A_{CF}A_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} A_{FF} & 0 \\ 0 & S \end{bmatrix} \begin{bmatrix} I_F & A_{FF}^{-1}A_{FC} \\ 0 & I_C \end{bmatrix},$$

where

$$S := (A/A_{FF}) := A_{CC} - A_{CF}A_{FF}^{-1}A_{FC}$$

is the Schur complement. If we now use an approximation  $\tilde{A}_{FF}$  of  $A_{FF}$  and an approximation  $\tilde{S}$  of  $S$ , or approximations of the inverses of these matrices, we obtain the matrix  $M$  with

$$M = \begin{bmatrix} I_F & 0 \\ A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} \tilde{A}_{FF} & 0 \\ 0 & \tilde{S} \end{bmatrix} \begin{bmatrix} I_F & \tilde{A}_{FF}^{-1}A_{FC} \\ 0 & I_C \end{bmatrix}. \quad (2)$$

This factorization is known as an approximate two-level block factorization [29]. Various multilevel methods use this two-level block approximate factorization as a major tool (see e.g. [4, 5, 2, 27, 7, 33] and references in [29]). One of these methods is the AMLI method introduced by Axelsson and Vassilevski in [4, 5]. With the use of matrix  $M$  in (2) one can define the stationary iteration given by the iteration matrix

$$T_{AMLI} = I - M^{-1}A. \quad (3)$$

Here the subscript AMLI is chosen with respect to the first use of this approximate block factorization by Axelsson and Vassilevski in the AMLI method. We point out that the

AMLI method itself contains a lot more ingredients like e.g. polynomial acceleration. Moreover, the AMLI method is designed originally in the framework of hierarchical bases to precondition the CG method.

For the iteration matrix  $T_{AMLI}$  in (3) we obtain

$$\begin{aligned} T_{AMLI} &= I - \left( \begin{bmatrix} I_F & 0 \\ A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} \tilde{A}_{FF} & 0 \\ 0 & \tilde{S} \end{bmatrix} \begin{bmatrix} I_F & \tilde{A}_{FF}^{-1}A_{FC} \\ 0 & I_C \end{bmatrix} \right)^{-1} A \\ &= I - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} A \\ &\quad - \begin{bmatrix} I_F \\ 0 \end{bmatrix} \tilde{A}_{FF}^{-1} \begin{bmatrix} I_F & 0 \end{bmatrix} A. \end{aligned}$$

Using the restriction and prolongation operators

$$\tilde{R} := [-A_{CF}\tilde{A}_{FF}^{-1} \ I_C], \quad \tilde{P}^T := \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix}, \quad \hat{R}^T := \begin{bmatrix} I_F \\ 0 \end{bmatrix} \quad (4)$$

and the matrices

$$P_1 := \tilde{P}^T \tilde{S}^{-1} \tilde{R}A \quad \text{and} \quad P_2 := \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}A \quad (5)$$

one obtains

$$T_{AMLI} = I - \tilde{P}^T \tilde{S}^{-1} \tilde{R}A - \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}A = I - P_1 - P_2. \quad (6)$$

Thus,  $T_{AMLI}$  can be written as an additive Schwarz method with inexact local solves (see [41, 19, 36] for details about Schwarz methods).

Hence, on one hand the matrices  $T_{AMLI}$  and  $M$  in (2) are constructed by an inexact block factorization, i.e. by using a *product* of matrices. Thus, the iteration given by  $T_{AMLI}$  can be seen as a multiplicative method. Moreover, there are also some other versions of the original AMLI method. These methods do not use the inexact block factorization in (2) and are based on a block-diagonal preconditioner. Therefore, these AMLI versions are called additive methods [1, 3].

But on the other hand  $T_{AMLI}$  can be written as an *additive* Schwarz method. To avoid confusion here, we will use the terminology *AMLI approach* for the inexact block factorization in (2) and the resulting iterative method. Moreover, we consider this AMLI approach as an additive Schwarz approach.

For some problems it is known that the multiplicative Schwarz method converges faster than the additive Schwarz method [17, 26]. So it is natural to consider a related multiplicative approach. The multiplicative version, which we call the MAMLI method or MAMLI approach [24], is then given by:

$$\begin{aligned} T_{MAMLI} &= (I - P_1)(I - P_2) \\ &= (I - \tilde{P}^T \tilde{S}^{-1} \tilde{R}A)(I - \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}A) \\ &= \left( I - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} A \right) \\ &\quad \cdot \left( I - \begin{bmatrix} I_F \\ 0 \end{bmatrix} \tilde{A}_{FF}^{-1} \begin{bmatrix} I_F & 0 \end{bmatrix} A \right). \end{aligned} \quad (7)$$

Closely related to the multiplicative version there are two other methods. The first one is the reverse MAMLI method (RMAMLI method) the other one is the symmetrized MAMLI method (SMAMLI) [24]. These variants are defined in terms of their iteration matrices given by

$$T_{RMAMLI} = (I - P_2)(I - P_1)$$

and

$$T_{SMAMLI} = (I - P_2)(I - P_1)(I - P_2).$$

The multiplicative versions are closely related to certain geometric and algebraic multigrid methods. The factor  $I - P_2$  in (7) can be seen as a relaxation or smoothing step, while  $I - P_1$  in (7) is a coarse grid correction. In particular, the MAMLI method can be viewed as a two-level V(1,0) cycle, the RMAMLI method as a two-level V(0,1), and the SMAMLI method as a two-level V(1,1) cycle with fine-grid only relaxation and special restriction and coarse grid operators. In the AMLI approach, the smoother and the coarse grid correction operator are combined in an additive way. While in the MAMLI approach, this is done in a multiplicative way. Hence, the MAMLI approach is closer to the original multigrid schemes. Nevertheless, general additive multigrid methods have been successfully developed in the last decades [10, 18, 37].

There is also a close relationship between the MAMLI techniques and the AMGr method introduced by MacLachlan, Manteuffel, and McCormick in [21]. AMGr uses fine-grid only relaxation, similar restriction operators, and Schur complements also. In [21] convergence results and bounds for the rate of convergence are established for the AMGr method applied to symmetric positive definite matrices.

There exist a lot of theoretical results for the AMLI method used as a solver as well as used as a preconditioner for the CG method [4, 5, 27, 2, 6, 13, 14]. However, the theoretical results can be applied to symmetric positive definite matrices only. Notay gives in [28] some results for the multilevel approximate block factorization applied to some special non-symmetric M-matrices which arise from a specific discretization of a certain PDE. But so far a general convergence analysis of the AMLI approach for a wide class of non-symmetric matrices is still missing. In this paper we give convergence results for the above mentioned AMLI approach applied to arbitrary non-symmetric M-matrices.

Moreover, we will establish a detailed convergence theory for the MAMLI, RMAMLI and SMAMLI methods applied to non-symmetric M-matrices.

By a recursive use of the above described two-level techniques one can easily construct the corresponding multilevel methods. For these multilevel methods convergence results are also established in this paper.

Of course, the choice of the coarsing algorithm used to get the partitioning in (1) has a major influence on the convergence behavior. But in this paper we assume that this partitioning is done already in an arbitrary way. We want to focus on the convergence behavior for general partitionings and will compare different multilevel methods starting with the same partitioning. Our convergence results are independent of the coarsing technique that is used.

Recently, additive and multiplicative Schwarz methods for non-symmetric matrices were analyzed in [16, 8, 26]. In these papers an algebraic convergence theory for the additive and multiplicative Schwarz was introduced. This theory yields a lot convergence results

and comparison results. However, this theory includes only special restriction and prolongation operators which are used in domain decomposition methods. This theory can not be applied to multilevel or multigrid methods. In contrast to [16, 8, 26] we use and analyze more general restriction and prolongation operators, here.

The paper is organized as follows. In the next section we give some notation and well-known results. Section 3 describes the properties of the approximations that we are using in the block factorization. In Section 4 we establish convergence and comparison results for two-level methods, while in Section 5 these results are extended to multilevel methods.

## 2 Notation and well-known results

A matrix  $B$  is nonnegative (positive), denoted  $B \geq 0$  ( $B > 0$ ), if its entries are non-negative (positive). We say that  $B \geq C$  if  $B - C \geq 0$ , and similarly with the strict inequality. These definitions carry over to vectors. A matrix  $A$  is a  $Z$ -matrix if its off-diagonal elements are non-positive. A  $Z$ -matrix  $A$  is called a nonsingular  $M$ -matrix if it is monotone, i.e.,  $A^{-1} \geq 0$ . It follows that if  $A$  and  $B$  are nonsingular  $M$ -matrices and  $A \geq B$ , then  $A^{-1} \leq B^{-1}$  [9, 38]. By  $\rho(B)$  we denote the spectral radius of the matrix  $B$ .

We say  $(M, N)$  is a splitting of  $A$  if  $A = M - N$  and  $M$  is nonsingular. A splitting is regular if  $M^{-1} \geq 0$  and  $N \geq 0$ ; it is weak regular of the first type if  $M^{-1} \geq 0$  and  $M^{-1}N \geq 0$ . [9, 38, 40].

Here, we consider stationary iterative methods to solve  $Ax = b$ . These methods start with a vector  $x^{(0)}$  and build a sequence of vectors  $x^{(i+1)}$  such that

$$x^{(i+1)} = Tx^{(i)} + c \quad \text{for } i = 1, 2, \dots \quad (8)$$

The matrix  $T$  is called iteration matrix. If  $\rho(T) < 1$  then, there exists a unique splitting  $(M, N)$  such that  $T = M^{-1}N$ . This splitting is given by  $M = A(I - T)^{-1}$  and  $N = M - A$ . We say that  $T$  is induced by this splitting  $(M, N)$ .

Related to the partitioning of  $A$  in (1), we will denote by  $I$  the  $n \times n$  identity matrix and with  $I_F$  and  $I_C$  the  $n_F \times n_F$  and  $n_C \times n_C$  the identity matrix, respectively.

For  $A = [a_{i,j}] \in \mathbb{R}^{n \times n}$  we define the matrices  $\text{diag}(A)$ ,  $\text{triu}(A)$  and  $\text{tril}(A) \in \mathbb{R}^{n \times n}$  by

$$\begin{aligned} (\text{diag}(A))_{i,j} &= \begin{cases} a_{i,j} & \text{for } i = j \\ 0 & \text{else} \end{cases} \\ (\text{tril}(A))_{i,j} &= \begin{cases} a_{i,j} & \text{for } i \geq j \\ 0 & \text{else} \end{cases} \\ (\text{triu}(A))_{i,j} &= \begin{cases} a_{i,j} & \text{for } i \leq j \\ 0 & \text{else} \end{cases} \end{aligned}$$

Next we recall the definition of the weighted max-norm. Given a positive vector  $w \in \mathbb{R}^n$ , denoted  $w > 0$ , the weighted max-norm is defined for any  $y \in \mathbb{R}^n$  as  $\|y\|_w = \max_{j=1, \dots, n} |\frac{1}{w_j} y_j|$ . The corresponding matrix norm is defined as  $\|T\|_w = \sup_{\|x\|_w=1} \|Tx\|_w$  and the following lemma holds.

**Lemma 2.1.** *Let  $A \in \mathbb{R}^{n \times n}$ , be nonnegative,  $w \in \mathbb{R}^n$ ,  $w > 0$ , and  $\gamma > 0$  such that*

$$Aw \leq \gamma w. \quad (9)$$

Then,  $\|A\|_w \leq \gamma$ . If the inequality in (9) is strict, then the bound on the norm is also strict. Moreover,

$$\|A\|_w = \max_{i=1..n} \frac{(Aw)_i}{w_i}.$$

**Proof:** see e.g. [15]. □

Most of our estimates hold for all positive vectors  $w$  of the form  $w = A^{-1}e$ , where  $e$  is any positive vector, i.e., for any positive vector  $w$  such that  $Aw$  is positive. In particular this would hold for an  $M$ -matrix  $A$  and  $e = (1, \dots, 1)^T$ , i.e., with  $w = A^{-1}e$  being the row sums of  $A^{-1}$ .

Moreover, we need the following well-known properties of  $M$ -matrices.

**Theorem 2.2.** *Let  $A \in \mathbb{R}^{n \times n}$  be a  $Z$ -matrix.  $A$  is a nonsingular  $M$ -matrix if and only if one of the following conditions holds*

- *There exist two nonsingular monotone matrices  $B_1, B_2$ , such that  $B_1 \leq A \leq B_2$ .*
- *Each principal submatrix of  $A$  is a nonsingular  $M$ -matrix.*

*Proof.* See e.g. [9]. □

**Theorem 2.3.** *Let  $A \in \mathbb{R}^{n \times n}$  be a nonsingular  $M$ -matrix partitioned as*

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}$$

where  $A_{FF} \in \mathbb{R}^{s \times s}$  and  $A_{CC} \in \mathbb{R}^{n-s \times n-s}$  for some  $s \in \{1, \dots, n-1\}$ . Then the Schur complement

$$(A/A_{FF}) = A_{CC} - A_{CF}A_{FF}^{-1}A_{FC}$$

is a nonsingular  $M$ -matrix.

*Proof.* see e.g. [9]. □

**Lemma 2.4.** *Let  $A$  be a nonsingular  $M$ -matrix and let  $(M, N)$  be a weak regular splitting of first type. Then*

$$M^{-1} \leq A^{-1}.$$

*Proof.* see e.g. [40, Theorem 3.2] □

**Lemma 2.5.** *Let  $A$  be a nonsingular  $M$ -matrix. Let  $M$  be a  $Z$ -matrix such that  $M \geq A$ . Then,  $(M, M - A)$  is a regular splitting, and therefore a weak regular splitting, of  $A$ .*

*Proof.* Since,  $M$  is a  $Z$ -matrix and  $M \geq A$ ,  $M$  is an  $M$ -matrix, see [9]. The statement then follows immediately from the definitions of a weak regular and regular splittings. □

### 3 The approximations

Of course the quality of the approximations  $\tilde{A}_{FF}$  of  $A_{FF}$  and  $\tilde{S}$  of  $S$  used in the inexact multilevel block factorizations will be important for the convergence behavior. Here we use the following properties of the approximations to prove convergence of the AMLI approach.

**Assumption 3.1.**

Let  $A$  be a nonsingular (non-symmetric)  $M$ -matrix and let  $A$  be partitioned in the following  $2 \times 2$  block structure

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}.$$

Furthermore, let  $\tilde{A}_{FF}$  and  $\tilde{S}$  be chosen such that the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - (A/\tilde{A}_{FF}))$  are weak regular of first type, i.e.

$$\tilde{A}_{FF}^{-1} \geq 0 \quad \text{and} \quad I_F - \tilde{A}_{FF}^{-1} A_{FF} \geq 0; \quad (10)$$

and

$$\tilde{S}^{-1} \geq 0 \quad \text{and} \quad I_C - \tilde{S}^{-1}(A/\tilde{A}_{FF}) \geq 0. \quad (11)$$

Here  $(A/\tilde{A}_{FF})$  is defined by  $(A/\tilde{A}_{FF}) := A_{CC} - A_{CF}\tilde{A}_{FF}^{-1}A_{FC}$ .

For the multiplicative versions we use a slightly modified set of approximations.

**Assumption 3.2.**

Let  $A$  be a nonsingular (non-symmetric)  $M$ -matrix and let  $A$  be partitioned in the following  $2 \times 2$  block structure

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}.$$

Furthermore, let  $\tilde{A}_{FF}$  and  $\tilde{S}$  be chosen such that the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - \tilde{R}A\tilde{P}^T)$  are weak regular of first type, i.e.

$$\tilde{A}_{FF}^{-1} \geq 0 \quad \text{and} \quad I_F - \tilde{A}_{FF}^{-1} A_{FF} \geq 0; \quad (12)$$

and

$$\tilde{S}^{-1} \geq 0 \quad \text{and} \quad I_C - \tilde{S}^{-1}(\tilde{R}A\tilde{P}^T) \geq 0, \quad (13)$$

where  $\tilde{R}$  and  $\tilde{P}^T$  are given as in (4).

Here we point out that these assumptions or properties we require for the approximations are very weak. We will give examples for these kind of approximations at the end of this section. Note that no special coarsing or no special partitioning of the matrix  $A$  is needed to find these kind of approximations.

If we compare Assumptions 3.1 and 3.2, we see that the only difference is the condition for the approximation  $\tilde{S}$ .

Using the relation

$$\tilde{R}A\tilde{P}^T = A_{CC} - A_{CF} \left( 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \right) A_{FC} \quad (14)$$

we will see in the following proposition, that Assumption 3.2 is weaker than Assumption 3.1.

**Proposition 3.3.** *Assumption 3.1 implies Assumption 3.2. In other words, if Assumption 3.1 holds than Assumption 3.2 is fulfilled also.*

**Proof:**

We only have to prove, that equation (10) together with equation (11) imply equation (13).

Hence, let the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - (A/\tilde{A}_{FF}))$  be weak regular of first type. Since

$$\tilde{R}A\tilde{P}^T = A_{CC} - A_{CF} \left( 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \right) A_{FC} ,$$

we obtain

$$\begin{aligned} & I_C - \tilde{S}^{-1}\tilde{R}A\tilde{P}^T \\ = & I_C - \tilde{S}^{-1} \left( A_{CC} - A_{CF} \left( 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \right) A_{FC} \right) \\ = & I_C - \tilde{S}^{-1} \left( A_{CC} - A_{CF}\tilde{A}_{FF}^{-1}A_{FC} \right) \\ & + \tilde{S}^{-1}A_{CF} \left( \tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \right) A_{FC} \\ = & I_C - \tilde{S}^{-1} \left( A/\tilde{A}_{FF} \right) + \tilde{S}^{-1}A_{CF} \left( I_F - \tilde{A}_{FF}^{-1}A_{FF} \right) \tilde{A}_{FF}^{-1}A_{FC}. \end{aligned}$$

Using the splitting property of  $\tilde{A}_{FF}$  and  $\tilde{S}$  and the sign pattern of the M-matrix  $A$  we obtain

$$I_C - \tilde{S}^{-1}\tilde{R}A\tilde{P}^T \geq 0.$$

Hence, the splitting  $(\tilde{S}, \tilde{S} - \tilde{R}A\tilde{P}^T)$  is weak regular of first type.  $\square$

Note that in both Assumptions there is a coupling between the approximations  $\tilde{A}_{FF}$  and  $\tilde{S}$ , but it is very mild. Indeed, starting with an  $M$ -matrix  $A$ , the approximations given by the Jacobi and the Gauss-Seidel methods and the incomplete LU-factorization are admissible approximations for example. To see this, one has to use Lemma 2.5. Thus admissible choices for  $\tilde{A}_{FF}$  are

$$\begin{aligned} \tilde{A}_{FF} &= \text{diag}(A_{FF}), \\ \tilde{A}_{FF} &= \text{tril}(A_{FF}), \\ \tilde{A}_{FF} &= \text{triu}(A_{FF}), \\ \tilde{A}_{FF} &= \tilde{L}\tilde{U}. \end{aligned}$$

Here,  $\tilde{L}$  and  $\tilde{U}$  are the factors of an incomplete  $LU$  factorization of  $A_{FF}$  [23].



We will show in Theorem 5.2 that

$$\tilde{R}A\tilde{P}^T = \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix}$$

is a nonsingular M-matrix, if  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  is a weak regular splitting, i.e. one of the above choices is used as an approximation of  $A_{FF}$ .

Hence, the following approximations  $\tilde{S}$  fulfill our Assumption 3.2

$$\begin{aligned} \tilde{S} &= \tilde{R}A\tilde{P}^T \\ \tilde{S} &= \text{diag}(\tilde{R}A\tilde{P}^T) \\ \tilde{S} &= \text{tril}(\tilde{R}A\tilde{P}^T) \\ \tilde{S} &= \hat{L}\hat{U} \end{aligned}$$

where  $\hat{L}$  and  $\hat{U}$  are the factors of an incomplete  $LU$  factorization of  $\tilde{R}A\tilde{P}^T$ . Even more, (see Theorem 5.2)  $\tilde{S}$  can be chosen as

$$\begin{aligned} \tilde{S} &= (A/\tilde{A}_{FF}) \\ \tilde{S} &= A_{CC} \\ \tilde{S} &= \text{diag}(A_{CC}) \end{aligned}$$

to fulfill Assumptions 3.1 and 3.2.

Hence, the Assumptions 3.1 and 3.2 allow a wide variety of approximations and all kind of approximations used in practice seems to be included.

Next, let us compare our approximations with those used at the theoretical analysis of approximate multilevel block factorization applied to symmetric positive definite systems. There a frequently used assumption is that

$$\tilde{A}_{FF} - A_{FF} \text{ is symmetric positive semidefinite} \quad (15)$$

see e.g. [4, 2, 29]. But this assumptions can be expressed also with the help of splittings as we did above. Equation (15) implies that the splitting  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  is a P-regular splitting of  $A_{FF}$ . P-regular splittings are introduced by Ortega in [31] see also [25]. A splitting  $A = M - N$  is called P-regular if  $M^T + N = M + M^T - A$  is positive definite. Note that a splitting of a symmetric positive definite matrix  $A$  is P-regular if and only if  $\|I - M^{-1}A\|_A < 1$ , see [42].

## 4 Two-level convergence-results

We start this section with a fundamental proposition which is the main tool in our convergence analysis.

**Proposition 4.1.** *Let  $A \in \mathbb{R}^{n \times n}$  be a nonsingular M-matrix. If*

- $C \in \mathbb{R}^{n \times n}$  is nonnegative,
- $I - CA$  is nonnegative,
- $C$  has no zero row,

then  $C$  is nonsingular and the splitting  $(C^{-1}, C^{-1} - A)$  of  $A$  is weak regular of first type. Moreover,

$$\rho(I - CA) \leq \|I - CA\|_w < 1, \quad (16)$$

where  $w := A^{-1}e$  for an arbitrary positive vector  $e \in \mathbb{R}^n$ .

**Proof:** Let  $e \in \mathbb{R}^n$  be an arbitrary positive vector. Then  $Ce$  is also positive. Since  $A$  is a nonsingular M-matrix the inverse  $A^{-1}$  is nonnegative and has no zero row. So the vector  $w := A^{-1}e$  is as well positive. Using these properties we get

$$0 \leq (I - CA)w = w - CAA^{-1}e = w - Ce < w.$$

Due to Lemma 2.1 this leads to

$$\rho(I - CA) \leq \|I - CA\|_w < 1. \quad (17)$$

Now assume, that  $C$  is singular. Then, there exists a nonzero vector  $z$  with  $Cz = 0$ . Let  $A^{-1}z = y$ , then

$$(I - CA)y = y - Cz = y.$$

But this contradicts (17). Hence,  $C$  is nonsingular. The splitting properties follow then directly from the assumptions.  $\square$

In the following we consider  $T_{AMLI}$  as given in (6).

**Lemma 4.2.** *Let Assumption 3.1 be satisfied, i.e.  $A$  is a nonsingular M-matrix partitioned as in (1) and the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - (A/\tilde{A}_{FF}))$  are weak regular of first type. Then*

$$T_{AMLI} \geq 0.$$

**Proof:** A computation leads to

$$\begin{aligned} T_{AMLI} &= I - \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \\ &\quad - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \\ &= I - \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FF} & \tilde{A}_{FF}^{-1}A_{FC} \\ 0 & 0 \end{bmatrix} \\ &\quad - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} A_{CF}(I_F - \tilde{A}_{FF}^{-1}A_{FF}) & (A/\tilde{A}_{FF}) \end{bmatrix} \\ &= \begin{bmatrix} I_F - \tilde{A}_{FF}^{-1}A_{FF} & 0 \\ 0 & 0 \end{bmatrix} \\ &\quad + \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FC}\tilde{S}^{-1}A_{CF}(I_F - \tilde{A}_{FF}^{-1}A_{FF}) & 0 \\ -\tilde{S}^{-1}A_{CF}(I_F - \tilde{A}_{FF}^{-1}A_{FF}) & 0 \end{bmatrix} \\ &\quad + \begin{bmatrix} 0 & -\tilde{A}_{FF}^{-1}A_{FC}(\tilde{S}^{-1}(A/\tilde{A}_{FF})) \\ 0 & I_C - \tilde{S}^{-1}(A/\tilde{A}_{FF}) \end{bmatrix}. \end{aligned}$$

Using the assumptions on the approximations,  $\tilde{A}_{FF}$  and  $\tilde{S}$  and the sign pattern of the M-matrix  $A$  we get that  $T_{AMLI}$  is nonnegative.  $\square$

Lemma 4.2 provide the non-negativity of the AMLI iteration matrix. This powerful property will be used in the convergence analysis below.

**Theorem 4.3.** *Let Assumption 3.1 be satisfied, i.e.  $A$  is a nonsingular  $M$ -matrix partitioned as in (1) and the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - (A/\tilde{A}_{FF}))$  are weak regular of first type. Then*

$$\rho(T_{AMLI}) \leq \|T_{AMLI}\|_w < 1,$$

where  $w = A^{-1}e$  for an arbitrary positive vector  $e$ . Moreover,  $T_{AMLI}$  is induced by a weak regular splitting of first type, i.e.

$$T_{AMLI} = I - C_{AMLI}A,$$

where  $C_{AMLI}$  is nonsingular and  $(C_{AMLI}^{-1}, C_{AMLI}^{-1} - A)$  is a weak regular splitting of first type of  $A$ .

**Proof:** In order to use Proposition 4.1 we first write  $T_{AMLI}$  as  $T_{AMLI} = I - C_{AMLI}A$ . We then establish that  $I - C_{AMLI}A$  satisfies the assumptions of Proposition 4.1.

By Lemma 4.2,  $T_{AMLI}$  is nonnegative. Thus it suffices to show that  $C_{AMLI}$  is nonnegative and  $C_{AMLI}$  has no zero row.

Since

$$T_{AMLI} = I - \tilde{P}^T \tilde{S}^{-1} \tilde{R}A - \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}A = I - P_1 - P_2,$$

the matrix  $C_{AMLI}$  is given by  $\tilde{P}^T \tilde{S}^{-1} \tilde{R} + \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}$ . With

$$M_S := \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R} = \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} \quad (18)$$

$$M_{CG} := \tilde{P}^T \tilde{S}^{-1} \tilde{R} = \begin{bmatrix} -\tilde{A}_{FF}^{-1} A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF} \tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \quad (19)$$

we obtain

$$T_{AMLI} = I - (M_S + M_{CG})A = I - C_{AMLI}A, \quad (20)$$

where  $C_{AMLI} := M_S + M_{CG}$ .

Using the  $M$ -matrix and splitting properties, we see that both matrices  $M_S$  and  $M_{CG}$  are nonnegative. As a sum of two nonnegative matrices,  $C_{AMLI}$  is also nonnegative.

Since  $\tilde{A}_{FF}^{-1}$  and  $\tilde{S}^{-1}$  are the inverses of nonsingular matrices they do not have zero rows. So the first  $n_F$  rows of

$$M_S = \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix}$$

are not zero rows. Moreover the last  $n_C$  rows of

$$M_{CG} = \begin{bmatrix} \tilde{A}_{FF}^{-1} A_{FC} \tilde{S}^{-1} A_{CF} \tilde{A}_{FF}^{-1} & -\tilde{A}_{FF}^{-1} A_{FC} \tilde{S}^{-1} \\ -\tilde{S}^{-1} A_{CF} \tilde{A}_{FF}^{-1} & \tilde{S}^{-1} \end{bmatrix}$$

are not zero rows. Since  $M_{CG}$  and  $M_S$  are nonnegative and  $C_{AMLI} = M_{CG} + M_S$ , the matrix  $C_{AMLI}$  has no zero row.

With Lemma 4.2 and Proposition 4.1 we obtain

$$\rho(T_{AMLI}) \leq \|T_{AMLI}\|_w < 1.$$

□

Starting with an  $M$ -matrix  $A$  and approximations as in (10) and (11) we proved convergence of the AMLI approach for a wide class of non-symmetric matrices. In the convergence proof the non-negativity of the iteration matrix  $T_{AMLI}$  was the major tool. Next we analyze the MAMLI iteration matrix as given in (7).

Before we consider the product  $(I - P_1)(I - P_2)$  we take a look at each factor separately. It is worth mentioning that not both factors  $(I - P_1)$  and  $(I - P_2)$  are nonnegative in general. This is a major difference to the convergence analysis for special Schwarz methods given in [16, 8, 26]. While factor  $I - P_2$  is nonnegative the other factor  $I - P_1$  need not to be nonnegative (see Example 4.5).

**Proposition 4.4.** *Let  $A$  be a nonsingular  $M$ -matrix partitioned as in (1).*

*If  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  is a weak regular splitting of  $A_{FF}$  of first type, then*

$$\begin{aligned} I - P_2 &\geq 0, \\ \|I - P_2\|_w &= 1, \end{aligned}$$

where  $w = A^{-1}e$  for an arbitrary positive vector  $e$ .

**Proof:**

Using the splitting properties and the  $M$ -matrix sign pattern, we get that

$$\begin{aligned} I - P_2 &= I - \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \\ &= \begin{bmatrix} I_F - \tilde{A}_{FF}^{-1}A_{FF} & -\tilde{A}_{FF}^{-1}A_{FC} \\ 0 & I_C \end{bmatrix} \end{aligned}$$

is nonnegative.

Let  $e$  be an arbitrary positive vector. Since  $A$  is a nonsingular  $M$ -matrix  $w = A^{-1}e$  is also positive. With  $\tilde{A}_{FF}^{-1} \geq 0$  we obtain

$$0 \leq (I - P_2)w = \left( I - \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} A \right) A^{-1}e = w - \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} e \leq w.$$

Using Lemma 2.1 we get  $\|I - P_2\|_w \leq 1$ . But since the last components of  $(I - P_2)w$  and  $w$  are the same, it holds that

$$\|I - P_2\|_w = \sup_{\|x\|_w=1} \|(I - P_2)x\|_w \geq \|(I - P_2)w\|_w = \max_i \frac{((I - P_2)w)_i}{w_i} \geq 1$$

This leads to

$$\|I - P_2\|_w = 1.$$

□

**Example 4.5.** *Consider the matrix*

$$A = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix} \left. \begin{array}{l} \} n_F \\ \} n_C \end{array} \right\}$$

and partition it as indicated. We use the approximations

$$\tilde{A}_{FF} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \quad \text{and} \quad \tilde{S} = \begin{bmatrix} \frac{3}{2} & 0 \\ 0 & 2 \end{bmatrix}.$$

These approximations fulfill (10) and (11). Then  $I - P_1$  is given by

$$I - P_1 = \frac{1}{6} \begin{bmatrix} 6 & 0 & 0 & 0 \\ 1 & 6 & -3 & 2 \\ 2 & 0 & 0 & 4 \\ 0 & 0 & 3 & 0 \end{bmatrix}.$$

Although  $(I - P_1)$  is not nonnegative in general, we are able to establish the non-negativity of the MAMLI iteration matrix.

**Lemma 4.6.** *Let Assumption 3.2 be satisfied, i.e.  $A$  is a nonsingular  $M$ -matrix partitioned as in (1) and the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - \tilde{R}\tilde{A}\tilde{P}^T)$  are weak regular of first type. Then*

$$T_{MAMLI} \geq 0.$$

**Proof:**

A computation leads to

$$\begin{aligned} & T_{MAMLI} \\ = & \left( I - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \right) \\ & \cdot \left( I - \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \right) \\ = & I - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FF} & -\tilde{A}_{FF}^{-1}A_{FC} \\ 0 & 0 \end{bmatrix} \\ & - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}(2I - \tilde{A}_{FF}^{-1}A_{FF}) & \tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \\ & \cdot \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \\ = & I - \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FF} & \tilde{A}_{FF}^{-1}A_{FC} \\ 0 & 0 \end{bmatrix} \\ & - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} A_{CF} (I_F - \tilde{A}_{FF}^{-1}A_{FF})^2 & \tilde{R}\tilde{A}\tilde{P}^T \end{bmatrix} \end{aligned}$$

$$\begin{aligned}
&= I - \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FF} & \tilde{A}_{FF}^{-1}A_{FC} \\ 0 & 0 \end{bmatrix} \\
&\quad - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} A_{CF} (I_F - \tilde{A}_{FF}^{-1}A_{FF})^2 & 0 \\ 0 & \tilde{R}A\tilde{P}^T \end{bmatrix} \\
&\quad - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} 0 & \tilde{R}A\tilde{P}^T \end{bmatrix} \\
&= \begin{bmatrix} I_F - \tilde{A}_{FF}^{-1}A_{FF} & 0 \\ 0 & 0 \end{bmatrix} \\
&\quad + \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FC}\tilde{S}^{-1}A_{CF} (I_F - \tilde{A}_{FF}^{-1}A_{FF})^2 & 0 \\ -\tilde{S}^{-1}A_{CF} (I_F - \tilde{A}_{FF}^{-1}A_{FF})^2 & 0 \end{bmatrix} \\
&\quad + \begin{bmatrix} 0 & -\tilde{A}_{FF}^{-1}A_{FC} (I_C - \tilde{S}^{-1}(\tilde{R}A\tilde{P}^T)) \\ 0 & I_C - \tilde{S}^{-1}(\tilde{R}A\tilde{P}^T) \end{bmatrix}
\end{aligned}$$

Using the assumption on the approximations  $\tilde{A}_{FF}$  and  $\tilde{S}$  and the sign pattern of the M-matrix  $A$  we get that  $T_{MAMLI}$  is nonnegative.  $\square$

Due to Proposition 3.3, Assumption 3.1 also leads to a nonnegative matrix  $T_{MAMLI}$ . Next we prove the convergence of the MAMLI method.

**Theorem 4.7.** *Let Assumption 3.2 be satisfied, i.e.  $A$  is a nonsingular M-matrix partitioned as in (1) and the splittings  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  and  $(\tilde{S}, \tilde{S} - \tilde{R}A\tilde{P}^T)$  are weak regular of first type. Then*

$$\rho(T_{MAMLI}) \leq \|T_{MAMLI}\|_w < 1,$$

where  $w = A^{-1}e$  for an arbitrary positive vector  $e$ . Moreover,  $T_{MAMLI}$  is induced by a weak regular splitting of first type, i.e.

$$T_{MAMLI} = I - C_{MAMLI}A,$$

where  $C_{MAMLI}$  is nonsingular and  $(C_{MAMLI}^{-1}, C_{MAMLI}^{-1} - A)$  is a weak regular splitting of first type of  $A$ .

**Proof:** As in the proof of Theorem 4.3 we will use Proposition 4.1. So we first write  $T_{MAMLI}$  as  $T_{MAMLI} = I - C_{MAMLI}A$ . Then we will establish that  $I - C_{MAMLI}A$  satisfies the assumptions of Proposition 4.1. Since with Lemma 4.6  $T_{MAMLI}$  is nonnegative, it suffices to show that  $C_{MAMLI}$  is nonnegative and  $C_{MAMLI}$  has no zero row. With

$$T_{MAMLI} = (I - P_1)(I - P_2)$$

where

$$P_1 = \tilde{P}^T \tilde{S}^{-1} \tilde{R}A \quad \text{and} \quad P_2 = \hat{R}^T \tilde{A}_{FF}^{-1} \hat{R}A,$$

we easily obtain

$$\begin{aligned}
T_{MAMLI} &= I - P_1 - P_2 + P_1P_2 \\
&= I - (M_{CG} + M_S - M_{CG}A M_S)A \\
&= I - C_{MAMLI}A.
\end{aligned} \tag{21}$$

where

$$C_{MAMLI} := M_{CG} + M_S - M_{CG}A M_S$$

and  $M_S$  and  $M_{CG}$  are as in (18) and (19).

Next we show that  $C_{MAMLI}$  is a nonnegative matrix. As seen in the proof of Theorem 4.3 both matrices  $M_S$  and  $M_{CG}$  are nonnegative. For  $-M_{CG}A M_S$  we obtain

$$\begin{aligned} & -M_{CG}A M_S \\ = & - \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I_C \end{bmatrix} \tilde{S}^{-1} \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I_C \end{bmatrix} \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix} \begin{bmatrix} \tilde{A}_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} \\ = & \begin{bmatrix} \tilde{A}_{FF}^{-1}A_{FC}\tilde{S}^{-1}A_{CF} \left( I_F - \tilde{A}_{FF}^{-1}A_{FF} \right) \tilde{A}_{FF}^{-1} & 0 \\ -\tilde{S}^{-1}A_{CF} \left( I_F - \tilde{A}_{FF}^{-1}A_{FF} \right) \tilde{A}_{FF}^{-1} & 0 \end{bmatrix}. \end{aligned} \quad (22)$$

Using the M-matrix and splitting properties we obtain that  $-M_{CG}A M_S$  is nonnegative. Consequently,

$$C_{MAMLI} = M_{CG} + M_S + (-M_{CG}A M_S) \quad (23)$$

is also nonnegative.

Next, we prove that  $C_{MAMLI}$  has no zero row. We already established that all three terms of  $C_{MAMLI}$  in (23) are nonnegative. So it suffices to prove that the term  $M_{CG} + M_S$  has no zero row, but this was already done in the proof of Theorem 4.3. Now using Proposition 4.1, we get that

$$\rho(T_{MAMLI}) \leq \|T_{MAMLI}\|_w < 1.$$

□

Starting with an M-matrix  $A$  and approximations as in (10) and (11) or (12) and (13), respectively, we proved convergence of the AMLI and the MAMLI method. As far as we know these are the first convergence results for these methods for a wide class of non-symmetric matrices.

Next we consider the RMAMLI iteration matrix as given in (8)

$$T_{RMAMLI} = (I - P_2)(I - P_1).$$

Since

$$T_{MAMLI} = (I - P_1)(I - P_2) \quad \text{and} \quad T_{RMAMLI} = (I - P_2)(I - P_1),$$

we immediately obtain

$$\rho(T_{RMAMLI}) = \rho(T_{MAMLI}). \quad (24)$$

Thus

**Corollary 4.8.** *If Assumption 3.2 is satisfied, then*

$$\rho(T_{RMAMLI}) < 1.$$

Although the spectral radii of the iteration matrices of the RMAMLI and the MAMLI method are the same, there are significant differences in the structure of the iteration matrices. The iteration matrix of the RMAMLI method is not nonnegative in general as shown in the next example.

**Example 4.9.** Consider the matrix used in Example 4.5

$$A = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}. \quad \left. \begin{array}{l} \phantom{A =} \\ \phantom{A =} \\ \phantom{A =} \\ \phantom{A =} \end{array} \right\} \begin{array}{l} n_F \\ n_C \\ n_C \end{array}$$

Again we use the approximations

$$\tilde{A}_{FF} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \quad \text{and} \quad \tilde{S} = \begin{bmatrix} \frac{3}{2} & 0 \\ 0 & 2 \end{bmatrix}.$$

We obtain

$$T_{RMAMLI} = \frac{1}{12} \begin{bmatrix} 1 & 6 & -3 & 2 \\ 8 & 0 & 0 & 4 \\ 4 & 0 & 0 & 8 \\ 0 & 0 & 6 & 0 \end{bmatrix}.$$

Now we will analyze the SMAMLI iteration matrix as given in (8)

$$T_{SMAMLI} = (I - P_2)(I - P_1)(I - P_2).$$

**Theorem 4.10.** Let Assumption 3.2 be satisfied. Then

$$\begin{aligned} T_{SMAMLI} &\geq 0, \\ \rho(T_{SMAMLI}) &\leq \|T_{SMAMLI}\|_w < 1, \end{aligned}$$

where  $w = A^{-1}e$  for an arbitrary positive vector  $e$ .

**Proof:** We easily obtain with Lemma 4.6 and Proposition 4.4

$$T_{SMAMLI} = (I - P_2)(I - P_1)(I - P_2) = (I - P_2)T_{MAMLI} \geq 0. \quad (25)$$

With Theorem 4.7 and Proposition 4.4 we get

$$\rho(T_{SMAMLI}) \leq \|T_{SMAMLI}\|_w \leq \|I - P_2\|_w \|T_{MAMLI}\|_w < 1. \quad (26)$$

□

In the remaining of this section we compare the SMAMLI, MAMLI, and AMLI methods with respect to the weighted maximum norm of their iteration matrices.

**Theorem 4.11.** Let Assumption 3.1 be satisfied. Then

$$\|T_{SMAMLI}\|_w \leq \|T_{MAMLI}\|_w \leq \|T_{AMLI}\|_w < 1,$$

where  $w = A^{-1}e$  for an arbitrary positive vector  $e$ .

**Proof:**

The inequality  $\|T_{AMLI}\|_w < 1$  was proved in Theorem 4.3. As shown in (20) and (21) we have

$$\begin{aligned} T_{AMLI} &= I - (M_S + M_{CG})A \\ T_{MAMLI} &= I - (M_S + M_{CG} - M_{CG}A M_S)A \\ &= T_{AMLI} + M_{CG}A M_S A, \end{aligned}$$



where  $M_S$  and  $M_{CG}$  are defined in (18) and (19).

Due to the Lemmata 4.2 and 4.6, both iteration matrices are nonnegative. Since  $A$  is a M-matrix and  $e$  is a positive vector,  $w = A^{-1}e$  is also positive. So both terms  $T_{AMLI}w$  and  $T_{MAMLI}w$  are positive.

As seen in (22) the term  $-M_{CG}AM_S$  is nonnegative. So we obtain that the vector  $M_{CG}AM_S e$  has to be non-positive. Therefore,

$$\begin{aligned}
\|T_{MAMLI}\|_w &= \max_{i=1\dots n} \frac{(T_{MAMLI}w)_i}{w_i} \\
&= \max_{i=1\dots n} \frac{(T_{AMLI}w + M_{CG}AM_S AA^{-1}e)_i}{w_i} \\
&\leq \max_{i=1\dots n} \frac{(T_{AMLI}w)_i}{w_i} + \min_{i=1\dots n} \frac{(M_{CG}AM_S AA^{-1}e)_i}{w_i} \\
&= \|T_{AMLI}\|_w + \min_{i=1\dots n} \frac{(M_{CG}AM_S e)_i}{w_i} \\
&\leq \|T_{AMLI}\|_w.
\end{aligned}$$

The remaining inequality  $\|T_{SMAMLI}\|_w \leq \|T_{MAMLI}\|_w$  follows directly from inequality (26).  $\square$

In the following we establish a result for the AMLI, MAMLI, RMAMLI and SMAMLI methods. All these methods coincide if the block  $A_{FF}$  is inverted exactly, i.e.  $\tilde{A}_{FF} = A_{FF}$ , independent of the quality of the approximation of the Schur complement  $S$ . This results holds for all system matrices  $A$ , symmetric or non-symmetric, and not for M-matrices only.

**Theorem 4.12.** *Let  $A$  be a nonsingular matrix, which is partitioned in the following  $2 \times 2$  block structure*

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}.$$

*If  $A_{FF}$  is nonsingular and  $\tilde{A}_{FF} = A_{FF}$  then*

$$T_{AMLI} = T_{MAMLI} = T_{RMAMLI} = T_{SMAMLI}.$$

*holds for any approximation  $\tilde{S}$  of  $(A/A_{FF})$ .*

**Proof:** We have with (20) and (21)

$$\begin{aligned}
T_{AMLI} &= I - (M_S + M_{CG})A \\
T_{MAMLI} &= I - (M_S + M_{CG} - M_{CG}AM_S)A.
\end{aligned}$$

Similarly, we obtain for the RMAMLI and the SMAMLI method

$$\begin{aligned}
T_{RMAMLI} &= I - (M_S + M_{CG} - M_SAM_{CG})A \\
T_{SMAMLI} &= I - (2M_S + M_{CG} - M_SAM_{CG} \\
&\quad - M_SAM_S - M_{CG}AM_S \\
&\quad + M_SAM_{CG}AM_S)A.
\end{aligned}$$

But a computation leads to

$$M_{CG}AM_S = M_SAM_{CG} = 0,$$

and

$$M_SAM_S = \begin{bmatrix} A_{FF}^{-1} & 0 \\ 0 & 0 \end{bmatrix} = M_S.$$

Hence,

$$T_{AMLI} = T_{MAMLI} = T_{RMAMLI} = T_{SMAMLI}.$$

□

## 5 Multilevel convergence results

In Section 4 we proved convergence for two-level methods. Here we extend the convergence results to multilevel methods, i.e. the two-level AMLI or MAMLI technique is used recursively on different levels or different coarse grids. The so called coarse grid system, the system that involves the Schur complement, is then solved or approximated by the AMLI or MAMLI approach again.

To prove convergence for the multilevel methods, we need the following preliminary results.

**Lemma 5.1.** *Let  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  be a weak regular splitting of first type of  $A_{FF}$ . Then*

$$\tilde{B}_{FF}^{-1} := 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1}$$

*is nonsingular and the splitting*

$$(\tilde{B}_{FF}, \tilde{B}_{FF} - A_{FF})$$

*of  $A_{FF}$  is weak regular of first type.*

*Proof.* We have

$$\begin{aligned} \tilde{B}_{FF}^{-1} &= 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \\ &= \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} + \underbrace{(I - \tilde{A}_{FF}^{-1}A_{FF})}_{\geq 0} \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} \\ &\geq 0. \end{aligned}$$

Thus,  $\tilde{B}_{FF}^{-1}$  can be written as a sum of nonnegative terms. Since  $\tilde{A}_{FF}^{-1}$  is nonsingular,  $\tilde{A}_{FF}^{-1}$  has no zero row. Hence,  $\tilde{B}_{FF}^{-1}$  has also no zero row.

Moreover,

$$\begin{aligned} I - \tilde{B}_{FF}^{-1}A_{FF} &= I - (2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1})A_{FF} \\ &= \underbrace{(I - \tilde{A}_{FF}^{-1}A_{FF})}_{\geq 0} \underbrace{(I - \tilde{A}_{FF}^{-1}A_{FF})}_{\geq 0} \\ &\geq 0. \end{aligned}$$

Thus, we have shown that

- $\tilde{B}_{FF}^{-1}$  is nonnegative,
- $I - \tilde{B}_{FF}^{-1}A_{FF}$  is nonnegative,
- $\tilde{B}_{FF}^{-1}$  has no zero row.

Hence, with Proposition 4.1 the matrix  $\tilde{B}_{FF}^{-1}$  is nonsingular and  $(\tilde{B}_{FF}, \tilde{B}_{FF} - A_{FF})$  is a weak regular splitting of first type.  $\square$

**Theorem 5.2.** *Let  $A$  be a nonsingular  $M$ -matrix partitioned as*

$$A = \begin{bmatrix} A_{FF} & A_{FC} \\ A_{CF} & A_{CC} \end{bmatrix}.$$

Let  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  be a weak regular splitting of first type  $A_{FF}$ . Then

$$\begin{aligned} & A_{CC}, \\ & (A/\tilde{A}_{FF}), \\ \tilde{R}A\tilde{P}^T &= \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \end{aligned}$$

are nonsingular  $M$ -matrices. Moreover,

$$A_{CC} \geq (A/\tilde{A}_{FF}) \geq \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \geq (A/A_{FF}). \quad (27)$$

*Proof.* With Theorem 2.2 we have that  $A_{CC}$  is a nonsingular  $M$ -matrix.

Next we consider  $(A/\tilde{A}_{FF})$ .

Since  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  is a weak regular splitting we obtain with Lemma 2.4 that

$$0 \leq \tilde{A}_{FF}^{-1} \leq A_{FF}^{-1}.$$

Using the  $M$ -matrix structure of  $A$  we get

$$\begin{aligned} (A/A_{FF}) &= A_{CC} - \underbrace{A_{CF}}_{\leq 0} \underbrace{A_{FF}^{-1}}_{\geq \tilde{A}_{FF}^{-1}} \underbrace{A_{FC}}_{\leq 0} \\ &\leq A_{CC} - A_{CF}^{-1} \tilde{A}_{FF}^{-1} A_{CF} = (A/\tilde{A}_{FF}). \end{aligned}$$

Similarly,

$$(A/\tilde{A}_{FF}) = A_{CC} - \underbrace{A_{CF}}_{\leq 0} \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} \underbrace{A_{FC}}_{\leq 0} \leq A_{CC}.$$

Thus, we get the inequalities

$$A_{CC} \geq (A/\tilde{A}_{FF}) \geq (A/A_{FF}). \quad (28)$$

By Theorem 2.2  $A_{CC}$  is a nonsingular  $M$ -matrix and with Theorem 2.3  $(A/A_{FF})$  is a nonsingular  $M$ -matrix also. Hence, by Theorem 2.2, we obtain the desired property of  $(A/\tilde{A}_{FF})$ .

Next consider  $\begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix}$ .

We obtain

$$\begin{aligned} & \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \\ &= A_{CC} - A_{CF} \left( \tilde{A}_{FF}^{-1} + \tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} \right) A_{FC} \\ &= A_{CC} - A_{CF}\tilde{B}_{FF}^{-1}A_{FC} \\ &= \left( A/\tilde{B}_{FF} \right), \end{aligned}$$

where  $\tilde{B}_{FF}^{-1} := \tilde{A}_{FF}^{-1} + \tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1}$ .

With Lemma 5.1,  $\left( \tilde{B}_{FF}, \tilde{B}_{FF} - A_{FF} \right)$  is a weak regular splitting of first type. Hence, replacing  $\tilde{A}_{FF}$  by  $\tilde{B}_{FF}$  we can follow the above part of the proof. Therefore, we obtain that  $\left( A/\tilde{B}_{FF} \right) = \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix}$  is a nonsingular  $M$ -matrix.

Thus, we get the inequalities

$$A_{CC} \geq \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \geq (A/A_{FF}). \quad (29)$$

Next we will prove (27). We have

$$\tilde{B}_{FF}^{-1} = 2\tilde{A}_{FF}^{-1} - \tilde{A}_{FF}^{-1}A_{FF}\tilde{A}_{FF}^{-1} = \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} + \underbrace{\left( I - \tilde{A}_{FF}^{-1}A_{FF} \right)}_{\geq 0} \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} \geq \tilde{A}_{FF}^{-1}.$$

Since  $A$  is an  $M$ -matrix, the blocks  $A_{CF}$  and  $A_{FC}$  are non-positive. Therefore

$$\begin{aligned} & \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \\ &= A_{CC} - \underbrace{A_{CF}}_{\leq 0} \underbrace{\tilde{B}_{FF}^{-1}}_{\geq \tilde{A}_{FF}^{-1} \geq 0} \underbrace{A_{FC}}_{\leq 0} \\ &\leq A_{CC} - \underbrace{A_{CF}}_{\leq 0} \underbrace{\tilde{A}_{FF}^{-1}}_{\geq 0} \underbrace{A_{FC}}_{\leq 0} = \left( A/\tilde{A}_{FF} \right). \end{aligned}$$

Together with (28) and (29), we then obtain (27)

$$A_{CC} \geq \left( A/\tilde{A}_{FF} \right) \geq \begin{bmatrix} -A_{CF}\tilde{A}_{FF}^{-1} & I \end{bmatrix} A \begin{bmatrix} -\tilde{A}_{FF}^{-1}A_{FC} \\ I \end{bmatrix} \geq (A/A_{FF}).$$

□

Now, we will turn to the multilevel methods. As mentioned in the beginning of this section, the coarse grid system is solved recursively with an iterative method, namely the same method as used for the original system. To describe the multilevel methods, we need also a hierarchy of matrices  $A^{(1)}, \dots, A^{(L)}$ , where  $A^{(1)} = A$  and  $A^{(2)}$  is a  $n_c \times n_c$  matrix, etc.

First we consider the AMLI approach. Similarly to Assumption 3.1 in the two-level case, we need assumptions for the approximations in the multilevel case.

**Assumption 5.3.** Let  $A^{(1)} \in \mathbb{R}^{n \times n}$  be a nonsingular  $M$ -matrix. For  $l = 1, \dots, L - 1$  assume that

- $A^{(l)}$  is partitioned in the  $2 \times 2$  block structure

$$A^{(l)} = \begin{bmatrix} A_{FF}^{(l)} & A_{FC}^{(l)} \\ A_{CF}^{(l)} & A_{CC}^{(l)} \end{bmatrix},$$

- $(\tilde{A}_{FF}^{(l)}, \tilde{A}_{FF}^{(l)} - A_{FF}^{(l)})$  is a weak regular splitting of first type,
- $A^{(l+1)}$  is given by

$$A^{(l+1)} = \left( A / \tilde{A}_{FF}^{(l)} \right). \quad (30)$$

Moreover  $(\tilde{S}^{(L)}, \tilde{S}^{(L)} - A^{(L)})$  is a weak regular splitting of first type.

Note that for  $L = 2$ , Assumption 5.3 is nothing else than Assumption 3.1. We immediately obtain the following Lemma.

**Lemma 5.4.** If Assumption 5.3 holds, all matrices  $A^{(l)}, l = 1, \dots, L$  are nonsingular  $M$ -matrices.

*Proof.* The lemma can be proved by induction and Theorem 5.2.  $\square$

The multilevel AMLI iteration matrix  $T_{AMLI}^M$  is then given by

$$T_{AMLI}^M = I - C_{AMLI}^{(1)} A^{(1)}$$

where for  $l = 1, \dots, L - 1$

$$C_{AMLI}^{(l)} := \begin{bmatrix} \tilde{A}_{FF}^{(l)-1} & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} -\tilde{A}_{FF}^{(l)-1} A_{FC}^{(l)} \\ I_C \end{bmatrix} C_{AMLI}^{(l+1)} \begin{bmatrix} -A_{CF}^{(l)} \tilde{A}_{FF}^{(l)-1} & I_C \end{bmatrix}$$

and  $C_{AMLI}^{(L)} := S^{(L)-1}$ .  
We obtain

**Theorem 5.5.** Let Assumption 5.3 be satisfied. Then

$$\rho(T_{AMLI}^M) \leq \|I - C_{AMLI}^{(1)} A^{(1)}\|_w < 1.$$

*Proof.* For level  $L - 1$  we have

$$C_{AMLI}^{(L-1)} := \begin{bmatrix} \tilde{A}_{FF}^{(L-1)-1} & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} -\tilde{A}_{FF}^{(L-1)-1} A_{FC}^{(L-1)} \\ I_C \end{bmatrix} C_{AMLI}^{(L)} \begin{bmatrix} -A_{CF}^{(L-1)} \tilde{A}_{FF}^{(L-1)-1} & I_C \end{bmatrix}.$$

Since  $C_{AMLI}^{(L)} := \tilde{S}^{(L)-1}$ , we can apply Theorem 4.3. Thus  $(\tilde{C}_{AMLI}^{(L)-1}, \tilde{C}_{AMLI}^{(L)-1} - A^{(L)})$  is a weak regular splitting of first type, and since  $A^{(L-1)}$  is an  $M$ -matrix (see Lemma 5.4) the assumptions of Theorem 4.3 are satisfied.

Thus, by Theorem 4.3,  $C_{AMLI}^{(L-1)}$  is nonsingular and  $(C_{AMLI}^{(L-1)^{-1}}, C_{AMLI}^{(L-1)^{-1}} - A^{(L-1)})$  is a weak regular splitting of first type.

Repeating this procedure for level  $L - 2$  and all other levels we obtain

$$\rho(T_{AMLI}^M) \leq \|I - C_{AMLI}^{(1)} A^{(1)}\|_w < 1.$$

□

Next we consider the MAMLI method. Similarly to Section 4, we can use a weaker assumption for the approximations used in the MAMLI method.

**Assumption 5.6.** *Let  $A^{(1)} \in \mathbb{R}^{n \times n}$  be a nonsingular  $M$ -matrix. For  $l = 1, \dots, L - 1$  assume that*

- $A^{(l)}$  is partitioned in the  $2 \times 2$  block structure

$$A^{(l)} = \begin{bmatrix} A_{FF}^{(l)} & A_{FC}^{(l)} \\ A_{CF}^{(l)} & A_{CC}^{(l)} \end{bmatrix},$$

- $(\tilde{A}_{FF}^{(l)}, \tilde{A}_{FF}^{(l)} - A_{FF}^{(l)})$  is a weak regular splitting of first type,
- $A^{(l+1)}$  is given by

$$A^{(l+1)} = \begin{bmatrix} -A_{CF}^{(l)} \tilde{A}_{FF}^{(l)-1} & I \end{bmatrix} A^{(l)} \begin{bmatrix} -\tilde{A}_{FF}^{(l)-1} A_{FC}^{(l)} \\ I \end{bmatrix}. \quad (31)$$

Moreover,  $(\tilde{S}^{(L)}, \tilde{S}^{(L)} - A^{(L)})$  is a weak regular splitting of first type.

Again, note that for  $L = 2$  Assumption 5.6 is nothing else than Assumption 3.2. We immediately obtain again

**Lemma 5.7.** *If Assumption 5.6 holds, all matrices  $A^{(l)}, l = 1, \dots, L$  are nonsingular  $M$ -matrices.*

*Proof.* The lemma can be proved by induction and Theorem 5.2. □

The multilevel MAMLI method is then given by the iteration matrix

$$T_{MAMLI}^M = I - C_{MAMLI}^{(1)} A^{(1)},$$

where for  $l = 1, \dots, L - 1$

$$C_{MAMLI}^{(l)} := \begin{bmatrix} \tilde{A}_{FF}^{(l)-1} & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} -\tilde{A}_{FF}^{(l)-1} A_{FC}^{(l)} \\ I_C \end{bmatrix} C_{MAMLI}^{(l+1)} \begin{bmatrix} -A_{CF}^{(l)} \tilde{A}_{FF}^{(l)-1} & I_C \end{bmatrix} \\ - \begin{bmatrix} -\tilde{A}_{FF}^{(l)-1} A_{FC}^{(l)} \\ I_C \end{bmatrix} C_{MAMLI}^{(l+1)} \begin{bmatrix} -A_{CF}^{(l)} \tilde{A}_{FF}^{(l)-1} & I_C \end{bmatrix} A^{(l)} \begin{bmatrix} \tilde{A}_{FF}^{(l)-1} & 0 \\ 0 & 0 \end{bmatrix}$$

and  $C_{MAMLI}^{(L)} := S^{(L)-1}$ .

We now are able to prove the convergence of the multilevel MAMLI method.

**Theorem 5.8.** *Let Assumption 5.6 be satisfied. Then*

$$\rho(T_{MAMLI}^M) \leq \|I - C_{MAMLI}^{(1)} A^{(1)}\|_w < 1.$$

*Proof.* The proof is similar to the proof of Theorem 5.5. Note, that with Theorem 5.2 all  $A^{(l+1)}$  are M-matrices.  $\square$

In our two level convergence theorems, the assumptions for the approximations of the coarse grid system, i.e. the Schur complement, are expressed with the use of the original matrix  $A$ . In detail, we used  $(A/\tilde{A}_{FF})$  for the AMLI method and  $\tilde{R}A\tilde{P}^T$  for the MAMLI method, see (11) in Assumption 3.1 and (13) in Assumption 3.2.

Thus for the multilevel convergence theorems above, we used the corresponding formulation for building the coarse matrices on each level, i.e.

$$A^{(l+1)} = \left( A/\tilde{A}_{FF}^{(l)} \right)$$

for the AMLI method, and

$$A^{(l+1)} = \begin{bmatrix} -A_{CF}^{(l)} \tilde{A}_{FF}^{(l)-1} & I \end{bmatrix} A^{(l)} \begin{bmatrix} -\tilde{A}_{FF}^{(l)-1} A_{FC}^{(l)} \\ I \end{bmatrix}$$

for the MAMLI method.

However, with similar arguments as in the proofs of the two-level and the multilevel convergence theorems, one can prove the following results.

**Theorem 5.9.** *Let Assumption 5.3 be satisfied, except that (30) is replaced by*

$$A^{(l+1)} = A_{CC}^{(l)}.$$

*Then*

$$\rho(T_{AMLI}^M) \leq \|I - C_{AMLI}^{(1)} A^{(1)}\|_w < 1.$$

*Proof.* Obviously, all matrices  $A^{(l+1)}$  are nonsingular M-matrices.

The proof for the multilevel method is based on the proof for the two-level method. Thus, for the two-level method we assume, that  $\tilde{A}_{FF}$  and  $\tilde{S}$  are chosen such that the splitting  $(\tilde{A}_{FF}, \tilde{A}_{FF} - A_{FF})$  is weak regular of first type and that the splitting

$$(\tilde{S}, \tilde{S} - A_{CC}) \text{ is weak regular of first type.} \quad (32)$$

We omit the level indices here.

Using  $I - \tilde{S}^{-1} A_{CC} \geq 0$  we get

$$\begin{aligned} I - \tilde{S}^{-1}(A/\tilde{A}_{FF}) &= I - \tilde{S}^{-1}(A_{CC} - A_{CF} \tilde{A}_{FF}^{-1} A_{FC}) \\ &= I - \tilde{S}^{-1} A_{CC} + \tilde{S}^{-1}(A_{CF} \tilde{A}_{FF}^{-1} A_{FC}) \\ &\geq 0. \end{aligned}$$

Thus,  $(\tilde{S}, \tilde{S} - (A/\tilde{A}_{FF}))$  is a weak regular splitting. Therefore, the assumptions of Theorems 4.2 and 4.3 are fulfilled and we can follow their proofs. Hence, we obtain that the new iteration matrix  $\hat{T}_{AMLI}$ , that is build by using the approximations (32), is nonnegative. Moreover, we also have  $\rho(\hat{T}_{AMLI}) < 1$  and  $\hat{T}_{AMLI}$  is induced by a weak regular splitting  $(\hat{C}_{AMLI-1}, \hat{C}_{AMLI-1}^{-1} - A)$ .

We can then follow the proof of Theorem 5.5 to establish the convergence of the multilevel method.  $\square$

Moreover, we have for the MAMLI method.

**Theorem 5.10.** *Let Assumption 5.6 be satisfied, except that (31) is replaced by*

$$A^{(l+1)} = A_{CC}^{(l)}$$

or

$$A^{(l+1)} = \left( A / \tilde{A}_{FF}^{(l)} \right).$$

Then

$$\rho(T_{MAMLI}^M) \leq \|I - C_{MAMLI}^{(1)} A^{(1)}\|_w < 1.$$

*Proof.* The proof is similar to the proof of Theorem 5.9. □

## 6 Conclusion

In this paper we have analyzed algebraic multilevel methods applied to non-symmetric M-matrices. We considered two types of AMG methods, the AMLI approach and the MAMLI method. We establish convergence results for these methods used as solvers applied to non-symmetric M-matrices. Moreover, some algebraic properties of these methods and the corresponding iteration matrices and splittings are given.

## Acknowledgement

We would like to thank the referees for helpful comments.

## References

- [1] O. Axelsson. Stabilization of algebraic multilevel iteration methods. *Numerical Algorithms*, 21:23–47, 1999.
- [2] O. Axelsson and M. Neytcheva. Algebraic multilevel iteration method for Stieltjes matrices. *Numer. Linear Algebra Appl.*, 1(3):213–236, 1994.
- [3] O. Axelsson and A. Padiy. On the additive version of the algebraic multilevel iteration method for anisotropic elliptic problems. *SIAM Journal on Scientific Computation*, 20:1807–1830, 1999.
- [4] O. Axelsson and P.S. Vassilevski. Algebraic Multilevel Preconditioning Methods, I. *Numerische Mathematik*, 56:157–177, 1989.
- [5] O. Axelsson and P.S. Vassilevski. Algebraic Multilevel Preconditioning Methods, II. *SIAM Journal Numerical Analysis*, 27:1569–1590, 1990.
- [6] O. Axelsson and P.S. Vassilevski. Variable-step multilevel preconditioning methods, I: self-adjoint and positive definite elliptic problems. *Numerical Linear Algebra and Applications*, 1:75–101, 1994.
- [7] R.E. Bank and C. Wagner. Multilevel ILU decomposition. *Numerische Mathematik*, 82:543–576, 1999.



- [8] M. Benzi, A. Frommer, R. Nabben, and D.B. Szyld. Algebraic theory of multiplicative schwarz methods. *Numerische Mathematik*, 89:605–639, 2001.
- [9] A. Berman and R. J. Plemmons. *Nonnegative Matrices in the Mathematical Sciences*. Academic Press, New York, 1979.
- [10] J.E. Bramble and J.E. Pasciak. New estimates for multilevel algorithms including the  $V$ -cycle. *Math. Comp*, 202:447–471, 1993.
- [11] A. Brandt. General highly accurate algebraic coarsening. *Electron. Trans. Numer. Anal.*, 10:1–20 (electronic), 2000.
- [12] A. Brandt, S. McCormick, and J. Ruge. Algebraic multigrid (AMG) for sparse matrix equations. In *Sparsity and its applications (Loughborough, 1983)*, pages 257–284. Cambridge Univ. Press, Cambridge, 1985.
- [13] R. Falgout and P.S. Vassilevski. On generalizing the AMG framework. *SIAM Journal on Numerical Analysis*, 42:1669–1693, 2004.
- [14] R. Falgout, P.S. Vassilevski, and L. Zikatanov. On two-grid convergence estimates. *Numerical Linear Algebra and Applications*, 12:471–494, 2005.
- [15] A. Frommer, H. Schwandt, and D.B. Szyld. Asynchronous weighted additive Schwarz methods. *Electron. Trans. Numer. Anal.*, 5(June):48–61 (electronic), 1997.
- [16] A. Frommer and D.B. Szyld. An algebraic convergence theory for restricted additive Schwarz methods using weighted max norms. *SIAM Journal on Numerical Analysis*, 39:463–479, 2001.
- [17] M. Griebel and P. Oswald. On the abstract theory of additive and multiplicative Schwarz algorithms. *Numerische Mathematik*, 70:163–180, 1995.
- [18] W. Hackbusch. *Multigrid methods and applications*, volume 4 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 1985.
- [19] W. Hackbusch. *Iterative Solution of Large Sparse Systems of Equations*. Springer Verlag, New York – Berlin – Heidelberg, 1994.
- [20] A. N. Langville and C. D. Meyer. *Google’s PageRank and beyond: the science of search engine rankings*. Princeton University Press, Princeton, NJ, 2006.
- [21] S. MacLachlan, T. Manteuffel, and S. McCormick. Adaptive reduction-based AMG. *Numerical Linear Algebra with Applications*, 13(8):599–620, 2006.
- [22] S. Margenov, L. Xanthis, and L. Zikatanov. On the optimality of the semi-coarsening AMLI algorithm. In *Iterative Methods in Linear Algebra II*, volume 3 of *IMACS Series in Computational and Applied Mathematics*, pages 254–269, New Jersey, USA, 1995.
- [23] J.A. Meijerink and H.A. Van der Vorst. An iterative solution method for linear systems of which the coefficient matrix is a symmetric  $m$ -matrix. *Math. Comp.*, 31:148–162, 1977.
- [24] C. Mense. Algebraisches Schwarz-Verfahren unter Verwendung des Schurkomplements. Master’s thesis, Universität Bielefeld, 2002.

- [25] R. Nabben. A note on comparison theorems for splittings and multisplittings of hermitian positive definite matrices. *Linear Algebra and its Applications*, 233:67–80, 1996.
- [26] R. Nabben. Comparisons between multiplicative and additive schwarz iterations in domain decompositon methods. *Numerische Mathematik*, 95:145–162, 2003.
- [27] M. Neytcheva, O. Axelsson, and K. Georgiev. An application of the AMLI method for solving convection-diffusion problems with potential velocity field. In *Algebraic multilevel iteration methods with applications, Vol. I, II (Nijmegen, 1996)*, pages 197–210. Katholieke Univ. Nijmegen, Dept. Math., Nijmegen, 1996.
- [28] Y. Notay. A robust algebraic multilevel preconditioner for non-symmetric  $M$ -matrices. *Numer. Linear Algebra Appl.*, 7(5):243–267, 2000.
- [29] Y. Notay. Algebraic multigrid and algebraic multilevel methods: a theoretical comparison. *Numerische Linear Algebra with Applications*, 12:419–451, 2005.
- [30] Y. Notay. Aggregation-based algebraic multilevel preconditioning. *SIAM J. Matrix Anal. Appl.*, 27(4):998–1018 (electronic), 2006.
- [31] J. M. Ortega. *Numerical analysis - a seconc course*. Accademic Press, New York, 1972.
- [32] J. W. Ruge and K. Stüben. Algebraic multigrid. In *Multigrid methods*, volume 3 of *Frontiers Appl. Math.*, pages 73–130. SIAM, Philadelphia, PA, 1987.
- [33] Y. Saad and B. Suchomel. ARMS: an algebraic recursive multilevel solver for general sparse linear systems. *Numer. Linear Algebra Appl.*, 9(5):359–378, 2002.
- [34] K. Stüben. Algebraic multigrid (AMG): experiences and comparisons. *Appl. Math. Comput.*, 13(3-4):419–451, 1983.
- [35] K. Stüben. An introduction to algebraic multigrid. In *U. Trottenberg, C. W. Oosterlee and A. Schüller Multigrid*, pages 413–528. Academic Press Inc., 2001.
- [36] A. Toselli and O. Widlund. *Domain decomposition methods—algorithms and theory*, volume 34 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 2005.
- [37] U. Trottenberg, C.W. Oosterlee, and A. Schüler. *Multigrid*. Academic Press, London, 2001.
- [38] R. S. Varga. *Matrix iterative analysis*, volume 27 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, expanded edition, 2000.
- [39] C. Wagner. On the algebraic construction of multilevel transfer operators. *Computing*, 65(1):73–95, 2000.
- [40] Z. I. Woźnicki. Nonnegative splitting theory. *Japan Journal of Industrial and Applied Mathematics*, 11:289–342, 1994.
- [41] J. Xu. Iterative methods by space decomposition and subspace correction. *SIAM Review*, 34:581–613, 1992.
- [42] D. M. Young. *Iterative Solution of Large Linear Systems*. Accademic Press, New York, 1971.