

On the linear response of mechanical systems with constraints

Carsten Hartmann,^{1, a)} Christof Schütte,¹ and Giovanni Ciccotti²

¹⁾*Institut für Mathematik, Freie Universität Berlin, 14195 Berlin, Germany*

²⁾*Dipartimento di Fisica, Università "La Sapienza", 00185 Roma, Italy*

We revisit the problem of the linear response of a constrained mechanical system. In doing so we show that the standard expressions of Green and Kubo carry over to the constrained case without any alteration. The argument is based on the appropriate definition of constrained expectations by means of which Liouville's theorem and the Green-Kubo relations naturally follow.

PACS numbers: 02.70.Ns, 45.10.-b, 64.70.qd

Linear response theory as originally formulated by Green¹ and Kubo^{2,3} provides a tool to describe the response of a mechanical systems to a small external perturbation. The typical derivation of the linear response is based on the Hamiltonian function generating the underlying dynamics⁴. In many instances, however, the equations of motion are only available in a non-Hamiltonian form, although the system is of mechanical origin. One such case is a constrained mechanical system in Cartesian (ambient-space) coordinates with explicit Lagrange multipliers in which case a Hamiltonian formulation is not obvious⁵.

Steps towards a formulation of constrained systems in the framework of non-Hamiltonian statistical mechanics have been taken in Ref. 6. Therein the authors demonstrate that it is still possible to derive a response result that has the familiar form, while involving additional terms that are hard to interpret and which are attributed to an apparent nonzero phase space compressibility.

In this communication we argue that in the presence of constraints the standard linear response result *does* in fact hold — unambiguously and without any alteration from the unconstrained case. Our reasoning is based on an appropriate definition of constrained expectation values that gives rise to the standard Liouville equation for probability densities and Liouville's theorem. The linear response result of Green and Kubo then naturally follows from these ingredients.

I. CONSTRAINED DYNAMICS

Consider a particle of unit mass assuming states $x \in \mathbf{R}^n$ with potential energy $V(x)$. The dynamics are confined to a hypersurface $\mathcal{S} \subset \mathbf{R}^n$ that is defined by

$$\mathcal{S} = \{x \in \mathbf{R}^n : \varphi(x) = 0\}$$

with a scalar function φ and $|\nabla\varphi| \neq 0$ everywhere on \mathcal{S} . Newton's equations for the particle read

$$\begin{aligned} \ddot{x}_t &= -\nabla V(x_t) - \lambda \nabla \varphi(x_t) \\ 0 &= \varphi(x_t) \end{aligned} \quad (1)$$

with initial values $x_0 = x$ and $\dot{x}_0 = v$ satisfying

$$\varphi(x) = 0, \quad v \cdot \nabla \varphi(x) = 0. \quad (2)$$

Here and in the following we use the notation $x \cdot y = x^T y$ and $A : B = \text{tr}(AB)$ to denote the inner products between vectors x and y or the double contraction of second-order tensors A and B .

In (1), we may easily eliminate the Lagrange multiplier λ by differentiating the constraint $\varphi(x_t) = 0$ twice with respect to time. This yields

$$\ddot{x}_t \cdot \nabla \varphi(x_t) + \dot{x}_t \otimes \dot{x}_t : \nabla \nabla \varphi(x_t) = 0$$

where we use the notation $\nabla \nabla = \nabla \otimes \nabla$ to denote the matrix of second derivatives. Hence, with (1),

$$\lambda \nabla \varphi(x) = -(\nabla V(x))^\perp + \mathbb{I}(\dot{x}, \dot{x})$$

with the abbreviations

$$(\nabla V)^\perp = \frac{\nabla \varphi \cdot \nabla V}{|\nabla \varphi|^2} \nabla \varphi, \quad \mathbb{I}(v, v) = \frac{v \otimes v : \nabla \nabla \varphi}{|\nabla \varphi|^2} \nabla \varphi.$$

Inserting the expression for the constraint force into the equations of motion (1) then gives

$$\ddot{x}_t = -(P \nabla V)(x_t) - \mathbb{I}(\dot{x}_t, \dot{x}_t). \quad (3)$$

with the notation $PX = X - X^\perp$. Eq. (3) is called an ambient-space formulation of the differential-algebraic system (1) as it is formulated in terms of the ambient-space coordinates (x, v) on \mathbf{R}^{2n} rather than generalized coordinates, and it does not involve Lagrange multipliers. Introducing the new variable $v = \dot{x}$ it can be recast as

$$\begin{aligned} \dot{x}_t &= v_t \\ \dot{v}_t &= -(P \nabla V)(x_t) - \mathbb{I}(v_t, v_t). \end{aligned} \quad (4)$$

Given initial values satisfying (2) its solution automatically stays on the set

$$\mathcal{M} = \{(x, v) \in \mathbf{R}^{2n} : \varphi(x) = 0, v \cdot \nabla \varphi(x) = 0\}.$$

^{a)}Corresponding author: chartman@mi.fu-berlin.de; supported by the DFG Research Center MATHEON "Mathematics for Key Technologies" (FZT86) in Berlin

II. LIOUVILLE EQUATION

Obviously (4) is not Hamiltonian although it can be shown to be equivalent to a Hamiltonian system (e.g., by using generalized coordinates). Nonetheless its flow shares basic properties of a Hamiltonian system such as being volume-preserving. First of all, note that

$$d\mu(z) = |\nabla\varphi(z)|^2 \delta(\varphi(z))\delta(\dot{\varphi}(z))dz \quad (5)$$

is the natural Liouville measure on \mathcal{M} expressed in terms of the ambient-space coordinates $z = (x, v)$; see Ref. 7 for details. Now, given an observable $f(z)$, we call

$$\langle f \rangle_\rho = \int_{\mathbf{R}^{2n}} f(z)\rho(z, t) d\mu(z)$$

the expectation of f with respect to the (possibly time-dependent) probability density ρ . It is convenient to write (4) as the differential equation

$$\dot{z}_t = B(z_t) \quad (6)$$

with the vector field $B = (v, -P\nabla V - \mathbb{I})^T$ and the solution $z_t = z_t(z)$, $z_0 = z$. The Liouville operator corresponding to (6) then assumes the standard form

$$L = B(z) \cdot \nabla \quad (7)$$

where as before and unless noted otherwise, ∇ denotes the derivative with respect to the argument (here: z).

A. Stationary distribution

Before we can state Liouville's theorem, we have to say what we mean by an invariant distribution. We employ what is known as the *Heisenberg picture* in quantum mechanics and call a distribution ρ invariant under (6) if the expectation of an observable at time t ,

$$f(z_t) = \exp(Lt)f(z),$$

is stationary, i.e., if

$$\langle f(z_t) \rangle_\rho = \langle f(z_0) \rangle_\rho, \quad (8)$$

where the average is understood over the initial values. By construction, f solves the differential equation

$$\begin{aligned} \frac{d}{dt}f(z_t) &= Lf(z_t) \\ f(z_0) &= f, \end{aligned} \quad (9)$$

so stationarity of expectation values is equivalent to say that ρ is a stationary solution of the Liouville equation

$$\begin{aligned} \frac{\partial}{\partial t}\rho(z, t) &= L^*\rho(z, t) \\ \rho(z, 0) &= \rho_0(z). \end{aligned} \quad (10)$$

Here ρ_0 is understood as a probability density with respect to the constrained Liouville measure μ , i.e.,

$$\int_{\mathbf{R}^{2n}} \rho_0 d\mu = 1,$$

and the adjoint L^* is defined by means of the duality relation $(Lf, g) = (f, L^*g)$ with the natural scalar product $(f, g) = \langle fg \rangle_{\rho=1}$ between functions f and g .

The Liouville equation (10) is in fact the usual one, for $(Lf, g) = -(f, Lg)$ or $L^* = -L$, respectively. To see this, it is helpful to note that (cf. Ref. 6, p. 750)

$$\int_{\mathbf{R}^{2n}} (Lf)g |\nabla\varphi|^2 dz = - \int_{\mathbf{R}^{2n}} (Lg)f |\nabla\varphi|^2 dz$$

which can be seen upon expanding the Liouvillian term by term and integrating by parts:

$$\begin{aligned} \int_{\mathbf{R}^{2n}} (Lf)g |\nabla\varphi|^2 dz &= \int_{\mathbf{R}^{2n}} (v \cdot \nabla_x f - (P\nabla V + \mathbb{I}) \cdot \nabla_v f) g |\nabla\varphi|^2 dz \\ &= - \int_{\mathbf{R}^{2n}} \left(v \cdot \nabla_x g - 2 \frac{g}{|\nabla\varphi|^2} v \otimes \nabla\varphi : \nabla\nabla\varphi - g \nabla_v \cdot \mathbb{I} - (P\nabla V + \mathbb{I}) \cdot \nabla_v g \right) f |\nabla\varphi|^2 dz \\ &= - \int_{\mathbf{R}^{2n}} (v \cdot \nabla_x g - (P\nabla V + \mathbb{I}) \cdot \nabla_v g) f |\nabla\varphi|^2 dz \\ &= - \int_{\mathbf{R}^{2n}} (Lg)f |\nabla\varphi|^2 dz \end{aligned}$$

where in the third line we have used that $|\nabla\varphi|^2 \nabla_v \cdot \mathbb{I} = 2v \otimes \nabla\varphi : \nabla\nabla\varphi$.

Hence it follows that

$$\int_{\mathbf{R}^{2n}} (Lf)g d\mu = - \int_{\mathbf{R}^{2n}} (Lg)f d\mu + \int_{\mathbf{R}^{2n}} (L\{\delta(\varphi)\delta(\dot{\varphi})\}) fg |\nabla\varphi|^2 dz$$

with

$$\begin{aligned}
& \int_{\mathbf{R}^{2n}} (L\{\delta(\varphi)\delta(\dot{\varphi})\}) fg|\nabla\varphi|^2 dz \\
&= \int_{\mathbf{R}^{2n}} (v \cdot \nabla_x \{\delta(\varphi)\delta(\dot{\varphi})\} - (P\nabla V + \mathbb{I}) \cdot \nabla_v \{\delta(\varphi)\delta(\dot{\varphi})\}) fg|\nabla\varphi|^2 dz \\
&= \int_{\mathbf{R}^{2n}} (v \cdot \nabla\varphi \delta'(\varphi)\delta(\dot{\varphi}) + v \otimes v : \nabla\nabla\varphi \delta(\varphi)\delta'(\dot{\varphi}) - (P\nabla V + \mathbb{I}) \cdot \nabla\varphi \delta(\varphi)\delta'(\dot{\varphi})) fg|\nabla\varphi|^2 dz \\
&= \int_{\mathbf{R}^{2n}} (v \cdot \nabla\varphi \delta'(\varphi)\delta(\dot{\varphi}) - (P\nabla V) \cdot \nabla\varphi \delta(\varphi)\delta'(\dot{\varphi})) fg|\nabla\varphi|^2 dz.
\end{aligned}$$

Here the second equality follows from $\mathbb{I} \cdot \nabla\varphi = v \otimes v : \nabla\nabla\varphi$. Finally $\dot{\varphi} = v \cdot \nabla\varphi$ with $\dot{\varphi} = 0$ and $(P\nabla V) \cdot \nabla\varphi = 0$, so the last integral vanishes which proves that

$$\int_{\mathbf{R}^{2n}} (Lf)g d\mu = - \int_{\mathbf{R}^{2n}} (Lg)f d\mu \quad (11)$$

or, in other words $L^* = -L$.

B. Liouville's theorem

Now Liouville's theorem readily follows: Let $A \neq \emptyset$ be any compact subset of \mathcal{M} . Then

$$\mu(A) = \int_{\mathbf{R}^{2n}} \chi_A d\mu \quad \text{with} \quad \chi_A(z) = \begin{cases} 1 & \text{if } z \in A \\ 0 & \text{if } z \notin A \end{cases}$$

is the phase space volume of A , and we have to show that $d\mu(A)/dt = 0$ under the flow of (6).

To this end we choose $f = \chi_A$ as initial condition in (9) and take the time derivative of (8) with $\rho = 1$. Since f solves (9), it remains to show that

$$\int_{\mathbf{R}^{2n}} Lf d\mu = 0.$$

But the integral equals $(Lf, \mathbf{1}) = -(f, L\mathbf{1})$ where $L\mathbf{1} = 0$. Hence the integral vanishes and, since A is arbitrary, we have proved conservation of volume.

Before we come to the formulation of the linear response, a final remark is in order: it is a common fallacy that the system (6) were not volume-preserving because its ambient-space divergence is not zero. Indeed,

a vector field B is volume-preserving if and only if it is divergence-free⁸. But here the solutions generated by the vector field B live on the constrained phase space $\mathcal{M} \subset \mathbf{R}^{2n}$, so the appropriate notion of divergence is the divergence on \mathcal{M} , whereas the (unconstrained) divergence in the ambient-space variables does not tell us much about volume-preservation.

III. LINEAR RESPONSE

To (6) we add a small perturbation in the way that

$$\dot{z}_t^\epsilon = B(z_t^\epsilon) + \epsilon G(z_t^\epsilon)u_t, \quad \epsilon \ll 1 \quad (12)$$

where u is a time-dependent scalar forcing (not necessarily smooth) and G is a Hamiltonian vector field compatible with the constraint (in other words, the form of G resembles that of B). Now set

$$L_{\text{eq}} = B(z) \cdot \nabla, \quad L_t = u_t G(z) \cdot \nabla.$$

The Liouville equation associated with (12) then reads

$$\begin{aligned}
\frac{\partial}{\partial t} \rho^\epsilon(z, t) &= (L_{\text{eq}}^* + \epsilon L_t^*) \rho^\epsilon(z, t) \\
\rho^\epsilon(z, 0) &= \rho_0(z).
\end{aligned}$$

with $L_{\text{eq}}^* = -L_{\text{eq}}$ and $L_t^* = -L_t$; the latter follows *mutatis mutandis* from Eq. (11). Applying variation of constants or Dyson's formula yields the formal solution

$$\rho^\epsilon(z, t) = \exp(tL_{\text{eq}}^*)\rho_0(z) + \epsilon \int_0^t \exp((t-s)L_{\text{eq}}^*)L_s^* \rho^\epsilon(z, s) ds$$

for which we seek a perturbative expansion of the form

$$\rho^\epsilon = \rho_0 + \epsilon\rho_1 + \epsilon^2\rho_2 + \dots \quad (13)$$

Let us suppose that the initial distribution $\rho(z, 0) = \rho_0(z)$ is independent of ϵ and is invariant under the unperturbed dynamics, i.e., $L_{\text{eq}}^*\rho_0 = 0$. By plugging the ansatz (13) into the Liouville equation and equating equal powers

ers of ϵ , we recover the $\mathcal{O}(\epsilon)$ -approximation

$$\rho^\epsilon(z, t) \approx \rho_0(z) + \epsilon \int_0^t \exp((t-s)L_{\text{eq}}^*) L_s^* \rho_0(z) ds.$$

Now let

$$H: \mathbf{R}^{2n} \rightarrow \mathbf{R}, \quad H = \frac{1}{2}|v|^2 + V$$

denote the Hamiltonian of the unperturbed system. Clearly, H is preserved for $\epsilon = 0$. The perturbed system (12) for $\epsilon > 0$ obeys the energy balance

$$\frac{d}{dt} H(z_t^\epsilon) = -\epsilon u_t J(z_t^\epsilon) \quad (14)$$

with $z_t^\epsilon = z_t^\epsilon(z)$ denoting the solution of (12) and $J(z) = G(z) \cdot \nabla H(z)$ being the dissipative flux.

A. Green-Kubo relations

We come to our main result that generalizes the classical result of Green¹ and Kubo^{2,3} for Hamiltonian systems to non-Hamiltonian systems of the form (1).

In Eq. (12), suppose that $u_t = \delta(t)$, i.e., the perturbation is impulsive at $t = 0$, and the initial values $z_0^\epsilon = z$ are drawn from the canonical distribution

$$\rho_0(z) \propto \exp(-\beta H(z)), \quad \int_{\mathbf{R}^{2n}} \rho_0(z) d\mu(z) = 1.$$

Let moreover

$$\langle f \rangle_{\rho^\epsilon} = \int_{\mathbf{R}^{2n}} f(z) \rho^\epsilon(z, t) d\mu(z)$$

denote the expectation with respect to the probability distribution ρ^ϵ . By replacing ρ^ϵ in the last equation by its $\mathcal{O}(\epsilon)$ -approximation we find

$$\langle f \rangle_{\rho^\epsilon} \approx \langle f \rangle_{\rho_0} + \epsilon \int_{\mathbf{R}^{2n}} \int_0^t f \exp((t-s)L_{\text{eq}}^*) L_s^* \rho_0 ds d\mu.$$

Using that $u_t = \delta(t)$ in the expression for $L_t^* = -L_t$, the double integral can be recast as

$$\begin{aligned} & \int_{\mathbf{R}^{2n}} \int_0^t f \exp((t-s)L_{\text{eq}}^*) L_s^* \rho_0 ds d\mu \\ &= - \int_{\mathbf{R}^{2n}} \int_0^t f \exp((t-s)L_{\text{eq}}^*) L_s \rho_0 ds d\mu \\ &= - \int_{\mathbf{R}^{2n}} \int_0^t (\exp((t-s)L_{\text{eq}}) f)(G \cdot \nabla \rho_0) \delta(s) ds d\mu \\ &= - \int_{\mathbf{R}^{2n}} (\exp(tL_{\text{eq}}) f)(G \cdot \nabla \rho_0) d\mu \\ &= \beta \int_{\mathbf{R}^{2n}} f(z_t)(G \cdot \nabla H) \rho_0 d\mu \end{aligned}$$

where last equality is due to $\rho_0 \propto \exp(-\beta H)$. Employing the definition of the dissipative flux $J(z) = -G(z) \cdot \nabla H(z)$, it follows that

$$\int_{\mathbf{R}^{2n}} f(G \cdot \nabla H) \rho_0 d\mu = -\langle Jf(z_t) \rangle_{\rho_0}$$

with $z_t = z_t(z)$ denoting the solution to the unperturbed problem. As a consequence we recover the classical linear response result of Green and Kubo, viz.,

$$\langle f \rangle_{\rho^\epsilon} \approx \langle f \rangle_{\rho_0} - \epsilon \beta \langle Jf(z_t) \rangle_{\rho_0}. \quad (15)$$

IV. CONCLUSIONS

In this communication we have demonstrated that the common practice, namely, employing the standard linear response result by Green and Kubo in case of a system with holonomic constraints is indeed justified, although some articles that can be found in the literature suggest otherwise (e.g., Ref. 9). Although there has been no doubt that, in principle, Hamiltonian systems that are subject to holonomic constraints behave like any other natural mechanical system from the viewpoint of statistical mechanics (which is certainly not true for nonholonomic systems), the result closes a loophole in the statistical mechanics of constrained systems involving Lagrange multipliers. Consequently the reader using linear response theory should not worry as to whether his system is subject to constraints or not.

This clearly raises the question whether the result carries over to systems that are either inherently non-Hamiltonian or which involve non-Hamiltonian perturbations. A typical representative of the first class is Nosé-Hoover dynamics while the SLLOD equations of motion belong to the second category. We stress that the ingredients we have employed to derive the Green-Kubo relations are relatively simple and are not at all tied to a Hamiltonian framework: a perturbation consistent with the equations of motion (e.g., divergence-free), a preserved measure that induces an inner product via an expectation and a formal expansion of the evolution equation for the corresponding probability densities. Addressing general dynamical systems with arbitrary non-Hamiltonian perturbations, however, is beyond the scope of this communication.

In the course of the derivation we have also revisited the misleading statement (see, e.g., Refs. 10,11) that flows of constrained systems with non-vanishing ambient-space compressibility do not conserve volume. This, in fact wrong, statement is based on the misconception of taking the ambient-space divergence of a constrained vector field as indicative of being volume-preserving. The reader should also not worry about this issue.

Last but not least all the results in this paper easily generalize to the case of multiple constraints, non-trivial mass matrices or more complicated types of perturbations (clearly, being compatible with the constraints). For the sake of readability we refrain from presenting our results in such generality and leave it to the interested reader to fill this gap.

REFERENCES

- ¹M.S. Green. *J. Chem. Phys.* **22**:398–413, 1954.
- ²R. Kubo, M. Yokota, and S. Nakajima. *J. Phys. Soc. Japan* **12**:570–586, 1957.
- ³R. Kubo, M. Yokota, and S. Nakajima. *J. Phys. Soc. Japan* **12**:1203–1211, 1957.
- ⁴K. Hayashi and S.-I. Sasa. *Physica A* **370**:407–429, 2006.
- ⁵M.E. Tuckerman, Yi Liu, G. Ciccotti, and G.J. Martyna. *J. Chem. Phys.* **115**:1678–1702, 2001.
- ⁶G. Ciccotti, R. Kapral, and A. Sergi. In S. Yip, editor, *Handbook of Materials Modeling*, pages 745–761. Springer, Berlin, 2005.
- ⁷P. Ryckaert and G. Ciccotti. *J. Chem. Phys.* **13**:7368–7374, 1983.
- ⁸L. Simon. *Lectures on Geometric Measure Theory*. Australian National University Centre for Mathematical Analysis, 1984.
- ⁹A. Sergi. *Phys. Rev. E* **67**: 021101, 2003.
- ¹⁰A. Sergi. *Phys. Rev. E* **69**: 021109, 2004.
- ¹¹L.H. de la Pena, R. van Zon, J. Schofield, and S.B Oppsi. *J. Chem. Phys.* **126**: 074105, 2007.